



Some Studies on the Effect of Copper (I) Nicotinate Complex on the Immune Response and Some Biochemical Parameters of *Sea Bass* Fish Vaccinated With *Yersinia Ruckeri* Bacterin

T.T. Saad^{1*}, Z.Kh. El-Meadawy², M.A. ALI³ and A.I.E. Fayza⁴

¹Poultry and Fish Diseases Department, Alexandria University, Alexandria, Egypt

²Pharmacology Department, Faculty of Veterinary Medicine, Alexandria University, Alexandria, Egypt

³Animal Wealth Development Department, Faculty of Veterinary Medicine, Zagazig University, Zagazig, Egypt

⁴Animal Health Research Institute, Alexandria University (Damanhour Branch), Alexandria, Egypt

*Corresponding author's email: talaat_fish@yahoo.com

ABSTRACT

The present study was designed to investigate the influence of dietary supplementation of Copper (I) nicotinate complex at different concentrations (30 and 60 mg/kg feed) for 4 weeks on the immune response and some biochemical parameters of *Sea bass* fish vaccinated with *Yersinia ruckeri* bacterin. The relative level of protection against challenge with virulent strain of *Yersinia ruckeri* was also determined. The present findings indicated that, the using highest concentration of Copper (I) nicotinate (60mg/kg feed) was preferable for stimulation of hemopiosis which reflected on the significant increase of some tested hematological parameters. Both concentrations Copper (I) nicotinate complex had no adverse effect on liver cells which reflected on the unchanged activity of liver enzymes and increase the serum total protein, albumin and globulin. The increased lymphocytes percentage and serum globulin along the course of experiment was elevated phagocytic index and activity gave a strong evidence of the immunostimulant effect of copper (I) nicotinate complex in the levels of cell mediated and humeral immunity. This immunostimulant effects were dose dependant and confirmed by the increase in both antibody titer and relative level of protection against the injected virulent strain of *Yersinia ruckeri*.

Key Words: Copper (I) nicotinate complex; Immune response; Biochemical parameters; *Sea bass* fish.

Abbreviations: ALT (Alanine aminotransferase); AST (Aspartate aminotransferase); ATP (Adenosine triphosphate); GTP (Guanosine triphosphate); Hb (Haemoglobin); NAD (Nicotinamide adenine dinucleotide); NADP (Nicotinamide adenine dinucleotide phosphate); NADPH (reduced form of NADP); PCV (Packed cell volume); RLP (Relative level of protection); SEM (Standard error of mean); TEC (Total erythrocytic count); TLC (Total leucocytic count); UTP (Uridine triphosphate).

INTRODUCTION

Immunostimulants represents an emerging class of chemicals that are designed to amplify immune responses against infectious diseases, pathogens and tumor cells. The field of essential metalloelement complexes has attracted many authors in the last decade. Biochemical mechanisms have been postulated concerning the mode of action of such vitamin metal complexes on enzyme reactivity (Franklin and Richardson, 1980). The same authors reported the medical benefits of the different copper complexes. Al-Mulla Hummadi *et al.* (2005 and 2006) reported that, an organic complex of copper chloride, ascorbic acid and nicotinamide has an immunomodulating effect similar to BCG and a direct antileishmanial effect resulting, partially or entirely from the inhibition of enzymes that are necessary for the parasites carbohydrate metabolism. Another copper complexes had been extensively studied as a potent antitumor, anti-inflammatory and antidiabetic (Greenaway *et al.*, 1999; Li *et al.*, 1999; Tsuji and Sakurai, 1998). Copper (I) nicotinate complex reduced the adverse effects of 5-fluorouracil on patients with hepatocellular carcinoma and enhanced defense mechanisms against oxidative stress (El-Saadani, 2004) and produced antiulcerogenic activity (El-Saadani *et al.*, 1993). Literatures concerning the effect of copper (I) nicotinate complexes in fish are so deficient. Our previous work on Big head carp fish (El-Ashmawy *et al.*, 2007) demonstrated that, fish supplemented with copper (I) nicotinate complex (either 30 or 60 mg/kg feed) showed a significant increase in the percentages of most hematological and biochemical

parameters. Therefore, the present study was planned to investigate the effect of dietary supplementation of different concentration of copper (I) nicotinate complex on the immune response and some biochemical parameters of another fish species (*sea bass*) vaccinated against the same bacterial strain used in the last work (*Yersinia ruckeri* bacteria).

MATERIALS AND METHODS

Copper (I) nicotinate complex

Copper nicotinic acid complex (Fig. 1) is a bright red pure crystalline compound $[\text{CuCl}(\text{HNA})_2]$. The complex has been synthesized by the method of Gohar and Dratovisky (1975) and prepared by Research, Development and Quality Control Division Pharco Pharmaceuticals, Alexandria, Egypt.

Vaccines and Virulent strains

Yersinia ruckeri bacterin was prepared and evaluated according to Badran (1990). Stained *Yersinia ruckeri* bacterin was prepared according to Collins *et al.* (1976) to increase the visibility of the serological reaction.

Fish, Aquaria and Experimental Design

A total number of two hundred and ten apparently healthy *Sea bass* fish with an average body weight of 60 ± 5 gm were obtained from private fish farm at Egypt and used in this study. Fish were acclimated for 2 weeks and fed on commercial fish food containing 25% crude protein. The diet was daily provided at a percentage of 3% of body weight as described by Eurell *et al.* (1979). No drugs were given to the fish along the course of the experiment except those under investigation. Glass aquaria measuring $90 \times 60 \times 50$ cm (10 fish each) were used for holding the fish groups during the experiment. The aquaria were supplied with chlorine free tap water and supplemented with continuous aeration. Throughout the experiment 3 separate subgroups of 10 fish per each treated main group were used. Each subgroup of fish only represents one observation. The present study was divided into two main experiments each of 4 weeks duration. In the first experiment, 90 fish were used and divided into 3 groups (30 fish each). The first group served as a control without drug treatment. The second and the third groups were supplied daily with 30 and 60 mg copper (I) nicotinate complex /kg feed, respectively along the experimental period (4 weeks). However, in the second experiment, the remaining 120 fish were divided into 4 groups (30 fish) each. Fish of the first group were injected intraperitoneally (I/P) with 0.2 ml/fish sterile saline and given feed without medicament and served as non medicated control. The fish in the second group were injected I/P with 0.2 ml/fish of formalin inactivated bacterin from *Yersinia ruckeri* and given feed without medicaments and served as vaccinated control. The fish in the third and fourth groups were similarly inoculated with 0.2 ml/fish of formalin inactivated *Yersinia ruckeri* bacterin and received daily 30 and 60 mg copper (I) nicotinate complex/kg diet, respectively along the experimental period (4 weeks). The booster dose of bacterin was given 2 weeks after the first injection. Afterwards, the same fish were challenged by inoculation of 0.2 ml of virulent strain of *Yersinia ruckeri* of the same strain used for bacterin preparation in the second experiment and fish mortalities were recorded during one week duration period.

Sampling and the analytical methods

In the first experiment, fish were weighed weekly, and at the end of experiment, the body of fish were cleaned and blotted dry with adsorbent paper. Blood samples were collected from the caudal vessels using disposable tuberculin syringe (Hawk *et al.*, 1965) for estimation of total erythrocytic count (TEC), total leucocytic count (TLC), packed cell volume (PCV) Stoskopf (1993). Haemoglobin percentage (Hb %) was assessed according to Drubkin (1947), and differential leucocytic count was determined according to Schalm (1986). Both phagocytic activity and index were also determined according to Hawk *et al.* (1965). Similarly, blood was collected without anticoagulant for serum separation as described by Leid *et al.* (1975). The obtained sera were used for spectrophotometric determination of the activities of AST and ALT as directed by Reitman and Frankel (1957). Serum total protein, albumin, globulin and glucose values were determined spectrophotometrically as implied by the methods of Doumas *et al.* (1981), Reinhold (1953), Coles (1974) and Trinder (1969), respectively.

After end of the second experiment, blood samples were collected and the obtained sera were stored at -20°C until used for detection of immune response against *Yersinia ruckeri* according to the method described by Badran (1990). Briefly, in a standard microtiter plate (U-shaped wells), serial two fold dilutions of serum were made in sterile saline solution using a 0.025 ml pipette dropper and 0.025 ml micro diluter. *Yersinia ruckeri* stained antigen (0.025 ml) was added to the diluted serum. The suspensions were kept in humidified clumber and incubated overnight in the refrigerator dilution still giving a clear agglutination. The negative controls consisted of: A) one drop of sterile physiological saline and one drop of tested serum. B) one drops of sterile physiological saline and one drop of stained antigen. The positive controls were carried out using collected positive antisera. The potency of bacterin were evaluated by calculating the relative level of protection (RLP) by subtracting the value resulted from the division of percent of mortality of vaccinated fish group on percent of mortality of non vaccinated control {RLP = 1 - (percent of mortality of vaccinated fish group/percent of mortality of non vaccinated control group); Newman and Majnarich (1982).

Statistical analysis

75




Fig. 1: Probable Cu^{+} conjugation to ribose (redrawn after McDonald, *et al.*, 1995)

The obtained data were compared between groups within different periods by using student-t test. All data are presented as mean \pm standard error of mean (SEM) by using repeated ANOVA. All tests were performed using computer package of the statistical analysis system (SAS, 1987).

RESULTS AND DISCUSSION

The present findings revealed that, highest concentration of copper (I) nicotinate (60mg/kg feed) only induced a significant increase in the body weight of *Sea bass* throughout the experimental period when compared with the control. However, the average body weight was increased significantly ($p < 0.05$) within the time and reaching its maximum at 4 weeks in all groups (Table 1). The obtained results revealed also that, Total Leucocytic Count (TLC) was significantly increased ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control throughout the experimental period (Table 2). This increase was more pronounced in fish supplemented with highest concentration of copper nicotinate (60 mg/kg diet) than the lowest concentration (30 mg/kg diet) throughout the experimental period (Table 2). However, the Total Erythrocytic Count (TEC) was not significantly changed ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control (Table 2).

Table 1. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on body weight (gm) in *Sea bass* fish.

Treatment	Duration of Treatment			
	1 st week	2 nd week	3 rd week	4 th week
Control	54.2 \pm .80Bd	57.8 \pm 0.58Bc	59.8 \pm .37Bb	59.6 \pm 0.24Ba
30 mg/kg diet	54.4 \pm .36Bd	58.4 \pm 1.03Bc	60.2 \pm 0.86Bb	60.4 \pm 0.24Ba
60 mg/kg diet	57.6 \pm .25Ad	60.2 \pm .97Ac	62.0 \pm 0.45Ab	64.6 \pm 0.68Aa

Capital letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean \pm SEM, n = 30 fish (1st experiment).

In addition, TLC and TEC within the same group were not changed significantly ($p < 0.05$) throughout the experimental period (Table 2). The data summarized in Table 2 indicated that, the percentages of Hb and PCV were significantly increased ($p < 0.05$) only in fish supplemented with highest copper nicotinate concentration (60 mg/kg diet) when compared with the control and lowest copper nicotinate concentration (30 mg/kg diet) which remained comparable throughout the experimental period. There were no significant changes in Hb and PCV values within the same group throughout the experimental period (Table 2).

Table 2. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on TLC (103/mm³), TEC (106/mm³), Hb (g/dl) and PCV (%) in sea bass fish.

Parameters	Period	Treatment		
		Control	30 mg/kg diet	60 mg/kg diet
TLC	1 st week	19.0 \pm 0.58Cc	21.0 \pm 0.33Ba	22.9 \pm 0.58Aa
	2 nd week	20.1 \pm 0.58Cc	20.0 \pm 0.58Ba	24.0 \pm 0.30Aa
	3 rd week	18.9 \pm 0.58Cc	21.7 \pm 0.33Ba	22.8 \pm 0.33Aa
	4 th week	20.0 \pm 0.88Cc	21.9 \pm 0.58Ba	24.0 \pm 0.87Aa
TEC	1 st week	2.0 \pm 0.16Aa	1.9 \pm 0.12Aa	1.9 \pm 0.18Aa
	2 nd week	2.1 \pm 0.03Aa	2.1 \pm 0.09Aa	2.1 \pm 0.03Aa
	3 rd week	2.0 \pm 0.12Aa	2.1 \pm 0.15Aa	2.2 \pm 0.02Aa
	4 th week	2.1 \pm 0.04Aa	1.8 \pm 0.03Aa	2.0 \pm 0.09Aa
Hb	1 st week	7.7 \pm 0.88Ba	8.5 \pm 0.67Ba	10.3 \pm 0.33Aa
	2 nd week	8.9 \pm 0.32Ba	8.6 \pm 0.58Ba	9.1 \pm 0.33Aa
	3 rd week	7.9 \pm 0.32Ba	8.7 \pm 0.88Ba	9.9 \pm 0.86Aa
	4 th week	8.5 \pm 0.58Ba	8.3 \pm 0.57Ba	10.0 \pm 0.33Aa
PCV	1 st week	31.7 \pm 1.19Ba	31.5 \pm 1.73Ba	33.9 \pm 1.15Aa
	2 nd week	30.3 \pm 0.88Ba	32.0 \pm 1.73Ba	35.1 \pm 1.45Aa
	3 rd week	31.7 \pm 0.33Ba	31.9 \pm 1.50Ba	36.3 \pm 1.20Aa
	4 th week	31.7 \pm 0.67Ba	30.3 \pm 1.20Ba	35.9 \pm 0.87Aa

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean \pm SEM, n = 30 fish (1st experiment).

The percentages of lymphocytes and esinophils were significantly increased ($P \leq 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control. However, the percentage of monocytes was not significantly changed in all groups ($p < 0.05$; Table 3). In addition, the percentage of basophils was significantly increased ($p < 0.05$) only in fish supplemented with highest copper nicotinate concentration (60 mg/kg feed) when compared with the control and lowest copper nicotinate concentration (30 mg/kg diet) which remained comparable throughout the experimental period (Table 3). The percentage of neutrophils was significantly decreased ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control. This decrease was more observed in fish

supplemented with highest concentration of copper nicotinate (60 mg/kg) than the lowest used concentration (30 mg/kg) throughout the experimental period (Table 3). There was no significant changes in all types of leucocytes within the same group throughout the experimental period (Table 3).

The values of phagocytic activity and index were significantly increased ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control. This increment was more marked in fish received highest concentration of copper nicotinate (60 mg) than the lowest used concentration (30 mg) throughout the experimental period (Table 4). However, the phagocytic index and activity values within the same group were significantly increased only at the fourth week of the experiment (Table 4).

In the present study, serum total proteins values were increased significantly ($p < 0.05$) in fish received copper (I) nicotinate when compared with the control group (Table 5). However, this increase was more pronounced in fish received highest concentration of copper nicotinate than the lowest used concentration (Table 5). The value of serum albumin of all fish supplemented with copper (I) nicotinate was significantly increased ($p \leq 0.05$) when compared with the control (Table 5).

Table 3. Effect of daily supplementation of copper (I) nicotinate for 4 weeks on the differential leucocytic count (%) in *Sea bass* fish.

Parameters	Period	Treatment		
		Control	30 mg/kg diet	60 mg/kg diet
Lymphocyte	1 st week	49.3 ± 0.33Ba	51.0 ± 0.58Aa	53.1 ± 0.33Aa
	2 nd week	47.3 ± 0.58Ba	52.9 ± 0.58Aa	53.0 ± 0.58Aa
	3 rd week	49.0 ± 0.58Ba	50.1 ± 0.58Aa	51.3 ± 0.33Aa
	4 th week	48.1 ± 0.03Ba	51.2 ± 0.88Aa	52.3 ± 0.33Aa
Monocyte	1 st week	1.3 ± 0.33Aa	1.3 ± 0.33Aa	1.2 ± 0.33Aa
	2 nd week	1.4 ± 0.31Aa	1.3 ± 0.33Aa	1.2 ± 0.88Aa
	3 rd week	1.2 ± 0.57Aa	1.1 ± 0.33Aa	1.1 ± 0.33Aa
	4 th week	1.2 ± 0.58Aa	1.3 ± 0.33Aa	1.3 ± 0.58Aa
Basophil	1 st week	8.1 ± 0.57Ba	8.3 ± 0.88Ba	10.3 ± 0.33Aa
	2 nd week	8.2 ± 0.58Ba	8.3 ± 0.88Ba	10.7 ± 0.33Aa
	3 rd week	8.3 ± 0.32Ba	8.3 ± 0.33Ba	10.3 ± 0.88Aa
	4 th week	8.0 ± 0.88Ba	7.9 ± 0.58Ba	10.3 ± 0.33Aa
Eosinophil	1 st week	6.3 ± 0.31Ba	8.3 ± 0.33Aa	8.1 ± 0.33Aa
	2 nd week	6.5 ± 0.33Ba	8.1 ± 0.33Aa	8.0 ± 0.33Aa
	3 rd week	6.0 ± 0.33Ba	7.7 ± 0.33Aa	8.0 ± 0.58Aa
	4 th week	5.9 ± 0.57Ba	7.8 ± 0.88Aa	8.0 ± 0.58Aa
Neutrophil	1 st week	36.0 ± 0.00Aa	30.1 ± 0.58Ba	26.2 ± 1.15Ca
	2 nd week	34.6 ± 2.33Aa	29.4 ± 1.15Ba	26.1 ± 0.33Ca
	3 rd week	35.5 ± 0.58Aa	30.8 ± 0.33Ba	27.3 ± 0.58Ca
	4 th week	35.8 ± 1.86Aa	30.8 ± 0.88Ba	28.1 ± 0.67Ca

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean ± SEM, $n = 30$ fish (1st experiment).

Table 4. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on phagocytic activity (%) and index in *Sea bass* fish.

Parameters	Period	Treatment		
		Control	30 mg/kg diet	60 mg/kg diet
Phagocytic activity	1 st week	21.0 ± 0.58Cb	21.5 ± 0.33Bb	23.5 ± 0.88Ab
	2 nd week	18.8 ± 0.58Cb	20.0 ± 0.58Bb	26.3 ± 0.33Ab
	3 rd week	19.9 ± 0.58Cb	21.0 ± 0.58Bb	25.1 ± 0.33Ab
	4 th week	20.0 ± 0.58Ca	24.3 ± 0.88Ba	26.3 ± 0.67Aa
Phagocytic index	1 st week	2.0 ± 0.05Cb	2.9 ± 0.06Bb	5.3 ± 0.03Ab
	2 nd week	1.8 ± 0.02Cb	2.6 ± 0.09Bb	4.0 ± 0.12Ab
	3 rd week	1.9 ± 0.07Cb	2.7 ± 0.06Bb	3.8 ± 0.09Ab
	4 th week	1.9 ± 0.22Ca	3.0 ± 0.06Ba	5.4 ± 0.12Aa

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean ± SEM, $n = 30$ fish (1st experiment).

The value of serum globulin of all fish supplemented with copper (I) nicotinate was significantly increased ($p < 0.05$) when compared with the control. This increment was more recorded in fish fed highest concentration of copper nicotinate (60 mg/kg) than the lowest used concentration (30 mg/kg) throughout the experimental period (Table 5). The data presented in Table 6 indicated that the activities of ALT and AST were not changed in all treated group when compared with the control group throughout the whole experimental period. In addition, total protein, Albumin, globulin, ALT and AST activities within the same group were not changed significantly ($p < 0.05$) throughout the experimental period (Table 5, 6). The present findings also indicated that, the values of glucose were significantly ($p < 0.05$) decreased in all fish supplemented with copper (I) nicotinate when compared with the control (Table 6). This decrease was more recorded in the highest concentration of copper nicotinate (60 mg) than the lowest used concentration (30 mg) throughout the experimental period (El-Ashmawy, et al., 2007; Table 6).

The values of antibody titer in fish vaccinated against *Yersinia ruckeri* bacterin was significantly increased ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control. This increment was more marked in the highest concentration of copper nicotinate (60 mg/kg) than the lowest used concentration (30 mg/kg) throughout the experimental period (4 weeks, Table 7). In addition, the values of antibody titer were time dependant reaching its maximum at fourth week of the experiment in all tested fish (Table 7). The percentage of RLP of fish challenged by *Yersinia ruckeri* virulent strain was significantly increased ($p < 0.05$) in all fish supplemented with copper (I) nicotinate when compared with the control. This increment was higher in fish fed highest concentration of copper nicotinate (60 mg/kg diet; 60%) than the lowest used concentration (30mg/kg diet; 40%) throughout the experimental period (one week; Table 8).

Table 5. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on total protein (g/dl), albumin (g/dl) and globulin (g/dl) in *Sea bass* fish.

Parameters	Period	Treatment		
		Control	30 mg/kg diet	60 mg/kg diet
Total proteins	1 st week	4.0 ± 0.06Ca	4.6 ± 0.03Ba	7.0 ± 0.15Aa
	2 nd week	2.0 ± 0.06Ca	5.8 ± 0.03Ba	5.8 ± 0.08Aa
	3 rd week	3.2 ± 0.12Ca	4.8 ± 0.03Ba	4.8 ± 0.09Aa
	4 th week	2.9 ± 0.06Ca	5.7 ± 0.03Ba	5.9 ± 0.09Aa
Albumin	1 st week	2.1 ± 0.03Ca	4.1 ± 0.06Ba	3.9 ± 0.24Aa
	2 nd week	2.2 ± 0.03Ca	3.2 ± 0.16Ba	3.9 ± 0.22Aa
	3 rd week	2.2 ± 0.63Ca	3.2 ± 0.06Ba	4.0 ± 0.10Aa
	4 th week	2.2 ± 0.15Ca	3.0 ± 0.19Ba	5.1 ± 0.06Aa
Globulin	1 st week	0.8 ± 0.03Ca	1.5 ± 0.03Ba	2.1 ± 0.17Aa
	2 nd week	0.7 ± 0.03Ca	1.6 ± 0.17Ba	1.9 ± 0.15Aa
	3 rd week	0.9 ± 0.54Ca	1.6 ± 0.09Ba	1.8 ± 0.15Aa
	4 th week	0.7 ± 0.10Ca	1.7 ± 0.20Ba	1.8 ± 0.03Aa

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean ± SEM, n = 30 fish (1st experiment).

Table 6. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on ALT (U/l), AST (U/l) and glucose (mg/dl) in *Sea bass* fish.

Parameters	Period	Treatment		
		Control	30 mg/kg diet	60 mg/kg diet
ALT	1 st week	67.3 ± 0.33Aa	67.7 ± 1.44Aa	66.7 ± 0.88Aa
	2 nd week	66.1 ± 1.45Aa	65.7 ± 0.32Aa	68.0 ± 0.58Aa
	3 rd week	65.9 ± 0.88Aa	68.1 ± 0.33Aa	66.0 ± 0.56Aa
	4 th week	67.8 ± 2.65Aa	67.3 ± 1.20Aa	67.0 ± 1.15Aa
AST	1 st week	82.7 ± 0.33Aa	82.3 ± 0.33Aa	82.0 ± 0.58Aa
	2 nd week	80.5 ± 1.53Aa	82.0 ± 0.57Aa	82.7 ± 1.45Aa
	3 rd week	81.1 ± 1.20Aa	81.3 ± 0.33Aa	80.0 ± 0.58Aa
	4 th week	80.3 ± 0.33Aa	82.5 ± 0.88Aa	82.0 ± 0.56Aa
Glucose	1 st week	81.1 ± 0.88Aa	80.7 ± 1.85Ba	81.3 ± 0.88Ca
	2 nd week	83.4 ± 1.20Aa	81.3 ± 0.32Ba	80.7 ± 0.34Ca
	3 rd week	81.3 ± 1.76Aa	82.7 ± 0.31Ba	81.3 ± 0.87Ca
	4 th week	82.3 ± 0.33Aa	80.5 ± 0.33Ba	81.0 ± 0.58Ca

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$). Values are expressed as mean ± SEM, n = 30 fish (1st experiment).

Table 7. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on antibody titer in *Sea bass* fish vaccinated with *Yersinia ruckeri* bacterin. Values are expressed as mean ± SEM, n = 30 fish (2nd experiment).

Period	Control (not vaccinated)	Control (vaccinated)	30 mg/kg diet	60 mg/kg diet
1 st week	0.00	4 ± 0.04Cb	5 ± 0.03Bb	6 ± 0.05Ab
2 nd week	0.00	4 ± 0.03Cb	5 ± 0.04Bb	6 ± 0.03Ab
3 rd week	0.00	5 ± 0.03Ca	6 ± 0.05Ba	7 ± 0.02Aa
4 th week	0.00	5 ± 0.03Ca	6 ± 0.03Ba	7 ± 0.04Aa

Capital letter: indicated that means within the same row carrying different letter are significantly differed at ($P < 0.05$). Small letter: indicated that means within the same column carrying different letter are significantly differed at ($P < 0.05$).

Table 8. Effect of daily supplementation of copper (I) nicotinate complex for 4 weeks on protection of *Sea bass* fish against virulent strain of *Yersinia ruckeri* after vaccination with *Yersinia ruckeri* bacterin. n = 30 fish (2nd experiment).

Groups	Total number of tested fish	Number of dead fish	Mortality (%)	RLP (%)
Control (not vaccinated)	30	30	100	1- 100/100 = 0
Control (vaccinated)	30	21	70	1- 70/100 = 30
30 mg/kg diet (vaccinated)	30	18	60	1- 60/100 = 40
60 mg/kg diet (vaccinated)	30	12	40	1- 40/100 = 60

DISCUSSION

The medical benefits particularly the immunostimulant effect of the different copper complexes were reported (Eduardo *et al.*, 2009). Our previous work (El-Ashmawy, *et al.*, 2007) demonstrated the beneficial effect of dietary copper (I) nicotinate complex (either 30 or 60 mg/kg diet) on overall performance of Big head carp fish. Therefore we decided to investigate such effect in Sea bass fish which considered as a highly consumed fish species in Egypt. The observed increase of TLC in fish kept on copper nicotinate complex may be attributed to the activation of lymphoid tissue. Liebert, (2009) reported that, copper was involved in hemoglobin formation by simulation of ferro-oxidase I enzyme which catalyzes the oxidation of ferrous iron and plays a role in the transfer of iron from storage to sites of hemoglobin synthesis. Concerning the present hematological findings, the same findings were observed in our previous study in Big head carp fish using the same concentrations of copper nicotinate complex (Eduardo *et al.*, 2009) except only for the value of monocytes which was increased in big head carp fish. Moreover, the increment of lymphocytes, basophils and eosinophils in fish kept on copper nicotinate complex indicated the direct stimulation of copper nicotinate to lymphoid tissue (El-Ashmawy, *et al.*, 2007). However, the efficiency of either highest or lowest concentrations of copper nicotinate for lymphocyte stimulation was the same while, that for the other cells was differed as the highest concentration only was able to induce such stimulation. Moreover, the highest concentration only able to stimulate the basophils after prolonged period of administration (2 weeks). All leucocytes were calculated as a percentage of the whole leucocytic count which constitute 100 %. The significant decrease of the percentage of neutrophils in copper nicotinate received groups was simply attributed to the significant increase of other leucocytes. Obviously, the increment of phagocytic index and activity in highest concentration copper nicotinate received group than the lowest concentration received one introduced another evidence of the superiority of that concentration for stimulation of phagocytic cells than the lowest concentration. Similar results concerning the phagocytic index and activity were obtained in Big head carp fish when fed the same concentrations of copper (I) nicotinate complex used in the present study, the increase in phagocytic assay mainly due to increase in blood parameter especially in lymphocytes and monocytes (Taoka *et al.*, 2006b). Hematological parameters of fish blood, are useful tools that aids in diagnosis of disease. It can also be used to study immuno-potentiators. Such tests are general but not conclusive and must be correlated with biochemical tests of the subject. The significant increase of total protein, albumin and globulin in fish administered copper nicotinate complex perhaps attributed to the role of copper in protein biosynthesis as it is vitally concerned in the growth process (Minganti *et al.*, 2010). In addition, copper is involved in the formation of disulphide linkage of collagen and elastin proteins (Shalaby *et al.*, 2006). Moreover, the significant increase of serum globulin indicated the immunostimulant effect of copper nicotinate particularly for the highest concentration used. The role of copper in humeral and cell mediated immunity was reported by Radostits *et al.* (2000). The same author and others reported that, copper deficiency caused alteration in humeral response Gatesoupe (2007). In addition, the results concerning the effect of copper nicotinate on the activities of ALT and AST indicated that, both used concentrations of copper nicotinate were safe to the liver that it preserve liver enzymes at normal values. In addition, the significant increase ($p < 0.05$) of total protein, albumin and globulin indicated that, the administered copper nicotinate did not disturb the liver functions. Similar results of liver enzymes and protein patterns were obtained in Big head carp fish fed the same used copper nicotinate complex doses Marzouk *et al.* (2008). In addition, the significant decrease of serum glucose in fish administered copper nicotinate complex indicated the hypoglycemic effect of copper nicotinate particularly for the highest concentration used. This decrease perhaps attributed either to decrease the absorption of glucose from the intestine or the stimulation of hypoglycemic hormone like insulin. This result is in accordance with those obtained by Patra (2011). The results concerning the effect of the administration of copper nicotinate complex on the values of antibody titer in fish vaccinated against *Yersinia ruckeri* bacterin are correlated with those obtained by El-Saadani (2004) and in big head carp kept in the same experimental situation (El-Ashmawy *et al.*, 2007). Furthermore, the RLP of the present vaccinated control fish challenged by *Yersinia ruckeri* virulent strain (30%) are disagree with those obtained by EL-Gamal (2005; 25%) and El-Ashmawy *et al.* (2007; 22%). The confliction between our previous (Salinas *et al.*, 2008) and our present findings perhaps attributed to species difference. The increment of antibody titer of copper nicotinate administered group against *Yersinia ruckeri* bacterin along with the significant increase of the percentage of RLP challenged by *Yersinia ruckeri* virulent strain confirmed the above mentioned immunostimulant effect of copper nicotinate. This immunostimulant effect of copper nicotinate was reported in big head carp fish exposed to the same experimental design (EL-Ashmawy *et al.*, 2007). From the biochemical point of view, it is suggested that most of the absorbed copper (I) nicotinate complex was biologically utilized as such. This suggestion is based up on the structural stability of the complex with regard to competing ligands Nouh *et al.* (2009). Moreover, the most probable modification of the ligand in this copper complex is reduction to the pentadienyl derivatives shown in Figure 1. Hence, this complex could be conjugated in hepatocytes with phosphoribose resembling that of nicotinic acid Eid and Mohamed, (2008). Accordingly, NAD and NADP like structures are suggested to be more reactive as hydrogen carriers.

This is due to the electronegativity of the chloride ion. The ion pair of electrons on nitrogen in turn are attracted toward the copper atom as shown in Figure 1. This electron shift enhances stability of the complex. The predicted higher reactivity of NAD and NADP-like structures enhances the activity of oxido-reductases which catalyze anaerobic and aerobic oxidation that result in accumulation of ATP, GTP and UTP. These nucleotides are essential compounds for biosynthesis and production of immunoglobulin (Al-Dohail *et al.*, 2009).

CONCLUSION

The present study can introduce a powerful evidence of the immunostimulant effect of copper nicotinate in the level of cell mediated and humeral immunity without adverse effect on the liver functions of Sea bass fish. This immunostimulant effect was dose dependent as highest concentration of administered copper nicotinate (60 mg/kg of fish diet) was preferable for the stimulation of humeral and cell mediated immunity in the examined fish than the lowest used concentration (30 mg/kg diet).

REFERENCES

- Al-Dohail, M.A.; Hashim, R. and Aliyu- Paiko, M. (2009) : Effects of the probiotic, Lactobacillus acidophilus, on the growth performance, hematology parameters and immunoglobulin concentration in African Catfish (*Clarias gariepinus*, Burchell 1822) fingerling .Aquaculture Research, 40: 1642-1652.
- Al-Mulla Hummadi, Y.M., AL-Bashir, N.M., Najim, R.A., (2006.) Leishmania major and Leishmania tropica: II. Effect of an immunomodulator, S(2) complex on the enzymes of the parasites. Exp. Parasitol. 112, 85–91.
- AL-Mulla Hummadi, Y.M., Najim, R.A., AL-Bashir, N.M., (2005). Leishmania major and Leishmania tropica: I. The in vitro effects of an immunomodulator, S2-complex. Exp. Parasitol. 111, 47–54.
- Badran, A. F., (1990). The role of adjuvants in the immune response of the fish. Zagazig Veterinary Medicine Journal 18, 126–136.
- Coles, E.H., (1974). Veterinary Clinical Pathology. W.B. Saunders Company, Philadelphia and London.
- Collins, M. T., Dawe, D. L., Gratzek, J. B., (1976). Immuno response of Channel catfish (*Ictalurus punctatus*) under different environmental conditions. J. Am. Vet. Med. Ass. 169, 991–994.
- Doumas, B.T., Bayson, D.D., Carter, R.J., Peters, T., Schaffer, R., (1981). Estimation of total serum protein. Clin. Chem. 27, 1642–1643.
- Drubkin, D., (1947). Spectrophotometric methods XIV. The crystographic and optical properties of the haemoglobin of man in comparison with those of other species. J. Biol. Chem. 164, 703–723.
- Eduardo, G. A.; Gisele, C. F.; Daniela, C.; Fabiana, G. and Jose, C. D. (2009) : Dietary supplementation of lysine and/or methionine on performance, nitrogen retention and excretion in pacu *Piaractus mesopotamicus* reared in cages. Aquaculture. 295: 266-270.
- El-Ashmawy, I.M., Saad , T.T., El-Bahr, S.M., Salama, O.M., (2007). Assessment of immune response of big head carp fed on copper (I) nicotinate and vaccinated with *Yersinia Ruckeri* bacterin. Veterinary Medicine Journal Giza 2, 507–22.
- EL-Gamal, H.L., (2005). Some studies on the effect of *Yersinia Ruckeri* in the cultured fresh water fish. Ph.D., Faculty of Veterinary Medicine, Alexandria University, Egypt.
- EL-Saadani, M.A., (2004). A combination therapy with copper (I) nicotinate complex reduces the adverse effects of 5-fluorouracil on patient with hepatocellular carcinoma. J. Exp. Ther. Oncol. 4, 19–24.
- EL-Saadani, M.A., Nassar, A.Y., Abou El-Ela, S.H., (1993). The protective effect of copper complexes against gastric mucosal ulcer in rats. Biochem. Pharmacol. 46, 1011–1018.
- Eurell, T. E., Lewis, S.D., Grumbles, L.C., (1979). Comparison of selected diagnostic tests for detection of motile *Aeromonas septicaemia* in fish. Am. J. Vet. Res. 39, 1384–1386.
- Franklin, T.A., Richardson, M.F., (1980). Reactions of metal ions and vitamins: III. Synthesis and infrared spectra of metal complexes with pyridoxamine and pyridoxine. Inorganic Chemica Acta 46, 191–197.
- Gatesoupe, F.J. (2007) : Live yeasts in the gut: Natural occurrence, dietary introduction, and their effects on fish health and development. Aquaculture, 267: 20-30.
- Gohar, M., Dratovisky, M., (1975). Perchlorates of Cu (I) complexes with some pyridine derivatives. Coll. Czech. Chem. Commun. 40, 26.
- Greenway, F.T., Riviere, E., Girerd, J.J., (1999). Copper (II) complexes of a non steroidal anti-inflammatory drug niflumic acid. J. Inorg. Biochem. 76, 19–27.
- Haselkorn, R., Rothman-Denes, L., (1973). Protein synthesis. Annual Review of Biochemistry 42, 397–438.
- Hawk, P.B., Oser, B.L., Summerson, W.H., (1965). Practical Physiological Chemistry, 14th ed., McGraw-Hill Book Co. Inc., New York.
- Li, C., Wu, J., Wang, L., (1999). Synthesis, characterization and antitumor activity of copper (II) complex with nicotinamid-4-bis (2-chloroethyl) aminobenzaldimine. J. Inorg. Biochem. 73, 195–202.
- Liebert, F. (2009) : Amino acid requirement studies in *Oreochromis niloticus* by application of principles of the diet dilution technique. J Anim Physiol Anim Nutr (Berl). 93 (6):787-93.
- Lied, E., Gezrde, Z., Braskan, O.R., (1975). Simple and rapid technique for repeated blood sampling in rainbow trout. J. Fish. Res., Board of Canada 32, 699–701.
- Marzouk, M.S.; Moustafa, M.M. and Nermeen, M. M. (2008) : Evaluation of immunomodulatory effects of some probiotics on cultured *Oreochromis niloticus*. 8th International Symposium on Tilapia in Aquaculture.

- McDonald, P., Edward, R.A. and Green balgh, J.F. (1995): Animal nutrition 5th ed. pp. 103-106, English language, Book society
- Minganti, V.; Drava, G.; Pellegrini, R. and Siccardi, C. (2010) : Trace elements in farmed and wild gilthead Seabream, *Sparus aurata*. Marine Pollution Bulletin, (60): 2022 - 2025.
- Newman, S. G., Majnarich, J. J., (1982). Direct immersion vaccination of Juvenile rainbow trout, *Salmo gairdneri* Richardson, and Juvenile coho salmon, *Onchorhynchus kisutch* (Walbaum), with a *Yersinia ruckeri* bacterin. J. Fish Diseases 5, 339–341.
- Nouh, W. G.; Mohamed, M. F. and Aly, S.M. (2009) : Pathological evaluation to the effect of some probiotics on the health and immune status of Nile Tilapia (*Oreochromis niloticus*). J. Comp. Path. And Clinic. Path. 22(2): 233–249.
- Petrack, B., Greengard, P., Craston, A., Kalinsky, H., (1963). Nicotinamide deamidase in rat liver and biosynthesis of NAD. Biochem. Biophys. Res. Commun. 13, 472–477.
- Prohaska, J.R., Failla, M.L., (1993). Copper and Immunity, 1st ed. New York, Plenum Press.
- Radostits, O.M., Blood, D.C., Gay C.C., Hinchcliff, K.W.,(2000). Veterinary Medicine: a textbook of the diseases of cattle, sheep, pigs, goats and horses, Baillière Tindall, London.
- Reinhold, R.P., (1953). Determination of serum albumin. Clin. Chem. 21, 1370–1372.
- Reitman, S., Frankel, S., (1957). A colorimetric method for determination of serum glutamate oxalo-acetic acid and pyruvic acid transaminases. Am. J. Clin. Pathol. 29, 56–63.
- Salinas, I.; Abelli, L.; Bertoni, F.; Picchietti, S.; Roque, A.; Furones, D.; Cuesta, A.; Meseguer, J. and Esteban, M. (2008) : Monospecies and multispecies probiotic formulations produce different systemic and local immunostimulatory effects in the gilthead seabream (*Sparus aurata* L.). Fish &Shellfish Immunol, 25 (1-2): 114 - 123.
- SAS (1987). Statistical analysis system. User's Guide Statistics. SAS Institute Cary, North Carolina.
- Schalm, O.W., (1986). Veterinary Hematology, 4th Ed., Lea and Febiger, Philadelphia .
- Shalaby, A.M.; khattab, Y.A. and Abdel Rahman, A. M. (2006) : Effects of garlic (*Allium sativum*) and chloramphenicol on growth performance, physiological parameters and survival of Nile tilapia (*Oreochromis niloticus*) J. Venom. Anim. Toxins incl. Trop. Dis., 12, 2, p.200.
- Stoskopf, M. K., (1993). Fish Medicine, W.B. Saunders Comp. Philadelphia . 311–315.
- Tanner, M., Jaya Wardena, P. A., Xusoff, F. M., Subasing, R., (1988). Immunological parameters of Japanese carp *puntius goionotus* (Bleeker) exposed to copper and challenged with *Aeromonas hydrophila*. Shell Fish Immunol. 11, 281–291.
- Taoka, Y.; Maeda, H.; Jo, J.Y.; Jeon, M.J.; Bai, S.C.; Lee, W.J.; Yuge, K. and Koshio, S. (2006b) : Growth, stress tolerance and non-specific immune response of Japanese flounder *Paralichthys olivaceus* to probiotics in a closed recirculating system. Fisheries Sci. 72: 310-321.
- Trinder, P., (1969). Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. Ann. Clin. Biochem. 6, 24–27
- Tsuji, A., Sakurai, H., (1998). Generation of nitric oxide from streptozotocin (STZ) in the presence of copper (II) plus ascorbate. Biochem. Biophys. Res. Commun. 245, 11–16.
- Underwood, E.J., (1977). Trace elements in human and animal nutrition, 3rd ed. Academic Press, New York.