



# Influence of Enzymatic and Mechanical Liquefaction of Seminal Plasma on Freezability of Dromedary Camel Semen

**Khalid Ahmed El-Bahrawy**

*Animal and Poultry Production Division, Desert Research Center (DRC), Cairo, Egypt*

Corresponding author's Email: [bhrawi@yahoo.com](mailto:bhrawi@yahoo.com)

## ABSTRACT

This study aimed to investigate the efficiency of mechanical and enzymatic elimination of semen viscosity in adult dromedary camel bulls' semen on cryopreservation potential of spermatozoa during the breeding season. Bulls showed reaction time  $40.0 \pm 8.23$  seconds and  $251 \pm 24$  seconds mating duration. Physical properties of raw semen showed volume mean value  $5.28 \pm 0.66$  ml, initial viability  $2.5 \pm 0.6$ , initial raw motility  $59.34 \pm 4.99\%$ , livability  $95.3 \pm 2.36\%$ , first and second abnormalities  $4.13 \pm 0.88\%$  and  $7.01 \pm 1.254\%$ , respectively and acrosomal integrity  $5.03 \pm 1.05\%$ . The researcher examined three different treatments for viscosity elimination; namely; Amylase Enzymatic Treatment (AET), Syringe Mechanical Treatment (SMT) and Amylase Syringe Mixed Treatment (ASMT). The results revealed that, a significant deleterious effect of the ASMT on the post-thaw motility ( $M_{PT}$ )  $25.00 \pm 3.69\%$  was observed, with sperm Recovery Rate (RR)  $35.02 \pm 5.02\%$ , contrary to a clear superiority of AET treatment on ( $M_{PT}$ )  $49.00 \pm 4.87\%$ , followed by the SMT treatment ( $M_{PT}$ )  $41.67 \pm 6.72\%$ , with significantly higher RR% ( $76.86 \pm 4.63\%$  and  $62.10 \pm 6.65\%$ ) respectively. The AET recorded the highest acrosomal reaction ( $10.17 \pm 1.11\%$ ), followed by the mixed treatment ( $8.33 \pm 0.14\%$ ), with the least significant effect ( $P < 0.05$ ) on the mechanically treated group ( $7.33 \pm 0.99\%$ ). The results also showed the same trend for first and second abnormalities. Computer assisted semen analysis showed a significant superiority for the AET on mostly all sperm kinetics (DCL, DAP, VAP, VSL), except for DSL, VCL that showed highest significant value for SMT treatment. Conversely, the study recorded the lowest significant values for LIN, STR and WOB in the SMT. These results clarified that both enzymatic and mechanical methods have a positive influence on dromedary camel semen cryopreservation.

**Key words:** Amylase, Cryopreservation, Dromedary, Semen syringing, Viscosity

## INTRODUCTION

Semen processing and cryopreservation are becoming a prerequisite in the application of assisted reproductive technologies in camelids, especially with the increasing of genetically prized valued camels involved in camel racing and beauty contests, although, the technology is yet being sub-optimal in camels due to several challenges. The steady development of Artificial Insemination (AI) with frozen-thawed sperm requires efforts to improve the quality of semen processing techniques. One of the major challenges restricting the development of assisted reproductive technologies (ART's) in camelids is the high viscous nature of seminal plasma (Skidmore et al., 2013; Rateb, 2016). Under natural conditions, complete liquefaction for dromedary camel semen was observed to be  $23.89 \pm 1.49$  hours, varying in a wide range from 18 to 41 hrs (Mal et al., 2016).

Several studies were performed during the last decade for the elimination of camelids' semen viscosity, with various degrees of success. Enzymatic treatments were the most used technique. The cause of the viscous nature of camels' seminal plasma used to be a controversial issue among different studies and authors, either depending on the

ORIGINAL ARTICLE  
pii: S232245681700014-7  
Received: 17 Jul 2017  
Accepted: 22 Aug 2017





### Computer-assisted sperm analysis

Sperm kinetics were measured by a computer-assisted semen analysis (CASA) system [Sperm Vision Lite, a registered trademark of Minitube, USA] attached to a Zeiss warm stage microscope. For samples' evaluation, a 3  $\mu$ l aliquot of the sperm sample was placed in a Lica disposable capillary counting chamber; three fields were analyzed. Kinematic parameters were recorded representing; distance curved line (DCL,  $\mu$ m), distance average path (DAP,  $\mu$ m), distance straight line (DSL,  $\mu$ m), sperm velocity curved line speed (VCL,  $\mu$ m/sec.), velocity average path (VAP  $\mu$ m/sec.), velocity straight line (VSL,  $\mu$ m/sec.), sperm linearity movement (LIN=VSL/VCL), sperm straightness movement (STR=VSL/VAP), sperm balanced movement, Wobble (WOB=VAP/VCL).

### Statistical analysis

The obtained data results were statistically analyzed using one-way analysis of variance using SAS<sup>®</sup> (1999) software program. ANOVA procedure of SAS was used. Mean differences were tested by Duncan's Multiple Range tests (Duncan, 1955) when significant P value was obtained.

## RESULTS

Fifty ejaculates were collected from male camels during this study. The males had a very good body condition with an average body weight of 622 $\pm$ 40.12 kg and body condition score exceeding 2.5 $\pm$ 0.5 (Faye et al., 2001).

Reaction time was of a mean value 40.0 $\pm$  8.23 seconds, with an approximate mating duration of 251 $\pm$ 24 seconds, ejaculate volume 5.28 $\pm$ 0.66 ml, initial viability 2.5 $\pm$ 0.6, initial raw motility 59.34 $\pm$ 4.99%, livability 95.3 $\pm$ 2.36%. First and second abnormalities were 4.13 $\pm$ 0.88% and 7.01 $\pm$ 1.254 %, respectively, while acrosomal integrity was 5.03 $\pm$ 1.05%. Results shown in figure [1(A) (B)] revealed that, soon after the treatments, the initial motility ( $M_i$ ) showed that the 3<sup>rd</sup> group the ASMT exhibited significantly ( $P < 0.05$ ) the highest recorded ( $M_i$ ) reaching 70.00 $\pm$ 4.26% compared to the AET (group 1) and SMT (group 2) being 56.00 $\pm$ 4.00% and 50.00 $\pm$ 3.25%, respectively, the same trend was recorded for the pre-freezing motility ( $M_{pf}$ ) after 4 hour equilibration period, as it showed the superiority of ASMT treatment prior to cryo-preservation followed by the AET and finally the SMT. Contrarily to the ( $M_i$ ) and ( $M_{pf}$ ) values, a significant deleterious effect of the ASMT was observed in post-thaw motility ( $M_{pt}$ ) 25.00 $\pm$ 3.69%, with sperm RR 35.02 $\pm$ 5.02%. Superiority of AET on ( $M_{pt}$ ) 49.00 $\pm$ 4.87% followed by SMT 41.67 $\pm$ 6.72% was observed, with significantly higher RR, 76.86 $\pm$ 4.63% for the AET group and 62.10 $\pm$ 6.65% for the SMT group.

Although the highest post thaw motility ( $M_{pt}$ ) was recorded for AET, but this was accompanied with the highest reacted acrosome (10.17 $\pm$ 1.11%), followed by the ASMT group (8.33 $\pm$ 0.14%), with the least significant effect for the mechanical treated group (7.33 $\pm$ 0.99%) at  $p < 0.05$ . Almost the same trend referred to the treatments effect was observed regarding the first and second abnormalities [Figure 1A and 1B)].

Results illustrated in Figures [2A, 2B, 2C] showed a significant superiority for the AET group on mostly all sperm kinetics. However, the SMT showed the highest distance straight line (DSL,  $\mu$ m), also recorded high sperm track speed, velocity curved line (VCL,  $\mu$ m/sec.), as compared with other treatments. Velocity average path (VAP,  $\mu$ m/sec.) and velocity straight line (VSL,  $\mu$ m/sec.) were significantly higher in AET group [Figure 2A, 2B and 2C)]. Moreover, the lowest values for linearity of sperm movement (LIN=VSL/VCL), straightness (STR =VSL/VAP) and sperm balance movement (wobble) percentage (WOB=VAP/VCL) were recorded in SMT.




Figure 1. (A)




Figure 1. (B)

Figure 1. (A) and (B): Effect of Amylase Enzymatic Treatment (AET), Syringe Mechanical Treatment (SMT) and Amylase Syringe Mixed Treatment (ASMT) on sperm physical properties.




Figure 2. (A)





Figure 2. (B)



**Figure 2. (C)**

**Figure 2.** Effect of amylase enzymatic treatment (AET), syringe mechanical treatment (SMT) and amylase syringe mixed treatment (ASMT) on: Distance curved line (DCL,  $\mu\text{m}$ ), distance average path (DAP,  $\mu\text{m}$ ), distance straight line (DSL,  $\mu\text{m}$ ) (A); Velocity curved line speed (VCL,  $\mu\text{m}/\text{sec}$ ), velocity average path (VAP  $\mu\text{m}/\text{sec}$ ), velocity straight line (VSL,  $\mu\text{m}/\text{sec}$ ) (B); Linearity movement (LIN = VSL/VCL), straightness movement (STR = VSL/VAP), balanced movement, wobble (WOB= VAP/VCL) (C).

## DISCUSSION

There is a general agreement that the difficulty of manipulating hyper viscous semen samples has been a reason to identify several methods being proposed to decrease viscosity for either initial evaluation or further processing steps of ARTs application. In mostly all species, semen rheological properties radically change immediately after collection. In this regard, viscous semen liquefaction is of fundamental importance for the application of a wide range of assisted reproductive technologies (ARTs) in different species, especially camelids (Crichton et al., 2015; Mal et al., 2016; Malo et al., 2017).

The aim of this investigation was to evaluate the sole enzymatic effect or the sole mechanical effect or a mixed in between treatment serving to reduce viscosity in dromedary camel ejaculates to attain the ability for further semen processing. Up to date, there has been a controversy regarding the reason of semen viscosity in camels.

Although camels have no seminal vesicles, but still mucin-like glycoproteins are generally known to be secreted from the bulbourethral glands (More', 1991). As reported by Owen and Katz (2005) and later by Dissanayake et al. (2010), the liquefaction factors responsible for clot lysis are derived from the prostate (plasminogen activator,  $\alpha$ -amylase and prostate-specific antigen PSA). (Behr et al., 2009) recommended the usage  $\alpha$ -amylase and collagenase without any effect on the quality parameters of spermatozoa, reporting that such enzymatic treatment allows better use of semen ejaculates in application sex sorting using flow cytometry technique. On the same basis, (Rateb, 2016), based his study of using high-power low-frequency ultrasound efficiency in eliminating camel semen viscosity since it causes particle size reduction and viscosity alteration in different substances, this was attributed to modification in the microstructure and functional properties of carbohydrates as well as hydrolysis and cleavage of di- and polysaccharides (Kardos and Luche, 2001; Kbbani et al., 2011; Kunaver et al., 2012). Also, (Mendeluk et al., 2000) assumed that proteins are the main components responsible for the rheological behavior of the semen in its normal form. Hence, the rheological properties of hyper viscous semen samples indicated the existence of a highly organized network in which disulfide bonds and oligosaccharide chains complexes to the peptide core. Consequently, this may play a key role affecting the spermatozoa physiology and the sperm motility (Mendeluk et al., 2000; Skidmore et al., 2013).

As reported in this study, a superiority in using amylase as an enzymatic treatment for viscosity elimination, not only is succeeding in improving the motility and the kinetic characteristics of the sperm, but also the remarked softness of the rich fraction sperm clot, that allows easy packing in the French straws. The superiority of amylase for routine work examination of viscous semen specimens was reported earlier by (Dougherty et al., 1978), as they noted that, the lowest level of amylase did not significantly alter semen parameters, with an 80% sufficient rate to liquefy viscous samples, but with careful regard to the level of the enzyme and the interval between addition and analysis must be controlled carefully. Based on the present data, the results elucidate that a high value of detached acrosome and abnormalities were observed when using amylase as compared to other treatments, but with no observation of sperm agglutination as reported lately by (Monaco et al., 2016) using proteolytic enzymes.

Despite the fact that the results of enzymatic treatments surpass those obtained by mechanical methods, but still have acceptable post thaw motility compared to enzymatic treatment. Moreover, the SMT showed the highest distance straight line (DSL,  $\mu\text{m}$ ), also recorded high sperm track speed, velocity curved line (VCL,  $\mu\text{m}/\text{sec}$ .), this is an indicative value for the ability of the mechanical effect to eliminate viscosity. Reduction of the seminal plasma viscosity may be attributed to the changes in the molecular behavior of semen protein fibrils (Amelar, 1962), making them less viscous and more liquefied without an observed reduction in sperm motility or morphological disorders. (Henkel and Schill, 2003) did not recommend the force elimination of viscous seminal fluid through expulsion of the semen ejaculates through a narrow-gauge needle as it caused sperm immotility. The aforementioned were totally contrary to the present data in this study showing an acceptable post-thaw motility after mechanical treatment, this finding was similar to those reported by (Kussler et al., 2014) who recently recommended the physical process of expulsion of semen through a syringe, but with a great attention to the idea that; this methodology may increase sperm DNA fragmentation. The mechanical stress (passage of diluted ejaculate through the syringe) may play a significant role in sperm DNA breaking. Unfortunately, the effect of the treatments on sperm DNA fragmentation in this study was not investigated. Taking into consideration that, although different studies had recorded high post thaw motility for eliminated viscosity with different methods, thus, still the pregnancy rates attempts depending on camelids semen processing show a very limited success (Aller et al., 2003; Deen et al., 2003; Vaughan et al., 2003; Miragaya et al., 2006; Crichton et al., 2016), as compared with fresh semen insemination, to the best of our knowledge the best of pregnancy rates using frozen semen was reported by Bravo et al. (2000b) when using collagenase for viscosity elimination or even for cryopreserved semen in different species. Literature surveys did not clarify any information regarding the reason of low pregnancy rates using frozen semen. Since the stages of embryonic development that depend on paternal genomes begin after the four-to eight-cell stage, the impact of sperm DNA fragmentation in the embryo is usually not observed in *in-vitro* embryo transfer if carried out preferably up to day three. Worth to mention that, the blastocyst stage development rate and implantation till pregnancy may be affected (Borini et al., 2006), in addition to, expected pregnancy losses (Tarozzi et al., 2007; Rougier et al., 2013). Probably the significant deleterious effect on post thaw motility when mixing between enzymatic and mechanical treatments (ASMT) in the current work was associated with the high stress that may lead to a significant DNA breaking. Lack of consistency in information should be addressed by more investigations concerning DNA fragmentation in cryo-preserved camel semen.

## CONCLUSION

From this study, it could be concluded that the viscosity of dromedary camel seminal plasma could be successfully reduced by the use of either mechanical or enzymatic effect, as both may be considered a reliable alternative with different rates of success, with care to the concentration and power of the enzyme and handling and semen ejaculate manipulation during mechanical treatment.

### Acknowledgments

The author is thankful to the director of the Tharb camel hospital Dr. Ashraf El-Sharabasy for his full support. Deepest thanks are due to Dr. Mostafa Osama for assisting the technical work, as well as all team members of the camel reproduction laboratory in the Tharb camel hospital. Special thanks are due to Prof Dr. Mohamed Moawad Khalifa, Dr. Sherif Rateb and Mrs. Arig Rakha for assisting statistical analysis and editing of the manuscript.

### Competing interests

The author declares that he has no conflict of interest with respect to the research, authorship, and/or publication of this article, the author declares that he has no competing interests.

## REFERENCES

- Agostini A, Tawfik E and Campana A (1996). Quantitative post-coital test: sperm counts in cervical mucus after enzymatic liquefaction. *Human Reproduction*, 11(2): 311-317. DOI:10.1093/HUMREP/11.2.311
- Ali HA, Moniem KA and Tingari MD (1976). Some histochemical studies on the prostate, urethral and bulbourethral glands of the one-humped camel (*Camelus dromedarius*). *The Histochemical Journal*, 8:565–578. <https://link.springer.com/content/pdf/10.1007/BF01003958.pdf>
- Aller JF, Rebuffi GE, Cancino AK and Alberio RH (2003). Influence of cryopreservation on motility viability and fertility of llama sperm (*Lama glama*). *Archivos Zootecnia*, 5:15–23. [http://www.uco.es/organiza/servicios/publica/az/php/img/web/30\\_10\\_28\\_02aller.pdf](http://www.uco.es/organiza/servicios/publica/az/php/img/web/30_10_28_02aller.pdf)
- Amelar RD (1962). Coagulation, liquefaction and viscosity of human semen. *The Journal of Urology*, 87:187–190.

- Behr B, Rath D, Mueller P, Hildebrandt TB, Goeritz F, Braun BC, Leahy T, De Graaf SP, Maxwell WMC and Hermes R (2009). Feasibility of sex-sorting sperm from the white and the black rhinoceros (*Ceratotherium simum*, *Diceros bicornis*). *Theriogenology*, 72:353–364. [Doi:10.1016/j.theriogenology.2009.03.001](https://doi.org/10.1016/j.theriogenology.2009.03.001)
- Borini A, Tarozzi N, Bizzaro D, Bonu MA, Fava L, Flamigni C and Coticchio G (2006). Sperm DNA fragmentation paternal effect on early post-implantation embryo development in ART. *Human Reproduction*, 21(11):2876–2881. [Doi:10.1093/humrep/del251](https://doi.org/10.1093/humrep/del251)
- Bravo PW, Ccallo M and Garnica J (2000a). The effect of enzymes on semen viscosity in Llamas and Alpacas. *Small Ruminants Research*. 38: 91-95. [Doi: https://doi.org/10.1016/S0921-4488\(00\)00142-5](https://doi.org/10.1016/S0921-4488(00)00142-5)
- Bravo PW, Skidmore JA and Zhao XX (2000b). Reproductive aspects and storage semen in Camelidae. *Animal Reproduction Science*, 62: 173 – 193. [Doi: https://doi.org/10.1016/S0378-4320\(00\)00158-5](https://doi.org/10.1016/S0378-4320(00)00158-5)
- Ccallo M, Garnica J and Bravo PW (1999). Effect of Fibrinolytics, Hyaluronidase, Collagenase and Trypsin on the semen viscosity in Alpacas. *Proceeding of the II World Congress on Camelids*, Cusco, Peru 1999; p78.
- Crichton EG, Malo C, Pukazhenthil BS, Nagy P and Skidmore JA (2016). Evaluation of cholesterol- treated dromedary camel sperm function by heterologous IVF and AI. *Animal Reproduction Science*. 2016 Nov; 174: 20-28. [Doi: 10.1016/j.anireprosci.2016.08.013](https://doi.org/10.1016/j.anireprosci.2016.08.013)
- Crichton EG, Pukazhenthil BS, Billah M and Skidmore JA (2015). Cholesterol addition aids the cryopreservation of dromedary camel (*Camelus dromedarius*) spermatozoa. *Theriogenology*. 2015 Jan, 15: 83(2):168-174. [Doi: 10.1016/j.theriogenology.2014.09.005](https://doi.org/10.1016/j.theriogenology.2014.09.005). Epub 2014 Sep 16
- Deen A, Vyas S and Sahani MS (2003). Semen collection, cryopreservation and artificial insemination in the dromedary camel. *Animal Reproduction Science*, 77(3-4): 223-233. [Doi: https://doi.org/10.1016/S03784320\(03\)00040-X](https://doi.org/10.1016/S03784320(03)00040-X)
- Desantis S, Monaco D, Accogli G, Albrizio M, El-Bahrawy KA, Rateb SA and Lacalandra GM (2016). Effect of Bromelain and Papain treatments on the glycan pattern of cryopreserved dromedary camel spermatozoa. *The ICAR Satellite Meeting, Camelid Reproduction*, 1-3 July, Tours, France, P.83-86.
- Dissanayake D, Wijesinghe P, Ratnasooriya W and Wimalasena S (2010). Relationship between Seminal Plasma Zinc and semen quality in a sub fertile population. *Journal of Human Reproductive Sciences*, 3(3):124–128. [Doi:10.4103/0974-1208.74153](https://doi.org/10.4103/0974-1208.74153)
- Dougherty KA, Cockett AT and Urry RL (1978). Effect of amylase on sperm motility and viability. *The Journal of Urology*, 120(4):4256.
- Duncan DB (1955). Multiple range and multiple F tests. *Biometrics*. 11:1-42.
- El-Bahrawy KA (2010) Cryopreservation of dromedary camel semen supplemented with  $\alpha$ -amylase enzyme. *Journal of Camel Practice and Research*, 17(2):211-216. [Doi: https://www.cabdirect.org/cabdirect/abstract/20113238297](https://www.cabdirect.org/cabdirect/abstract/20113238297)
- El-Bahrawy KA and El-Hassanein EE (2009). Effect of different mucolytic agents on viscosity and physical characteristics of dromedary camel semen. *Alexandria Journal Agriculture Research*, 54(3):1-6. [Doi: http://agris.fao.org/agris-search/search.do?recordID=EG2012000501](http://agris.fao.org/agris-search/search.do?recordID=EG2012000501)
- Esfandiari N, Burjaq H, Gotlieb L and Casper RF (2008). Seminal hyperviscosity is associated with poor outcome of in-vitro fertilization and embryo transfer: A prospective study. *Fertility and Sterility*, 90(5):1739–1743. [Doi:10.1016/j.fertnstert.2007.09.032](https://doi.org/10.1016/j.fertnstert.2007.09.032)
- Faye B, Bengoumi M, Cleradin A, Tabarani A and Chilliard Y (2001). Body condition score in dromedary camel a tool for management of reproduction. *Emirate Journal of Agriculture Science*, 13:1–6.
- Henkel RR and Schill WB (2003). Sperm preparation for ART. *Reproductive Biology and Endocrinology*, 1:108-142. [Doi: http://dx.doi.org/10.1186/1477-7827-1-108](http://dx.doi.org/10.1186/1477-7827-1-108)
- Johnson L, Berndtson WE and Bickett BW (1976). An improved method for evaluating acrosomes of bovine spermatozoa. *Journal of Animal Science*, 42: 951-954. [doi:10.2527/jas1976.424951x](https://doi.org/10.2527/jas1976.424951x)
- Kardos N and Luche JL (2001). Sonochemistry of carbohydrate compounds. *Carbohydrate Research*, 332:115–131. [Doi: https://doi.org/10.1016/S0008-6215\(01\)00081-7](https://doi.org/10.1016/S0008-6215(01)00081-7)
- Kbbani D, Sepulcre F and Wedekind J (2011). Ultrasound assisted liquefaction of rosemary honey: Influence on rheology and crystal content. *Journal of Food Engineering*, 107:173–178. [Doi: https://doi.org/10.1016/j.jfoodeng.2011.06.027](https://doi.org/10.1016/j.jfoodeng.2011.06.027)
- Kershaw-Young CM and Maxwell WMC (2012). Seminal plasma components in camelids and comparisons with other species. *Reproduction in Domestic Animals*, 47(Suppl.4):369–375. [Doi: https://www.ncbi.nlm.nih.gov/pubmed/22827394](https://www.ncbi.nlm.nih.gov/pubmed/22827394)
- Kershaw-Young CM, Evans G and Maxwell WMC (2012). Glycosaminoglycans in the accessory sex glands, testes and seminal plasma of alpaca and ram. *Reproduction, fertility and development*, 24:362–369. [Doi: http://www.publish.csiro.au/rd/RD11152](http://www.publish.csiro.au/rd/RD11152)



- Keshavarz M, Niasari-Naslaji A, Zare H, Ziapour S, Mirtavoosi M, Omidi M, Kalantari A and Moosavi-Movahedi AA (2016). Effect of Ficin enzyme on semen viscosity in dromedary camel. *Journal of Camel Practice and Research*, 23(2): 219-222. Doi :[10.5958/2277-8934.2016.00037.0](https://doi.org/10.5958/2277-8934.2016.00037.0)
- Kunaver M, Jasiukaityt'e E and Cuk N (2012). Ultrasonically assisted liquefaction of lignocellulosic materials. *Bioresource Technology*, 103: 360–366. Doi: <https://doi.org/10.1016/j.biortech.2011.09.051>
- Kussler AP, Pimentel AM, Alcoba DD, Liu IP, Brum IS, Capp E and Corleta HV (2014). Mechanical processing of hyper viscous semen specimens can negatively affect sperm DNA fragmentation. *International Urology and Nephrology*, 46: 737–742. Doi:[10.1007/s11255-013-0578-9](https://doi.org/10.1007/s11255-013-0578-9)
- Mal G, Vyas S, Srinivasan A, Patil NV and Pathak KML (2016). Studies on liquefaction time and proteins involved in the improvement of seminal characteristics in dromedary camels (*Camelus dromedarius*) Hindawi Publishing Corporation Scientifica, Article ID 4659358, 6 pages. Doi: <http://dx.doi.org/10.1155/2016/4659358>
- Malo C, Crichton EG, Morrell JM, Pukazhenthil BS and Skidmore JA (2017). Single layer centrifugation of fresh dromedary camel semen improves sperm quality and in vitro fertilization capacity compared with simple sperm washing. *Reproduction of Domestic Animals*. 2017 Aug 12. Doi: [10.1111/rda.13036](https://doi.org/10.1111/rda.13036)
- Mendeluk G, Gonzalez Flecha FL, Castello PR and Bregni C (2000). Factors involved in the biochemical etiology of human seminal plasma hyper-viscosity. *Journal Andrology*, 21(2): 262–267. Doi: <http://onlinelibrary.wiley.com/doi/10.1002/j.1939-4640.2000.tb02104.x/epdf>
- Miragaya M, Chaves MG and Aguero A (2006). Reproductive biotechnology in South American camelids. *Small Ruminant Research*, 61: 299–310. Doi: <https://doi.org/10.1016/j.smallrumres.2005.07.017>
- Monaco D, Meriem F, Barbara P, Hammadi M, Khorchani T and Lacalandra GM (2016). Effect of  $\alpha$ -Amylase, Papain, and Spermfluid® treatments on viscosity and semen parameters of dromedary camel ejaculates. *Research in Veterinary Science*, 105: 5–9. Doi: <https://doi.org/10.1016/j.rvsc.2016.01.003>
- More J (1991). Lectin histochemistry of mucus-secreting cells in the calf bulbourethral gland. *Acta Anatomica*, 142: 147–151. Doi: <https://doi.org/10.1159/000147180>
- Morton KM, Vaughan JL and Maxwell WMC (2008). The continued development of artificial insemination technology in alpacas. Rural Industries Research and Development Corporation, Kingston, ACT. [https://www.researchgate.net/publication/242113086\\_The\\_continued\\_development\\_of\\_artificial\\_insemination\\_technologies\\_in\\_alpacas](https://www.researchgate.net/publication/242113086_The_continued_development_of_artificial_insemination_technologies_in_alpacas)
- Niasari-Naslaji A, Mosaferi S, Bahmani N, Gerami A, Gharahdaghi AA, Abarghani A and Ghanbari A (2007). Semen cryopreservation in bactrian camel (*Camelus bactrianus*) using Shotor diluent: Effects of cooling rates and glycerol concentrations. *Theriogenology*, 68:618–625. Doi: <http://dx.doi.org/10.1016/j.theriogenology.2007.04.059>
- Owen DH and Katz DF (2005). A review of the physical and chemical properties of human semen and the formulation of a semen simulant. *Journal of Andrology*. 26(4): 459–469. Doi: [10.2164/jandrol.04104](https://doi.org/10.2164/jandrol.04104)
- Rateb SA (2016). Ultrasound-assisted liquefaction of dromedary camel semen. *Small Ruminant Research*, 141: 48–55. Doi: <http://dx.doi.org/10.1016/j.smallrumres.2016.07.005>
- Rougier N, Uriondo H, Papier S, Checa MA, Sueldo C and Alvarez SC (2013). Changes in DNA fragmentation during sperm preparation for intra-cytoplasmic sperm injection over time. *Fertility and Sterility*, 100(1): 69–74. Doi:[10.1016/j.fertnstert.2013.03.005](https://doi.org/10.1016/j.fertnstert.2013.03.005)
- Santiani A, Huanca W, Sapano R, Huanca T, Sepulveda N and Sanchez R (2005). Effects on the quality of frozen-thawed alpaca (*Lama pacos*) semen using two different cryoprotectants and extenders. *Asian Journal of Andrology*, 7: 303-309. Doi:[10.1111/j.1745-7262.2005.00021.x](https://doi.org/10.1111/j.1745-7262.2005.00021.x)
- SAS (1999). Statistical analysis system user' guide statistics. SAS Institute Inc. Cary NC 27513USA.
- Skidmore JA, Morton KM and Billah M (2013). Artificial insemination in dromedary camels. *Animal Reproduction Science*, 136: 178– 186. Doi: <https://doi.org/10.1016/j.anireprosci.2012.10.008>
- Tarozzi N, Bizzaro D, Flamigni C and Borini A (2007). Clinical relevance of sperm DNA damage in assisted reproduction. *Reproductive Biomedicine Online*, 14(6): 746–757. <https://www.ncbi.nlm.nih.gov/pubmed/17579991>
- Vaughan J, Galloway D and Hopkins D (2003). Artificial insemination in alpacas (*Lama pacos*). Rural Industries Research and Development Corporation, Kingston, ACT, Australia. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.461.1783&rep=rep1&type=pdf>
- WHO (2010). Laboratory manual for the examination and processing of human semen. Vol. I, 5<sup>th</sup> edition. World Health Organization, Geneva. [http://apps.who.int/iris/bitstream/10665/44261/1/9789241547789\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44261/1/9789241547789_eng.pdf)