



Influence of Enzymatic and Mechanical Liquefaction of Seminal Plasma on Freezability of Dromedary Camel Semen

Khalid Ahmed El-Bahrawy

Animal and Poultry Production Division, Desert Research Center (DRC), Cairo, Egypt

Corresponding author's Email: bhrawi@yahoo.com

ABSTRACT

This study aimed to investigate the efficiency of mechanical and enzymatic elimination of semen viscosity in adult dromedary camel bulls' semen on cryopreservation potential of spermatozoa during the breeding season. Bulls showed reaction time 40.0 ± 8.23 seconds and 251 ± 24 seconds mating duration. Physical properties of raw semen showed volume mean value 5.28 ± 0.66 ml, initial viability 2.5 ± 0.6 , initial raw motility $59.34 \pm 4.99\%$, livability $95.3 \pm 2.36\%$, first and second abnormalities $4.13 \pm 0.88\%$ and $7.01 \pm 1.254\%$, respectively and acrosomal integrity $5.03 \pm 1.05\%$. The researcher examined three different treatments for viscosity elimination; namely; Amylase Enzymatic Treatment (AET), Syringe Mechanical Treatment (SMT) and Amylase Syringe Mixed Treatment (ASMT). The results revealed that, a significant deleterious effect of the ASMT on the post-thaw motility (M_{PT}) $25.00 \pm 3.69\%$ was observed, with sperm Recovery Rate (RR) $35.02 \pm 5.02\%$, contrary to a clear superiority of AET treatment on (M_{PT}) $49.00 \pm 4.87\%$, followed by the SMT treatment (M_{PT}) $41.67 \pm 6.72\%$, with significantly higher RR% ($76.86 \pm 4.63\%$ and $62.10 \pm 6.65\%$) respectively. The AET recorded the highest acrosomal reaction ($10.17 \pm 1.11\%$), followed by the mixed treatment ($8.33 \pm 0.14\%$), with the least significant effect ($P < 0.05$) on the mechanically treated group ($7.33 \pm 0.99\%$). The results also showed the same trend for first and second abnormalities. Computer assisted semen analysis showed a significant superiority for the AET on mostly all sperm kinetics (DCL, DAP, VAP, VSL), except for DSL, VCL that showed highest significant value for SMT treatment. Conversely, the study recorded the lowest significant values for LIN, STR and WOB in the SMT. These results clarified that both enzymatic and mechanical methods have a positive influence on dromedary camel semen cryopreservation.

Key words: Amylase, Cryopreservation, Dromedary, Semen syringing, Viscosity

INTRODUCTION

Semen processing and cryopreservation are becoming a prerequisite in the application of assisted reproductive technologies in camelids, especially with the increasing of genetically prized valued camels involved in camel racing and beauty contests, although, the technology is yet being sub-optimal in camels due to several challenges. The steady development of Artificial Insemination (AI) with frozen-thawed sperm requires efforts to improve the quality of semen processing techniques. One of the major challenges restricting the development of assisted reproductive technologies (ART's) in camelids is the high viscous nature of seminal plasma (Skidmore et al., 2013; Rateb, 2016). Under natural conditions, complete liquefaction for dromedary camel semen was observed to be 23.89 ± 1.49 hours, varying in a wide range from 18 to 41 hrs (Mal et al., 2016).

Several studies were performed during the last decade for the elimination of camelids' semen viscosity, with various degrees of success. Enzymatic treatments were the most used technique. The cause of the viscous nature of camels' seminal plasma used to be a controversial issue among different studies and authors, either depending on the

ORIGINAL ARTICLE
pII: S232245681700014-7
Received: 17 Jul 2017
Accepted: 22 Aug 2017

usage of proteolytic enzymes like pepsin, trypsin, alfa-chemotripsin (Ccallo et al.,1999; El-Bahrawy and El-Hassanein, 2009), and lately, the use of Papaine, Bromilaine and Ficin (Desantis et al., 2016; Keshavarz et al., 2016; Monaco et al., 2016), as their theory based on referring viscosity to proteoglycans (protein molecules) (Kershaw-Young and Maxwell, 2012; Mal et al., 2016). Although the superior effect of proteolytic enzymes on seminal plasma viscosity elimination, but still the drastic effect of such enzymes on sperm parameters is to be taken into consideration, (Monaco et al., 2016) observed sperm head agglutination, as well as a direct effect on the sperm's acrosomal integrity, these effects have limited the wide usage of these enzymes in a routine work protocol. Other researchers have depended on using polysaccharides degradation active enzymes (amylase, collagenase, hyaluronidase). As semen viscosity has been postulated that it is caused by glycosaminoglycans (GAGs), which are defined as long un-branched polysaccharides consisting of a repeating disaccharide, (Ali et al., 1976). Lately, (Kershaw-Young et al., 2012) reported in alpaca seminal plasma, a fifteen times higher concentration of GAGs compared to that of rams. Also the claim that camel semen viscosity is attributed to the presence of mucopolysaccharides from the secretions of the bulbourethral gland or the prostate gland (Skidmore et al., 2013). For decades, amylase has been routinely used for the liquefaction of semen hyper-viscosity in humans with different and varied concentrations (Dougherty et al., 1978; Agostini et al., 1996; Henkel and Schill, 2003) with controversial results for its effect. In Camels, (EL-Bahrawy, 2010) and (Monaco et al., 2016) used amylase to eliminate seminal plasma viscosity with different rates of success. In general, enzymatic treatment also affects the diluent components for their long acting effect during the equilibration period of semen processing for cryopreservation as well as sperm cells.

Mechanical methods were also used for seminal plasma viscosity elimination; for Bactrian camel semen, (Niasari-Naslaji et al., 2007) recommended mechanical stirring for viscosity elimination. The WHO manual in 2010 recommended gently running of human semen with hyper viscosity through a 5–10 ml syringe fitted with an 18-gauge needle for several times before initial raw semen parameters examination prior to any processing steps. (Kussler et al., 2014) used the process of expulsion of semen through a 10 ml syringe and an 18-gauge (18G) needle to reduce the seminal viscosity but reported a doubt on the safety of that procedure. In Camelids, early trails were used by Santiani et al., (2005) using the syringe technique prior to cryopreservation of *Lama Pacos* semen. Morton et al., (2008) reported that gentle pipetting of raw semen reduced semen viscosity with a moderate effect on semen parameters after processing. Although semen liquefaction by syringing is easy to be performed, with a quick, effective, and cheap methodology, but it still has not been confirmed whether the process is entirely harmless to the semen samples especially; after cryopreservation (Mendeluk et al., 2000; Esfandiari et al.,2008).

In the view of these facts, and as the etiology and the impact of seminal hyper viscosity elimination procedures on camel semen characteristics and functional capacity have not yet been fully understood with contradictory results, this investigation aims to figure out the effect of combining both methods of enzymatic and mechanical treatment or using each solely to investigate their effect on the physical characteristics of cryopreserved semen.

MATERIALS AND METHODS

Location and experimental animals

The experiment was carried out in an arid area, in the reproduction camel center of the Tharb camel hospital, Qatar. Semen was collected from five dromedary camel bulls of 9 to 15 years of age and 622 ± 40.12 kg average body weight during December 2016 to February 2017. Animals were daily fed at 10 am on a pelleted concentrate mixture (crude protein, 14%), and were further supplemented with barley and dried dates as sources of energy. Also, dry Berseem hay was offered *adlib.* as roughage, and the animals were allowed to drink twice daily.

Ethical approval

This experiment was a routine field work in animal reproduction considering all rules and regulations in conformity with the European Union Directive for the protection of experimental animals (2010/63/EU).

Semen collection

Semen samples were collected three times a week at 6:30 am in a clean area adjacent to the reproduction laboratory. Female teaser was used for semen collection, where it was physically restrained in sternal recumbency position, using a 42 cm bovine Artificial Vagina (AV) adjusted to 40-45°C (El-Bahrawy, 2010). The AV latex inner liner was lubricated from inside using sperm-friendly vaseline assigned for semen collection (Minitube Vaseline, 1000 g, REF.: 11905/0100) to avoid direct contact of spermatozoa with latex inner liner to overcome latex toxicity suspicious. Any contaminated, oligospermic or azospermic specimens were discarded from processing.

Semen diluents preparation and sample processing

Unless stated otherwise, all chemicals and reagents were obtained from Sigma (Sigma-Aldrich) for preparation of Tris-lactose egg yolk extender composed of Tris buffer (3.025%), lactose (5.5%), Citric acid (1.67%), glucose (1%), and supplemented with fresh egg-yolk (20%), the diluent was then divided into two portions A and B. Portion A represented the cooling extender which was added initially to the collected ejaculate with a ratio of 1:1, while portion B was supplemented with 6% glycerol to be added after 2 hours from equilibration (at 5°C) with a ratio 1:1 prior to freezing to make the diluted semen reach a final glycerol level of 3% and a full equilibration period of 4 hours at 5°C, and a final dilution rate 1:3 according to (El-Bahrawy, 2010).

Experimental design

Semen samples were allocated to three groups [Amylase Enzymatic Treatment (AET), Syringe Mechanical Treatment (SMT) and Amylase Syringe Mixed Treatment (ASMT)], with equal fractions for each group.

Enzymatic Procedure [Amylase Enzymatic Treatment (AET)]

After primary check in a pre-test study of the enzyme power units. A concentration of 2.5 ul/ml TERMAMYLSUPRA enzyme, [(a trademarked, amylase available from Termamyl Supra) by Novozymes (Novo Nordisk), Denmark] extracted from, *Bacillus licheniformis* was set up to be used in this experiment. Immediately after collection, Tris lactose extender (Portion A) supplemented with 2.5ul/ml amylase was added to the semen with a 1:1 ratio, according to (EL-Bahrawy, 2010).

Mechanical procedure [Syringe Mechanical Treatment (SMT)]

Soon after the semen was collected, Tris lactose diluent (Portion A) was added with a ratio 1:1, then diluted ejaculates were immediately submitted to a physical process of expulsion of the semen and extender using a 10- or 20-mL (according to the ejaculate volume) for two or three times. This process was repeated two to three times depending on viscosity and visual observation for mixing of semen with the extender.

Mechanical /Enzymatic Procedure [Amylase Syringe Mixed Treatment (ASMT)]

Immediately after collection, Tris lactose extender (Portion A) supplemented with 2.5 µl/ml amylase was added to the semen with a 1:1ratio, according to (EL-Bahrawy, 2010), then the diluted ejaculates were immediately submitted to a physical process of expulsion of the semen using a 10- or 20 ml (according to the ejaculate volume) for two or three times.

Semen cryopreservation

An automatic filling and sealing machine (Minitube, model MPP UNO) and a computer controlled cryo-freezer, with comfortable data input and automatic recording of the freezing curve (Minitube type: Ice Cube 14S) was used for cryopreservation before transporting the cryopreserved straws to the storage tanks for further investigations. Semen straws were subjected to slow thawing temperatures of 40°C for 40 seconds in water bath.

Samples assessment schedule

Immediately after collection, viscosity, viability, motility, abnormalities, livability and acrosomal integrity were initially assessed in raw semen, 10 minutes after initial dilution and performing the treatments (either mechanical, enzymatic or mixed treatments) the initial motility (M_I) was assessed, the pre-freezing motility (M_{PF}) was assessed after 4 hours of equilibration at 5°C and prior to the cryopreservation of samples, and finally post-thawing motility (M_{PT}), recovery rate post thawing (RR%), abnormalities and acrosomal integrity were examined post-thawing.

Semen characteristics assessment

Semen volume was recorded using graduated collecting glass tubes for semen collection. A phase-contrast microscope (Carl ZEISS, AX10_Lab.A1, Germany) with a warm stage adjusted at 37°C was used for the assessment of sperm motility in five different fields at 400X magnification. Both mass motility (viability) in raw semen on a scale from 1 to 3 and motility of freely moving sperm were assessed to the nearest 5%. Sperm livability (live and dead sperm, %) and abnormalities (1st and 2nd abnormalities) were examined using eosin-nigrosin differential staining technique. Acrosomal reaction was examined following the procedure reported by (Johnson et al., 1976).

Semen viscosity was examined according to the (Bravo et al., 2000a) method, using a scale of 1 to 3, where 1 represented low viscosity, 2 for intermediate viscosity and 3 represented the highest viscous samples.

Computer-assisted sperm analysis

Sperm kinetics were measured by a computer-assisted semen analysis (CASA) system [Sperm Vision Lite, a registered trademark of Minitube, USA] attached to a Zeiss warm stage microscope. For samples' evaluation, a 3 μ l aliquot of the sperm sample was placed in a Lica disposable capillary counting chamber; three fields were analyzed. Kinematic parameters were recorded representing; distance curved line (DCL, μ m), distance average path (DAP, μ m), distance straight line (DSL, μ m), sperm velocity curved line speed (VCL, μ m/sec.), velocity average path (VAP μ m/sec.), velocity straight line (VSL, μ m/sec.), sperm linearity movement (LIN=VSL/VCL), sperm straightness movement (STR=VSL/VAP), sperm balanced movement, Wobble (WOB=VAP/VCL).

Statistical analysis

The obtained data results were statistically analyzed using one-way analysis of variance using SAS[®] (1999) software program. ANOVA procedure of SAS was used. Mean differences were tested by Duncan's Multiple Range tests (Duncan, 1955) when significant P value was obtained.

RESULTS

Fifty ejaculates were collected from male camels during this study. The males had a very good body condition with an average body weight of 622 \pm 40.12 kg and body condition score exceeding 2.5 \pm 0.5 (Faye et al., 2001).

Reaction time was of a mean value 40.0 \pm 8.23 seconds, with an approximate mating duration of 251 \pm 24 seconds, ejaculate volume 5.28 \pm 0.66 ml, initial viability 2.5 \pm 0.6, initial raw motility 59.34 \pm 4.99%, livability 95.3 \pm 2.36%. First and second abnormalities were 4.13 \pm 0.88% and 7.01 \pm 1.254 %, respectively, while acrosomal integrity was 5.03 \pm 1.05%. Results shown in figure [1(A) (B)] revealed that, soon after the treatments, the initial motility (M_i) showed that the 3rd group the ASMT exhibited significantly ($P < 0.05$) the highest recorded (M_i) reaching 70.00 \pm 4.26% compared to the AET (group 1) and SMT (group 2) being 56.00 \pm 4.00% and 50.00 \pm 3.25%, respectively, the same trend was recorded for the pre-freezing motility (M_{pf}) after 4 hour equilibration period, as it showed the superiority of ASMT treatment prior to cryo-preservation followed by the AET and finally the SMT. Contrarily to the (M_i) and (M_{pf}) values, a significant deleterious effect of the ASMT was observed in post-thaw motility (M_{pt}) 25.00 \pm 3.69%, with sperm RR 35.02 \pm 5.02%. Superiority of AET on (M_{pt}) 49.00 \pm 4.87% followed by SMT 41.67 \pm 6.72% was observed, with significantly higher RR, 76.86 \pm 4.63% for the AET group and 62.10 \pm 6.65% for the SMT group.

Although the highest post thaw motility (M_{pt}) was recorded for AET, but this was accompanied with the highest reacted acrosome (10.17 \pm 1.11%), followed by the ASMT group (8.33 \pm 0.14%), with the least significant effect for the mechanical treated group (7.33 \pm 0.99%) at $p < 0.05$. Almost the same trend referred to the treatments effect was observed regarding the first and second abnormalities [Figure 1A and 1B)].

Results illustrated in Figures [2A , 2B, 2C] showed a significant superiority for the AET group on mostly all sperm kinetics. However, the SMT showed the highest distance straight line (DSL, μ m), also recorded high sperm track speed, velocity curved line (VCL, μ m /sec.), as compared with other treatments. Velocity average path (VAP, μ m/sec.) and velocity straight line (VSL, μ m/sec.) were significantly higher in AET group [Figure 2A, 2B and 2C)]. Moreover, the lowest values for linearity of sperm movement (LIN=VSL/VCL), straightness (STR =VSL/VAP) and sperm balance movement (wobble) percentage (WOB=VAP/VCL) were recorded in SMT.




Figure 1. (A)




Figure 1. (B)

Figure 1. (A) and (B): Effect of Amylase Enzymatic Treatment (AET), Syringe Mechanical Treatment (SMT) and Amylase Syringe Mixed Treatment (ASMT) on sperm physical properties.




Figure 2. (A)




Figure 2. (B)




Figure 2. (C)

Figure 2. Effect of amylase enzymatic treatment (AET), syringe mechanical treatment (SMT) and amylase syringe mixed treatment (ASMT) on: Distance curved line (DCL, μm), distance average path (DAP, μm), distance straight line (DSL, μm) (A); Velocity curved line speed (VCL, $\mu\text{m}/\text{sec}$), velocity average path (VAP $\mu\text{m}/\text{sec}$), velocity straight line (VSL, $\mu\text{m}/\text{sec}$) (B); Linearity movement (LIN = VSL/VCL), straightness movement (STR = VSL/VAP), balanced movement, wobble (WOB = VAP/VCL) (C).

DISCUSSION

There is a general agreement that the difficulty of manipulating hyper viscous semen samples has been a reason to identify several methods being proposed to decrease viscosity for either initial evaluation or further processing steps of ARTs application. In mostly all species, semen rheological properties radically change immediately after collection. In this regard, viscous semen liquefaction is of fundamental importance for the application of a wide range of assisted reproductive technologies (ARTs) in different species, especially camelids (Crichton et al., 2015; Mal et al., 2016; Malo et al., 2017).

The aim of this investigation was to evaluate the sole enzymatic effect or the sole mechanical effect or a mixed in between treatment serving to reduce viscosity in dromedary camel ejaculates to attain the ability for further semen processing. Up to date, there has been a controversy regarding the reason of semen viscosity in camels.

Although camels have no seminal vesicles, but still mucin-like glycoproteins are generally known to be secreted from the bulbourethral glands (More', 1991). As reported by Owen and Katz (2005) and later by Dissanayake et al. (2010), the liquefaction factors responsible for clot lysis are derived from the prostate (plasminogen activator, α -amylase and prostate-specific antigen PSA). (Behr et al., 2009) recommended the usage α -amylase and collagenase without any effect on the quality parameters of spermatozoa, reporting that such enzymatic treatment allows better use of semen ejaculates in application sex sorting using flow cytometry technique. On the same basis, (Rateb, 2016), based his study of using high-power low-frequency ultrasound efficiency in eliminating camel semen viscosity since it causes particle size reduction and viscosity alteration in different substances, this was attributed to modification in the microstructure and functional properties of carbohydrates as well as hydrolysis and cleavage of di- and polysaccharides (Kardos and Luche, 2001; Kbbani et al., 2011; Kunaver et al., 2012). Also, (Mendeluk et al., 2000) assumed that proteins are the main components responsible for the rheological behavior of the semen in its normal form. Hence, the rheological properties of hyper viscous semen samples indicated the existence of a highly organized network in which disulfide bonds and oligosaccharide chains complexes to the peptide core. Consequently, this may play a key role affecting the spermatozoa physiology and the sperm motility (Mendeluk et al., 2000; Skidmore et al., 2013).

As reported in this study, a superiority in using amylase as an enzymatic treatment for viscosity elimination, not only is succeeding in improving the motility and the kinetic characteristics of the sperm, but also the remarked softness of the rich fraction sperm clot, that allows easy packing in the French straws. The superiority of amylase for routine work examination of viscous semen specimens was reported earlier by (Dougherty et al., 1978), as they noted that, the lowest level of amylase did not significantly alter semen parameters, with an 80% sufficient rate to liquefy viscous samples, but with careful regard to the level of the enzyme and the interval between addition and analysis must be controlled carefully. Based on the present data, the results elucidate that a high value of detached acrosome and abnormalities were observed when using amylase as compared to other treatments, but with no observation of sperm agglutination as reported lately by (Monaco et al., 2016) using proteolytic enzymes.

Despite the fact that the results of enzymatic treatments surpass those obtained by mechanical methods, but still have acceptable post thaw motility compared to enzymatic treatment. Moreover, the SMT showed the highest distance straight line (DSL, μm), also recorded high sperm track speed, velocity curved line (VCL, $\mu\text{m}/\text{sec}$.), this is an indicative value for the ability of the mechanical effect to eliminate viscosity. Reduction of the seminal plasma viscosity may be attributed to the changes in the molecular behavior of semen protein fibrils (Amelar, 1962), making them less viscous and more liquefied without an observed reduction in sperm motility or morphological disorders. (Henkel and Schill, 2003) did not recommend the force elimination of viscous seminal fluid through expulsion of the semen ejaculates through a narrow-gauge needle as it caused sperm immotility. The aforementioned were totally contrary to the present data in this study showing an acceptable post-thaw motility after mechanical treatment, this finding was similar to those reported by (Kussler et al., 2014) who recently recommended the physical process of expulsion of semen through a syringe, but with a great attention to the idea that; this methodology may increase sperm DNA fragmentation. The mechanical stress (passage of diluted ejaculate through the syringe) may play a significant role in sperm DNA breaking. Unfortunately, the effect of the treatments on sperm DNA fragmentation in this study was not investigated. Taking into consideration that, although different studies had recorded high post thaw motility for eliminated viscosity with different methods, thus, still the pregnancy rates attempts depending on camelids semen processing show a very limited success (Aller et al., 2003; Deen et al., 2003; Vaughan et al., 2003; Miragaya et al., 2006; Crichton et al., 2016), as compared with fresh semen insemination, to the best of our knowledge the best of pregnancy rates using frozen semen was reported by Bravo et al. (2000b) when using collagenase for viscosity elimination or even for cryopreserved semen in different species. Literature surveys did not clarify any information regarding the reason of low pregnancy rates using frozen semen. Since the stages of embryonic development that depend on paternal genomes begin after the four-to eight-cell stage, the impact of sperm DNA fragmentation in the embryo is usually not observed in *in-vitro* embryo transfer if carried out preferably up to day three. Worth to mention that, the blastocyst stage development rate and implantation till pregnancy may be affected (Borini et al., 2006), in addition to, expected pregnancy losses (Tarozzi et al., 2007; Rougier et al., 2013). Probably the significant deleterious effect on post thaw motility when mixing between enzymatic and mechanical treatments (ASMT) in the current work was associated with the high stress that may lead to a significant DNA breaking. Lack of consistency in information should be addressed by more investigations concerning DNA fragmentation in cryo-preserved camel semen.

CONCLUSION

From this study, it could be concluded that the viscosity of dromedary camel seminal plasma could be successfully reduced by the use of either mechanical or enzymatic effect, as both may be considered a reliable alternative with different rates of success, with care to the concentration and power of the enzyme and handling and semen ejaculate manipulation during mechanical treatment.

Acknowledgments

The author is thankful to the director of the Tharb camel hospital Dr. Ashraf El-Sharabasy for his full support. Deepest thanks are due to Dr. Mostafa Osama for assisting the technical work, as well as all team members of the camel reproduction laboratory in the Tharb camel hospital. Special thanks are due to Prof Dr. Mohamed Moawad Khalifa, Dr. Sherif Rateb and Mrs. Arig Rakha for assisting statistical analysis and editing of the manuscript.

Competing interests

The author declares that he has no conflict of interest with respect to the research, authorship, and/or publication of this article, the author declares that he has no competing interests.

REFERENCES

- Agostini A, Tawfik E and Campana A (1996). Quantitative post-coital test: sperm counts in cervical mucus after enzymatic liquefaction. *Human Reproduction*, 11(2): 311-317. DOI:10.1093/HUMREP/11.2.311
- Ali HA, Moniem KA and Tingari MD (1976). Some histochemical studies on the prostate, urethral and bulbourethral glands of the one-humped camel (*Camelus dromedarius*). *The Histochemical Journal*, 8:565–578. <https://link.springer.com/content/pdf/10.1007/BF01003958.pdf>
- Aller JF, Rebuffi GE, Cancino AK and Alberio RH (2003). Influence of cryopreservation on motility viability and fertility of llama sperm (*Lama glama*). *Archivos Zootecnia*, 5:15–23. http://www.uco.es/organiza/servicios/publica/az/php/img/web/30_10_28_02aller.pdf
- Amelar RD (1962). Coagulation, liquefaction and viscosity of human semen. *The Journal of Urology*, 87:187–190.

- Behr B, Rath D, Mueller P, Hildebrandt TB, Goeritz F, Braun BC, Leahy T, De Graaf SP, Maxwell WMC and Hermes R (2009). Feasibility of sex-sorting sperm from the white and the black rhinoceros (*Ceratotherium simum*, *Diceros bicornis*). *Theriogenology*, 72:353–364. [Doi:10.1016/j.theriogenology.2009.03.001](https://doi.org/10.1016/j.theriogenology.2009.03.001)
- Borini A, Tarozzi N, Bizzaro D, Bonu MA, Fava L, Flamigni C and Coticchio G (2006). Sperm DNA fragmentation paternal effect on early post-implantation embryo development in ART. *Human Reproduction*, 21(11):2876–2881. [Doi:10.1093/humrep/del251](https://doi.org/10.1093/humrep/del251)
- Bravo PW, Ccallo M and Garnica J (2000a). The effect of enzymes on semen viscosity in Llamas and Alpacas. *Small Ruminants Research*. 38: 91-95. [Doi: https://doi.org/10.1016/S0921-4488\(00\)00142-5](https://doi.org/10.1016/S0921-4488(00)00142-5)
- Bravo PW, Skidmore JA and Zhao XX (2000b). Reproductive aspects and storage semen in Camelidae. *Animal Reproduction Science*, 62: 173 – 193. [Doi: https://doi.org/10.1016/S0378-4320\(00\)00158-5](https://doi.org/10.1016/S0378-4320(00)00158-5)
- Ccallo M, Garnica J and Bravo PW (1999). Effect of Fibrinolytics, Hyaluronidase, Collagenase and Trypsin on the semen viscosity in Alpacas. *Proceeding of the II World Congress on Camelids*, Cusco, Peru 1999; p78.
- Crichton EG, Malo C, Pukazhenthil BS, Nagy P and Skidmore JA (2016). Evaluation of cholesterol- treated dromedary camel sperm function by heterologous IVF and AI. *Animal Reproduction Science*. 2016 Nov; 174: 20-28. [Doi: 10.1016/j.anireprosci.2016.08.013](https://doi.org/10.1016/j.anireprosci.2016.08.013)
- Crichton EG, Pukazhenthil BS, Billah M and Skidmore JA (2015). Cholesterol addition aids the cryopreservation of dromedary camel (*Camelus dromedarius*) spermatozoa. *Theriogenology*. 2015 Jan, 15: 83(2):168-174. [Doi: 10.1016/j.theriogenology.2014.09.005](https://doi.org/10.1016/j.theriogenology.2014.09.005). Epub 2014 Sep 16
- Deen A, Vyas S and Sahani MS (2003). Semen collection, cryopreservation and artificial insemination in the dromedary camel. *Animal Reproduction Science*, 77(3-4): 223-233. [Doi: https://doi.org/10.1016/S03784320\(03\)00040-X](https://doi.org/10.1016/S03784320(03)00040-X)
- Desantis S, Monaco D, Accogli G, Albrizio M, El-Bahrawy KA, Rateb SA and Lacalandra GM (2016). Effect of Bromelain and Papain treatments on the glycan pattern of cryopreserved dromedary camel spermatozoa. *The ICAR Satellite Meeting, Camelid Reproduction*, 1-3 July, Tours, France, P.83-86.
- Dissanayake D, Wijesinghe P, Ratnasooriya W and Wimalasena S (2010). Relationship between Seminal Plasma Zinc and semen quality in a sub fertile population. *Journal of Human Reproductive Sciences*, 3(3):124–128. [Doi:10.4103/0974-1208.74153](https://doi.org/10.4103/0974-1208.74153)
- Dougherty KA, Cockett AT and Urry RL (1978). Effect of amylase on sperm motility and viability. *The Journal of Urology*, 120(4):4256.
- Duncan DB (1955). Multiple range and multiple F tests. *Biometrics*. 11:1-42.
- El-Bahrawy KA (2010) Cryopreservation of dromedary camel semen supplemented with α -amylase enzyme. *Journal of Camel Practice and Research*, 17(2):211-216. [Doi: https://www.cabdirect.org/cabdirect/abstract/20113238297](https://www.cabdirect.org/cabdirect/abstract/20113238297)
- El-Bahrawy KA and El-Hassanein EE (2009). Effect of different mucolytic agents on viscosity and physical characteristics of dromedary camel semen. *Alexandria Journal Agriculture Research*, 54(3):1-6. [Doi: http://agris.fao.org/agris-search/search.do?recordID=EG2012000501](http://agris.fao.org/agris-search/search.do?recordID=EG2012000501)
- Esfandiari N, Burjaq H, Gotlieb L and Casper RF (2008). Seminal hyperviscosity is associated with poor outcome of in-vitro fertilization and embryo transfer: A prospective study. *Fertility and Sterility*, 90(5):1739–1743. [Doi:10.1016/j.fertnstert.2007.09.032](https://doi.org/10.1016/j.fertnstert.2007.09.032)
- Faye B, Bengoumi M, Cleradin A, Tabarani A and Chilliard Y (2001). Body condition score in dromedary camel a tool for management of reproduction. *Emirate Journal of Agriculture Science*, 13:1–6.
- Henkel RR and Schill WB (2003). Sperm preparation for ART. *Reproductive Biology and Endocrinology*, 1:108-142. [Doi: http://dx.doi.org/10.1186/1477-7827-1-108](http://dx.doi.org/10.1186/1477-7827-1-108)
- Johnson L, Berndtson WE and Bickett BW (1976). An improved method for evaluating acrosomes of bovine spermatozoa. *Journal of Animal Science*, 42: 951-954. [doi:10.2527/jas1976.424951x](https://doi.org/10.2527/jas1976.424951x)
- Kardos N and Luche JL (2001). Sonochemistry of carbohydrate compounds. *Carbohydrate Research*, 332:115–131. [Doi: https://doi.org/10.1016/S0008-6215\(01\)00081-7](https://doi.org/10.1016/S0008-6215(01)00081-7)
- Kbbani D, Sepulcre F and Wedekind J (2011). Ultrasound assisted liquefaction of rosemary honey: Influence on rheology and crystal content. *Journal of Food Engineering*, 107:173–178. [Doi: https://doi.org/10.1016/j.jfoodeng.2011.06.027](https://doi.org/10.1016/j.jfoodeng.2011.06.027)
- Kershaw-Young CM and Maxwell WMC (2012). Seminal plasma components in camelids and comparisons with other species. *Reproduction in Domestic Animals*, 47(Suppl.4):369–375. [Doi: https://www.ncbi.nlm.nih.gov/pubmed/22827394](https://www.ncbi.nlm.nih.gov/pubmed/22827394)
- Kershaw-Young CM, Evans G and Maxwell WMC (2012). Glycosaminoglycans in the accessory sex glands, testes and seminal plasma of alpaca and ram. *Reproduction, fertility and development*, 24:362–369. [Doi: http://www.publish.csiro.au/rd/RD11152](http://www.publish.csiro.au/rd/RD11152)

- Keshavarz M, Niasari-Naslaji A, Zare H, Ziapour S, Mirtavoosi M, Omidi M, Kalantari A and Moosavi-Movahedi AA (2016). Effect of Ficin enzyme on semen viscosity in dromedary camel. *Journal of Camel Practice and Research*, 23(2): 219-222. Doi :[10.5958/2277-8934.2016.00037.0](https://doi.org/10.5958/2277-8934.2016.00037.0)
- Kunaver M, Jasiukaityt'e E and Cuk N (2012). Ultrasonically assisted liquefaction of lignocellulosic materials. *Bioresource Technology*, 103: 360–366. Doi: <https://doi.org/10.1016/j.biortech.2011.09.051>
- Kussler AP, Pimentel AM, Alcoba DD, Liu IP, Brum IS, Capp E and Corleta HV (2014). Mechanical processing of hyper viscous semen specimens can negatively affect sperm DNA fragmentation. *International Urology and Nephrology*, 46: 737–742. Doi:[10.1007/s11255-013-0578-9](https://doi.org/10.1007/s11255-013-0578-9)
- Mal G, Vyas S, Srinivasan A, Patil NV and Pathak KML (2016). Studies on liquefaction time and proteins involved in the improvement of seminal characteristics in dromedary camels (*Camelus dromedarius*) Hindawi Publishing Corporation Scientifica, Article ID 4659358, 6 pages. Doi: <http://dx.doi.org/10.1155/2016/4659358>
- Malo C, Crichton EG, Morrell JM, Pukazhenthii BS and Skidmore JA (2017). Single layer centrifugation of fresh dromedary camel semen improves sperm quality and in vitro fertilization capacity compared with simple sperm washing. *Reproduction of Domestic Animals*. 2017 Aug 12. Doi: [10.1111/rda.13036](https://doi.org/10.1111/rda.13036)
- Mendeluk G, Gonzalez Flecha FL, Castello PR and Bregni C (2000). Factors involved in the biochemical etiology of human seminal plasma hyper-viscosity. *Journal Andrology*, 21(2): 262–267. Doi: <http://onlinelibrary.wiley.com/doi/10.1002/j.1939-4640.2000.tb02104.x/epdf>
- Miragaya M, Chaves MG and Aguero A (2006). Reproductive biotechnology in South American camelids. *Small Ruminant Research*, 61: 299–310. Doi: <https://doi.org/10.1016/j.smallrumres.2005.07.017>
- Monaco D, Meriem F, Barbara P, Hammadi M, Khorchani T and Lacalandra GM (2016). Effect of α -Amylase, Papain, and Spermfluid® treatments on viscosity and semen parameters of dromedary camel ejaculates. *Research in Veterinary Science*, 105: 5–9. Doi: <https://doi.org/10.1016/j.rvsc.2016.01.003>
- More´ J (1991). Lectin histochemistry of mucus-secreting cells in the calf bulbourethral gland. *Acta Anatomica*, 142: 147–151. Doi: <https://doi.org/10.1159/000147180>
- Morton KM, Vaughan JL and Maxwell WMC (2008). The continued development of artificial insemination technology in alpacas. Rural Industries Research and Development Corporation, Kingston, ACT. https://www.researchgate.net/publication/242113086_The_continued_development_of_artificial_insemination_technologies_in_alpacas
- Niasari-Naslaji A, Mosaferi S, Bahmani N, Gerami A, Gharahdaghi AA, Abarghani A and Ghanbari A (2007). Semen cryopreservation in bactrian camel (*Camelus bactrianus*) using Shotor diluent: Effects of cooling rates and glycerol concentrations. *Theriogenology*, 68:618–625. Doi: <http://dx.doi.org/10.1016/j.theriogenology.2007.04.059>
- Owen DH and Katz DF (2005). A review of the physical and chemical properties of human semen and the formulation of a semen simulant. *Journal of Andrology*. 26(4): 459–469. Doi: [10.2164/jandrol.04104](https://doi.org/10.2164/jandrol.04104)
- Rateb SA (2016). Ultrasound-assisted liquefaction of dromedary camel semen. *Small Ruminant Research*, 141: 48–55. Doi: <http://dx.doi.org/10.1016/j.smallrumres.2016.07.005>
- Rougier N, Uriondo H, Papier S, Checa MA, Sueldo C and Alvarez SC (2013). Changes in DNA fragmentation during sperm preparation for intra-cytoplasmic sperm injection over time. *Fertility and Sterility*, 100(1): 69–74. Doi:[10.1016/j.fertnstert.2013.03.005](https://doi.org/10.1016/j.fertnstert.2013.03.005)
- Santiani A, Huanca W, Sapano R, Huanca T, Sepulveda N and Sanchez R (2005). Effects on the quality of frozen-thawed alpaca (*Lama pacos*) semen using two different cryoprotectants and extenders. *Asian Journal of Andrology*, 7: 303-309. Doi:[10.1111/j.1745-7262.2005.00021.x](https://doi.org/10.1111/j.1745-7262.2005.00021.x)
- SAS (1999). Statistical analysis system user' guide statistics. SAS Institute Inc. Cary NC 27513USA.
- Skidmore JA, Morton KM and Billah M (2013). Artificial insemination in dromedary camels. *Animal Reproduction Science*, 136: 178– 186. Doi: <https://doi.org/10.1016/j.anireprosci.2012.10.008>
- Tarozzi N, Bizzaro D, Flamigni C and Borini A (2007). Clinical relevance of sperm DNA damage in assisted reproduction. *Reproductive Biomedicine Online*, 14(6): 746–757. <https://www.ncbi.nlm.nih.gov/pubmed/17579991>
- Vaughan J, Galloway D and Hopkins D (2003). Artificial insemination in alpacas (*Lama pacos*). Rural Industries Research and Development Corporation, Kingston, ACT, Australia. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.461.1783&rep=rep1&type=pdf>
- WHO (2010). Laboratory manual for the examination and processing of human semen. Vol. I, 5th edition. World Health Organization, Geneva. http://apps.who.int/iris/bitstream/10665/44261/1/9789241547789_eng.pdf