



# Prevalence of *Vibrio parahaemolyticus* in seabass (*Dicentrarchus Labrax*) and seabream (*Sparus aurata*) and Detection of Streptomycin-resistant Strains

Adel M. El-Gamal<sup>1\*</sup> and Engy F. EL-Bahi<sup>2</sup>

<sup>1</sup>Bacteriology unit, Animal Health Research Institute, Kafr El-Sheikh branch, Agriculture research center (ARC), Egypt.

<sup>2</sup>Food hygiene unit, Animal Health Research Institute, Kafr El-Sheikh branch, Agriculture research center (ARC), Egypt.

\*Corresponding author's Email: [adelelgamal5544@yahoo.com](mailto:adelelgamal5544@yahoo.com); ORCID: 0000-0002-7697-929X

## ABSTRACT

*Vibrio* species are the most common and serious pathogens in fish and shellfish marine aquaculture worldwide. The present study aimed to determine the prevalence of *Vibrio* spp. in seabass and seabream in fish markets, especially streptomycin-resistant strains that have great public health importance. A total of 30 seabass (*Dicentrarchus Labrax*) and 30 seabream (*Sparus aurata*) were purchased from fish markets at Kafr El Sheikh Governorate and subjected to bacteriological examination. The PCR assay was used for the detection of virulence genes (*tdh* and *trh*), aminoglycoside resistance gene (*aadA1*), and *toxR* gene. The results indicated that the total prevalence of *Vibrio* spp. was 26.66%, including *V. parahaemolyticus* (8.3%), *V. alginolyticus* (8.3%), *V. mimicus* (3.3%), *V. harveyi* (5%) and *V. vulnificus* (1.6%). The *toxR*, *trh*, and *aadA1* genes were found in all *V. parahaemolyticus* isolates while *tdh* gene was found in 80% of isolates. Antimicrobial sensitivity test of *V. parahaemolyticus* isolates showed sensitivity to ciprofloxacin, norfloxacin, cefotaxime, and chloramphenicol. *Vibrio parahaemolyticus* isolates were resistant to ampicillin, erythromycin, streptomycin, and gentamycin. The present results indicated that good hygienic measures should be taken to avoid infection with *Vibrio* species, especially *V. parahaemolyticus* that can pose a great risk to human health.

**Keywords:** Antibiotic resistance, Seabass, Seabream, Streptomycin, *Vibrio parahaemolyticus*.

## INTRODUCTION

*Vibrio* genus contains Gram-negative, halophilic, rod-shaped, non-spore forming, oxidase-positive bacteria, which widespread in the coastal and estuarine environments (Austin and Austin, 2007). *Vibrio parahaemolyticus* is the most recorded pathogenic species of *Vibrio* genus and affects persons who consume improperly cooked or raw seafood (Raissy et al., 2015). This foodborne bacteria is reported as the main cause of seafood-borne illness in Egypt and many other countries around the world such as United States, Malaysia, Thailand, Korea, China, and Japan (Yoon et al., 2008; Iwahori et al., 2010; Abdel-Azeem et al., 2016). Infection with *V. parahaemolyticus* may cause acute human gastroenteritis, the major symptoms of which are headache, diarrhea, abdominal pain, and in some cases, septicemia (Broberg et al., 2011; Wang et al., 2015; Su and Liu, 2017). In coastal areas of the world, like Japan, *V. parahaemolyticus* has been regularly recognized as the main cause of sporadic cases of gastroenteritis (Qadri et al., 2005; Wang et al., 2017). In China, about 322 gastroenteritis outbreaks due to *V. parahaemolyticus* infection were reported from 2003 to 2008 (Wu et al., 2014). Multiplication of *V. parahaemolyticus* is related to water temperature and season (Deepanjali et al., 2005; Angela et al., 2006), with the highest prevalence in summer due to the higher salinity of water than other seasons (Zulkifli et al., 2009).

The pathogenicity of bacteria depends mainly on some virulence factors and virulence genes, which act together as major orchestrators. The most virulence genes leading to pathogenicity of *V. parahaemolyticus* are hemolysin genes (*tdh* and *trh*) (Hiyoshi et al., 2010). Molecular epidemiological studies demonstrated a clear relation between the hemolysin genes and disease-causing ability of *V. parahaemolyticus* (Kishishita et al., 1992; DePaola et al., 2003; Vongxay et al., 2008; Chao et al., 2009; Han et al., 2015; Hasrimi et al., 2018). These two genes were recorded in the most isolates from clinical cases of *V. parahaemolyticus* infections (Bej et al., 1999; Rojas et al., 2011). The *tdh* and *trh* genes encode virulence factors of thermostable direct hemolysin (TDH), and TDH-related hemolysin (TRH), respectively, which are involved in important pathogenic activities, such as enterotoxicity, hemolytic activity, cytotoxicity and cardiotoxicity (Shirai et al., 1990; Osawa et al., 1996).

The *toxR* gene is a pandemic marker gene for all *V. parahaemolyticus* strains either pathogenic or nonpathogenic one, and it was recorded in some other *Vibrio* species (Kim et al., 1999). The sequence of *toxR* gene can be used for molecular identification of *V. parahaemolyticus* (Yung et al., 1999; Hubbard et al., 2016). The *aadA1* and *aadA2* encode

ORIGINAL ARTICLE  
 pii: S232245682000042-10  
 Received: 28 Jun. 2020  
 Accepted: 10 Aug 2020

aminoglycoside adenylyl transferase and confer resistance to streptomycin, already are detected in *Vibrio* species isolates (Dalsgaard et al., 2001).

Cooking and frying of marine fish reduce the count of *V. parahaemolyticus*. After cooking (in oven 120 °C for 35 min), the percentage reduction in total count of *V. parahaemolyticus* was 98.2%, while after frying for 10 min at 190 °C, *V. parahaemolyticus* was completely destroyed and the percentage reduction was 100% (Saad et al., 2015); or even boiling at 64 °C for more than 90 seconds can kill *V. parahaemolyticus* (ICMSF, 1996). The bacteria will be removed by using high cooking temperature, although the toxin might remain in the foodstuff depending on the processing conditions (Raissy et al., 2015). The current study aimed to determine the prevalence of *Vibrio* spp., especially streptomycin-resistant strains, in seabass and seabream in fish markets of Kafr El Sheikh Governorate, Egypt.

## MATERIALS AND METHODS

### Samples collection

Thirty seabass and 30 seabream with a weight range of 100-250 g were purchased from fish markets at Kafr El Sheikh Governorate from February to August 2019. All samples were transferred in ice box to Animal Health Research Institute, Kafr El Sheikh laboratory, Egypt.

### Bacteriological examination

Bacteriological examinations were done according to ISO/ TS 21872-1 (2007) and ISO/ TS 21872-2 (2007).

### Samples preparation

After skin sterilization with alcohol, the muscles above the lateral line were removed, 25 g of each fish sample were mixed with 225 ml of alkaline saline peptone water (APW, pH 8.6) in a Stomacher bag. After that, these mixtures were incubated at 37 °C for 8-16 hours.

### Isolation of *Vibrio* species

After the incubation period, the upper layer of the alkaline saline peptone water (APW) enrichment broth was inoculated on Thiosulfate-citrate-bile salts-sucrose (TCBS) agar (Oxoid, UK), and then these plates were incubated at 37 °C for another 18-24 hours. After that, growing colonies were used for further screening tests including Gram staining, oxidase and catalase tests.

### Biochemical identification

Suspected colonies of *Vibrio* spp. on TCBS media and positive oxidase test were subjected to further identification by Microbact GNB kit (Oxoid, UK).

### Polymerase chain action

Suspected isolates of the *V. parahaemolyticus* were examined by using PCR for the detection of virulence genes (*tdh* and *trh*), *toxR* gene, and *aadA1* gene. DNA extraction were performed according to the manufacturer's recommendations by using the QIA amp DNA Mini kit (Qiagen, Germany, GmbH). Oligonucleotide primers were supplied from Metabion (Germany). The primers were utilized in a 25- µl reaction containing 12.5 µl of Emerald Amp Max PCR Master Mix (Takara, Japan) using an Applied Biosystem 2720 thermal cycler. Primers used and PCR conditions are presented in Table 1. The products of PCR were separated by electrophoresis on 1% Agarose gel (Appllichem, Germany, GmbH) in 1x TBE buffer at room temperature using gradients of 5 V/cm. The gel was photographed by a gel documentation system (Alpha Innotech, Biometra).

### Antimicrobial susceptibility test

Antimicrobial disk susceptibility test were performed as described by the Clinical and Laboratory Standards Institute (CLSI, 2012).

**Table 1.** Primers sequences, target genes, amplicon sizes and cycling conditions.

Target genes	Primers Sequence (5'-3')	Amplified segment (base pair)	Primary denaturation	Amplification (35 cycles)			Final extension	Reference
				Secondary denaturation	Annealing	Extension		
<i>toxR</i>	F <i>GTCTTCTGACGCAATCGTTG</i>	368	94°C	94°C	55°C	72°C	72°C	Kim et al., 1999
	R <i>ATACGAGTGGTTGCTGTCATG</i>		5 min.	30 sec.	40 sec.	40 sec.		
<i>aadA1</i>	F <i>TATCAGAGGTAGTTGGCGTCAT</i>	484	94°C	94°C	54°C	72°C	72°C	Randall et al. 2004
	R <i>GTTCCATAGCGTTAAGGTTTCATT</i>		5 min.	30 sec.	40 sec.	45 sec.		
<i>trh</i>	F <i>GGCTCAAAATGGTTAAGCG</i>	250	94°C	94°C	54°C	72°C	72°C	Mustapha et al., 2013
	R <i>CATTCCGCTCTCATATGC</i>		5 min.	30 sec.	30 sec.	30 sec.		
<i>tdh</i>	F <i>CCATCTGTCCCTTTTCTCTGC</i>	373	94°C	94°C	54°C	72°C	72°C	
	R <i>CAAATACATTTTACTTGG</i>		5 min.	30 sec.	30 sec.	40 sec.		

## RESULTS AND DISSCUSION

*Vibrio* spp. commonly inhabit the marine environments and can be found in the fresh water (Sujeewa et al., 2009). Seafood may be a vehicle for most of the bacterial pathogens such as *Vibrio* spp. (Huss, 1997). Various outbreaks of bacterial disease associated with seafood consumption have been reported (Friesema et al., 2012). Recently, *V. parahaemolyticus* recoded as an important species causing seafood infection associated with gastroenteritis illness in humans.

Table 2 shows that the total incidence of *Vibrio* spp. isolated from the examined seabass and seabream samples is 26.66% (16 out of 60 samples). Raissy et al. (2015) and Azwail et al. (2016) recorded nearly similar results (22% and 22.9%, respectively). however, the result of present study is lower than that recorded by Pal and Das (2010), Saad et al. (2015), Abdel-Azeem et al. (2016), Fri et al. (2017), and Hemmat et al. (2018). These differences may be due to difference in the type of examined fish, difference in the method of bacterial isolation, or difference in the hygienic state of fish sources. Additionally, the difference in results can be attributed to difference in season of sampling, as *Vibrio* spp. has been reported to have higher concentrations in summer seasons due to higher water salinity levels than other seasons (Zulkifli et al., 2009). As presented in table 2, several *Vibrio* strains were isolated from examined seabass and seabream, including *V. parahaemolyticus* (8.3%), *V. alginolyticus* (8.3%), *V. mimicus* (3.3%), *V. harveyi* (5%) and *V. vulnificus* (1.6%). *Vibrio cholera* was not detected in the studied samples. The examined seabass fish were more infected with *V. parahaemolyticus* than the examined seabream fish which may be due to the hygienic state of each fish source.

Similarly, Saad et al. (2015) isolated *V. parahaemolyticus* (10%), *V. fluvialis*, *V. vulnificus*, *V. alginolyticus*, *V. mimicus*, and *V. damsela* from *Tilapia nilotica* and *Mugil Cephalus*. Hemmat et al. (2018) isolated *V. parahaemolyticus* (12%), *V. mimicus*, *V. alginolyticus*, *V. cholera*, *V. vulnificus*, and *V. fluvialis* from *Oreochromis niloticus*, *Mugil Cephalus*, shrimp and crab. Raissy et al. (2015) isolated *V. harveyi* that was the most frequent species isolated, followed by *V. parahaemolyticus* (3.5%), *V. mimicus*, *V. vulnificus*, and *V. alginolyticus* from some marine fish and shrimps. Fri et al. (2017) isolated *V. fluvialis*, *Vibrio vulnificus*, and *V. parahaemolyticus* (5.45%) from dusky kop fish and sea water. Pal and Das (2010) isolated *Vibrio parahaemolyticus* with a high prevalence (35%) from shrimp, prawn, bhetki, pamfret and hilsa. According to the Egyptian Organization for Standardization and Quality Control (EOSQC, 2005), any seafood products must be free from *V. parahaemolyticus*.

As shown in table 3 and figure 1, all examined *Vibrio parahaemolyticus* isolates were positive for *toxR* gene. This result support finding of Yung et al. (1999); Pal and Das (2010), who reported that *toxR*-targeted PCR protocol can be used for *V. parahaemolyticus* detection. Also, all examined *Vibrio parahaemolyticus* isolates were positive for *aadAI* gene (Figure 2). Taviani et al. (2008) stated that *aadAI* gene is responsible for antibiotic resistance against aminoglycoside group including streptomycin in *Vibrio* spp. isolates from shellfish and other marine fish.

Pathogenicity of *V. parahaemolyticus* is conferred either by *tdh*, and/or *trh* (Yamaichi et al., 1999). As shown in table 3, all examined *V. parahaemolyticus* isolates were positive for *trh* gene (figure 3), and 80% were positive for *tdh* gene (Figure 4). The results did not match with that reported by Rojas et al. (2011) who detected *tdh* gene in 10.5% of *V. parahaemolyticus* isolates, while *trh* gene was not found. Also, Pal and Das (2010) recorded *tdh* gene in 35% of *V. parahaemolyticus* isolated from fish samples while *trh* gene was found only in 1.7% of *V. parahaemolyticus* isolates. Wang et al. (2017) recorded the virulence genes; *tdh* and *trh* with 87.9% and 3.7% of examined *V. parahaemolyticus* strains, respectively. Fri et al. (2017) recorded *trh* gene as 9.46% in examined *V. parahaemolyticus* strains, while Wong et al. (2000) recorded only one *V. parahaemolyticus* isolate (1.4%) harboring *trh* gene, but did not detect *tdh* gene among the examined *V. parahaemolyticus* isolates.

Antimicrobial susceptibility test showed that *V. parahaemolyticus* isolates were sensitive to ciprofloxacin, norfloxacin, cefotaxime, and chloramphenicol while they were resistant to ampicillin, erythromycin, streptomycin, and gentamycin (Table 4). These results indicate that the examined strains were resistant to most members of the aminoglycoside group, which may be due to the fact that *aadAI* gene was detected in all examined *V. parahaemolyticus* isolates. This result is nearly similar to that recorded by Rojas et al. (2011), who reported that *V. parahaemolyticus* had resistance to streptomycin and ampicillin with intermediate susceptibility to gentamicin.

**Table 2.** Prevalence of *Vibrio* species in examined seabass and seabream fish, Egypt.

<i>Vibrio</i> spp.	Number of positive samples		
	Seabass (n=30)	Seabream (n=30)	Total (%)
<i>V. parahaemolyticus</i>	1	4	5 (8.3)
<i>V. alginolyticus</i>	3	2	5 (8.3)
<i>V. mimicus</i>	0	2	2 (3.3)
<i>V. harveyi</i>	1	2	3 (5)
<i>V. vulnificus</i>	1	0	1 (1.6)
Total	6	10	16(26.6)


**Table 3.** Distribution of virulence genes among examined isolates of *Vibrio parahaemolyticus* isolated from seabass and seabream fish.

Sample No.	<i>toxR</i>	<i>tdh</i>	<i>trh</i>	<i>aadA1</i>
1	+	+	+	+
2	+	-	+	+
3	+	+	+	+
4	+	+	+	+
5	+	+	+	+


**Table 4.** Results of agar disc diffusion test of *Vibrio parahaemolyticus* isolated from marine fish

Antibiotic	Disc symbol & concentration (µg/disc)	Result
Norfloracin	Nor (10)	S
Erythromycin	E (15)	R
Ampicillin	AMP (10)	R
Amoxicillin + clavulanic acid	AMC (30)	S
Cefotaxime	CTX(30)	S
Doxycycline	DO (30)	R
Streptomycin	S(10)	R
Sulpamethazol + Trimethoprim	SXT(25)	R
Chloramphenicol	C (30)	S
Gentamycin	CN(10)	R
Ciprofloxacin	Cip (5 )	S

S: Sensitive R: Resistant




**Figure 1.** Agarose gel electrophoresis of PCR amplification of *toxR* gene (368 bp) of *Vibrio parahaemolyticus*. Lane L: 100-600 bp DNA Ladder. Neg.: Negative control. Pos.: Positive control. Lane 1-5: Positive samples.




**Figure 2.** Agarose gel electrophoresis of PCR amplification of *aadA1* gene (484 bp) of *Vibrio parahaemolyticus*. Lane L: 100-600 bp DNA Ladder. Neg.: Negative control. Pos.: Positive control. Lane 1-5: Positive samples.





**Figure 3.** Agarose gel electrophoresis of PCR amplification of *trh* gene (250 bp) of *Vibrio parahaemolyticus*. Lane L: 100-600 bp DNA Ladder. Neg.: Negative control. Pos.: Positive control. Lane 1-5: Positive samples.



**Figure 4.** Agarose gel electrophoresis of PCR amplification of *tdh* gene (373bp) of *Vibrio parahaemolyticus*. Lane L: 100-600 bp DNA Ladder. Neg.: Negative control. Pos.: Positive control. Lane 1,3,4 and 5: positive samples, Lane 2: negative sample.

## CONCLUSION

*Vibrio* spp. especially *V. parahaemolyticus*, *V. alginolyticus*, *V. mimicus*, and *V. vulnificus* are commonly isolated from seabass and seabream fish, which affects persons who consume improperly cooked or raw seafood. Most of these bacteria have antibiotic resistance genes that pose a great risk to human health; therefore, good hygienic measures should apply to avoid such infections.

## DECLARATIONS

### Acknowledgments

The authors would like to thank all member of Animal Health Research Institute, Egypt, for their kindly help during the investigation and paper preparation.

### Authors' contributions

All authors participated equally in study design, data collection, data analysis, writing, and approving the final manuscript.

### Competing interests

The authors declare that they have no competing interests.

## REFERENCES

- Angela D, Schulzea Abayomi O, Alabib Adele R, Sheldrake T and Miller KM (2006). Bacterial diversity in a marine hatchery: Balance between pathogenic and potentially probiotic bacterial strains. *Aquaculture*, 256: 50-73. DOI: <https://doi.org/10.1016/j.aquaculture.2006.02.008>
- Austin B and Austin DA (2007). *Bacterial fish pathogens, disease of farmed and wild fish*, 4th ed. Springer Praxis, Godalming. Available at: <https://www.springer.com/gp/book/9781402060687>
- Azwai SM, Alfallani EA, Abolghait SK, Garbaj AM, Naas HT, Moawad AA, Gammoudi FT, Rayes HM, Barbieri I and Eldaghayes IM (2016). Isolation and molecular identification of *Vibrio* spp. by sequencing of 16S RNA from seafood, meat and meat products in Libya. *Open Veterinary Journal*, 6(1): 36-43. DOI: <https://doi.org/10.4314/ovj.v6i1.6>
- Bej AK, Patterson DP, Brasher CW, Vickery MCL, Jones DD, and Kaysner CA (1999). Detection of total and hemolysin-producing *Vibrio parahaemolyticus* in shellfish using multiplex PCR amplification of *tlh*, *tdh*, and *trh*. *Journal of Microbial Methods*, 36: 215 – 225. DOI: [https://doi.org/10.1016/S0167-7012\(99\)00037-8](https://doi.org/10.1016/S0167-7012(99)00037-8)
- Broberg CA, Calder TJ and Orth K (2011). *Vibrio parahaemolyticus* cell biology and pathogenicity determinants. *Microbes and infection / Institute Pasteur*;13(12–13): 992-1001. DOI: <https://doi.org/10.1016/j.micinf.2011.06.013>
- Chao G, Jiao X, Zhou X, Yang Z, Huang J, and Pan Z (2009). Serodiversity, pandemic O3:K6 clone, molecular typing, and antibiotic susceptibility of food borne and clinical *Vibrio parahaemolyticus* isolates in Jiangsu, China. *Foodborne Pathogenic Disease*.6(8): 1021–1028. DOI: <https://doi.org/10.1089/fpd.2009.0295>
- Clinical and Laboratory Standards Institute (CLSI) (2012). Performance standards for antimicrobial disk susceptibility tests; approved standard, 11<sup>th</sup> ed (M02-A11). Clinical and Laboratory Standards Institute, Wayne, Pennsylvania. Available at: <http://www.facm.ucl.ac.be/intranet/CLSI/CLSI-M100S22>
- Dalsgaard A, Forslund A, Sandvang D, Arntzen L and Keddy K (2001). *Vibrio cholera* O1 outbreak isolates in Mozambique and South Africa in 1998 are multiple-drug resistant, contain the *STX* element and the *aadA2* gene located on class 1 integrons. *Journal of Antimicrobial Chemotherapy*, 48: 827–838. DOI: <https://doi.org/10.1093/jac/48.6.827>
- Deepanjali A, Sanath K and Karunasagar I (2005). Seasonal variation in abundance of total and pathogenic *Vibrio parahaemolyticus* bacteria in oysters along the Southwest coast of India. *Applied and Environmental Microbiology*, 71: 3575-3580. DOI: <https://doi.org/10.1128/AEM.71.7.3575-3580.2005>
- DePaola A, Nordstrom JL, Bowers JC, Wells JG and Cook DW (2003). Seasonal Abundance of Total and Pathogenic *V. parahaemolyticus* in Alabama Oysters. *Journal of Applied and Environmental Microbiology*, 69 (3): 1521 –1526. DOI: <https://doi.org/10.1128/AEM.69.3.1521-1526.2003>
- Egyptian Organization for Standardization and quality control (EOSQC) (2005). The Egyptian standard number 2005/3494. Available at: <https://www.unesco.org/egyptian-organization-standards-and-quality>
- Taviani E, Ceccarelli D, Lazaro N, Bani S, Cappuccinelli P, Colwell RR and Colombo MM (2008). Environmental *Vibrio* spp., isolated in Mozambique, contain a polymorphic group of integrative conjugative elements and class 1 integrons. *FEMS Microbiology Ecology*, 64: 45–55. DOI: <https://doi.org/10.1111/j.1574-6941.2008.00455.x>
- Qadri F, Chowdhury NR, Takeda Y and Nair BG (2005). *Vibrio parahaemolyticus*-Seafood Safety and Associations with Higher Organisms. *Oceans and Health: Pathogenes in the Marine Environment*. Edited by Belkin and Colwell, Springer, New York, 2005. DOI: <https://doi.org/10.1007/0-387-23709-711>
- Fri J, Ndip RN, Njom HA and Clarke AM (2017). Occurrence of virulence genes associated with human pathogenic *Vibrio* isolated from two commercial Dusky Kob (*Argyrosomus japonicus*) farms and Kareiga Estuary in the Eastern Cape Province, South Africa. *International Journal of Environmental Research Public Health*, 14: 1111. DOI: <https://doi.org/10.3390/ijerph14101111>
- Friesema IH, De Jong AE, Fitz James IA, Heck ME, Van den Kerkhof, JH and Notermans DW (2012). Outbreak of *Salmonella thompson* in the Netherlands since July 2012. *Eurosurveillance*, 17: 1-4. Available at: <http://eurosurveillance.org/images/dynamic/EE/V17N43/art20303.pdf>
- Han H, Li F, Yan W, Guo Y, Li N, and Liu X (2015):. Temporal and spatial variation in the abundance of total and pathogenic *Vibrio parahaemolyticus* in shellfish in China. *PLoS One*, 10(6): e0130302. DOI: <https://doi.org/10.1371/journal.pone.0130302>
- Hasrimi AN, Budiharjo A and Jannah SN (2018). Detection of *tlh* and *tdh* genes in *Vibrio parahaemolyticus* inhabiting farmed water ecosystem used for *L. Vannamei* aquaculture. *Journal of Physics: Conference Series*, 1025: 012058. DOI: <https://doi.org/10.1088/1742-6596/1025/1/012058>
- Hemmat M, Ibrahim, Reham A, Amin I, Nesreen Z, Eleiwa and Hanan RM Ghanaym (2018). *Vibrio* species in fish and shell fish . *Benha Veterinary Medical Journal*, 34 (2): 246-254. DOI: <https://doi.org/10.21608/bvmj.2018.29435>
- Hiyoshi H, Kodama T, Iida T, and Honda T (2010). Contribution of *Vibrio parahaemolyticus* virulence factors to cytotoxicity, enterotoxicity, and lethality in mice. *Infection and Immunity*, 78: 1772–1780. DOI: <https://doi.org/10.1128/IAI.01051-09>
- Hubbard TP, Chao MC, Abel S, Blondel CJ, zur Wiesched PA, Zhoue X, Davis BM and Waldor MK (2016). Genetic analysis of *Vibrio parahaemolyticus* intestinal colonization, *Proceedings of the National Academy of Sciences*, 113 (22): 6283–6288. DOI: <https://doi.org/10.1073/pnas.1601718113>
- Huss HH (1997). Control of indigenous pathogenic bacteria in seafood. *Food Control*, 8(2): 91-98. DOI: [https://doi.org/10.1016/S0956-7135\(96\)00079-5](https://doi.org/10.1016/S0956-7135(96)00079-5)
- International Commission on Microbiological Specifications for Foods (ICMSF) (1996). *Vibrio cholera*. In: *Microorganisms in Foods 5. Characteristics of Microbial Pathogens*. London: Blackie Academic & Professional, pp. 414-425. Available at: <https://www.iso.org/organization/9260.html>
- International Organization for Standardization/technical specification (ISO/TS21872-1) (2007). Specifies a horizontal method for the detection of the two main pathogenic *Vibrio* species causing intestinal illness in humans: *V. parahaemolyticus* and *V. cholera*. Available at: <https://www.iso.org/standard/38278.html>
- International Organization for Standardization/technical specification (ISO/TS21872-2) (2007). Specifies a horizontal method for the detection of the enteropathogenic *Vibrio* species, causing illness in or via the intestinal tract , other than *V. parahaemolyticus* and *V. cholera*. Include *V. fluvialis* , *V. mimicus* and *V. vulnificus*. Available at: <https://www.iso.org/standard/38279.html>
- Iwahori J, Yamamoto A, Suzuki H, Yamamoto T, Tsutsui T and Motoyama K (2010). Quantitative risk assessment of *Vibrio parahaemolyticus* in finfish: a model of raw horse mackerel consumption in Japan. *Risk Analysis*, 30(12): 1817–1832. DOI: <https://doi.org/10.1111/j.1539-6924.2010.01444.x>
- Kim YB, Okuda J, Matsumoto C, Takahashi T, Hashimoto S and Nishibuchi M (1999). Identification of *Vibrio parahaemolyticus* Strains at the Species Level by PCR Targeted to the *toxR* Gene. *Journal of Clinical Microbiology*, 37 (4): 1173-1177. DOI: <https://doi.org/10.1128/JCM.37.4.1173-1177.1999>
- Kishishita M, Matsuoka N, Kumagai K, Yamasaki S, Takeda Y and Nishibuchi M (1992). Sequence variation in the thermostable direct hemolysin related hemolysin (*trh*) gene of *Vibrio parahaemolyticus*. *Applied and Environmental Microbiology*, 58: 2449–2457. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC195802/>

- Liao Y, Li Y, Wu S, Mou J, Xu Z and Cui R (2015). Risk Factors for *Vibrio parahaemolyticus* Infection in a Southern Coastal Region of China. *Food borne Pathogenic Disease*, 12(11): 881–886. DOI: <https://doi.org/10.1089/fpd.2015.1988>
- Rojas MVR, Mattè MH, Drorpa M, da Silva ML and Mattè GR (2011). Characterization of *Vibrio parahaemolyticus* isolated from Oysters and Mussels in SÃO PAULO, BRAZIL. *Rev. Inst. Med. Trop. Sao Paulo*. 53(4): 201-205. DOI: <https://doi.org/10.1590/S0036-46652011000400005>
- Abdel-Azeem MW, Attaya A, Manal I, El-Barbary and Sultan S (2016). Isolation and molecular detection of pathogenic *Vibrio* species among economic fish from Red Sea in Egypt. *British Microbiology Research Journal*, 12(6): 1-8. DOI: <https://doi.org/10.9734/BMRJ/2016/42016>
- Mustapha S, Mustapha EM and Nozha C (2013). *Vibrio Alginolyticus*: An Emerging Pathogen of Food borne Diseases. *International Journal of Science and Technology*, 2 (4). 302-309.
- Osawa R, Okitsu T, Morozumi H and Yamai S (1996). Occurrence of urease-positive *Vibrio parahaemolyticus* in Kanagawa, Japan, with specific reference to presence of thermostable direct hemolysin (TDH) and the TDH-related hemolysin genes. *Applied and Environmental Microbiology*, 62:725–727. Available at” <https://www.ncbi.nlm.nih.gov/pubmed/8593076>
- Pal D and Das N (2010). Isolation, identification and molecular characterization of *Vibrio parahaemolyticus* from fish samples in Kolkata; European Review for Medical and Pharmaceutical Sciences.14: 545-549. Available at: <https://www.europeanreview.org/article/778>
- Raissy M, Rahimi E, Azargun R, Moumeni M, Rashedi M and Sohrabi HR (2015). Molecular detection of *Vibrio* spp. in fish and shrimp from the Persian Gulf. *Journal of Food Biosciences and Technology*, 5(2): 49-52. Available at: [http://jfbt.srbiau.ac.ir/article\\_7003.html](http://jfbt.srbiau.ac.ir/article_7003.html)
- Randall LP, Cooles SW, Osborn MK, Piddock LJV and Woodward MJ (2004). Antibiotic resistance genes, integrons and multiple antibiotic resistance in thirty-five serotypes of *Salmonella enteric* isolated from humans and animals in the UK. *Journal of Antimicrobial Chemotherapy*,53: 208–216. DOI: <https://doi.org/10.1093/jac/dkh070>.
- Saad MS, Samir MM and Abd El Maksod HES (2015). Incidence of *Vibrio* species in fish with special emphasis on the effect of heat Treatments. *Benha Veterinary Medical Journal*, 29 (1): 38-44. Available at: <http://www.bvmj.bu.edu.eg>
- Shirai H, Ito H, Hirayama T, Nakabayashi Y, Kumagai K, Takeda Y, and Nishibuchi M (1990). Molecular epidemiologic evidence for association of thermostable direct hemolysin (TDH) and TDH-related hemolysin of *Vibrio parahaemolyticus* with gastroenteritis. *Infection and Immunity*, 58: 3568–3573. Available at” <https://pubmed.ncbi.nlm.nih.gov/2228229/>
- Su YC and Liu C (2017). *Vibrio parahaemolyticus*: a concern of seafood safety. *Food Microbiology*, 24(6): 549–558. DOI: <https://doi.org/10.1016/j.fm.2007.01.005>
- Sujeewa AKW, Norrakiah AS and Laina M (2009). Prevalence of toxic genes of *Vibrio parahaemolyticus* in shrimps (*Penaeus monodon*) and culture environment. *International Food Research Journal*, 16: 89-95. Available at: <http://www.ifrj.upm.edu.my/20IFRJ-2008-Sujeewa>
- Vongxay K, Pan Z, Zhang X, Wang S, Cheng S and Mei L (2008). Occurrence of pandemic clones of *Vibrio parahaemolyticus* isolates from seafood and clinical samples in a Chinese coastal province. *Foodborne Pathogenic Disease*, 5(2):127–134. DOI: <https://doi.org/10.1089/fpd.2007.0045>
- Wang H, Tang X, Su Y-C, Chen J and Yan J (2017). Characterization of clinical *Vibrio parahaemolyticus* strains in Zhoushan, China, from 2013 to 2014. *PLoS One*, 12(7): e0180335. DOI: <https://doi.org/10.1371/journal.pone.0180335>
- Wang R, Zhong Y, Gu X, Yuan J, Saeed A Fand and Wang S (2015). The pathogenesis, detection and prevention of *Vibrio parahaemolyticus*. *Frontiers in Microbiology*, 6: 144. DOI: <https://doi.org/10.3389/fmicb.2015.00144>
- Wong HC, Lui SH, Wang TK, Lee CL, Chiou CS, Lui DP, Nishibuchi M and Lee BK (2000). Characterization of *Vibrio parahaemolyticus* O3:K6 from Asia. *Applied and Environmental Microbiology*, 66: 3981–3986. DOI: <https://doi.org/10.1128/AEM.66.9.3981-3986.2000>
- Wu Y, Wen J, Ma Y, Ma X and Chen Y (2014). Epidemiology of food borne disease outbreaks caused by *Vibrio parahaemolyticus*, China, 2003–2008. *Food Control*, 46: 197–202. DOI: <https://doi.org/10.1016/j.foodcont.2014.05.023>
- Yamaichi Y, Iida T, Park KS, Yamamoto K and Honda T (1999). Physical and genetic map of the genome of *Vibrio parahaemolyticus*: Presence of two chromosomes in *Vibrio* species. *Molecular Microbiology*, 31: 1513–1521. DOI: <https://doi.org/10.1046/j.1365-2958.1999.01296.x>.
- Yoon KS, Min KJ, Jung YJ, Kwon KY, Lee JK and Oh SW (2008). A model of the effect of temperature on the growth of pathogenic and nonpathogenic *Vibrio parahaemolyticus* isolated from oysters in Korea. *Food Microbiology*, 25(5): 635–641. DOI: <https://doi.org/10.1016/j.fm.2008.04.007>.
- Yung BK, Jun O, Chiho M, Naoki T, Saouri H, Mitsuaki IN, Hashimoto S and Nishibuchi M (1999). Identification of *Vibrio parahaemolyticus* Strains at the Species Level by PCR Targeted to the *toxR* Gene. *Journal of Clinical Microbiology*, 37 (4): 1173–1177. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC88669/>
- Zulkifli Y, Alitheen NB, Son R, Raha A R, Samuel L, Yeap SK and Nishibuchi M (2009). Random amplified polymorphic DNA-PCR and ERIC PCR analysis on *Vibrio parahaemolyticus* isolated from cockles in Padang, Indonesia. *International Food Research Journal*, 16: 141-150.