



Coccidiosis: A Parasitic Disease of Significant Importance in Rabbits

Wafaa A. Abd El-Ghany

Poultry Diseases Department, Faculty of Veterinary Medicine, Cairo University, 12211 Giza, Egypt

Corresponding author's Email: wafaa.ghany@yahoo.com; ORCID: 0000-0003-1686-3831

REVIEW ARTICLE
p11: S232245682000060-10
Received: 04 Nov 2020
Accepted: 20 Dec 2020

ABSTRACT

Rabbits are considered an important and healthy source of animal protein all over the world. They are susceptible to important diseases that can reduce their productivity, causing severe economic losses. Coccidiosis is one of the important protozoan diseases caused by *Eimeria* species. Rabbits are highly susceptible to coccidiosis, especially after weaning time. Coccidiosis in rabbits has two forms, namely hepatic and intestinal. Affected animals indicated the symptoms of diarrhea, reduced appetite, dehydration, and weight loss as well as liver and intestinal lesions. Diagnosis is based on the detection of the infective stages of the protozoan in feces or affected tissues. Prevention and control are achieved by adopting hygienic measures and using different anticoccidial drugs. The use of natural alternatives for the prophylaxis of coccidiosis in rabbits indicated promising results. Vaccine production trials are still under investigation. Accordingly, this review article aims to shed light on coccidiosis in rabbits considering pathology, diagnosis, and control.

Keywords: *Eimeria*, Intestine, Liver, Rabbits, Treatment

INTRODUCTION

Rabbits (*Oryctolagus cuniculus*) are regarded as a potential source of animal protein for human consumption. The meat of rabbits is recommended for human consumption more than other sources of proteins due to its high nutritious protein, calcium, phosphorus, and linoleic acid, with low fat and cholesterol contents (Nistor et al., 2013). In addition to the commercial use of rabbits, they can be used for wool production and in medical research as laboratory animals, and they are raised as pets for hobby purposes (Al-Mathal, 2008).

Rabbits are susceptible to dangerous viral, bacterial, and parasitic diseases that drastically affect their production. Coccidiosis or eimeriosis is one of the most important and widely distributed parasitic diseases in rabbits (Grès et al., 2003; El-Shahawi et al., 2012; Okumu et al., 2014; Yin et al., 2016; Bachene et al., 2019; Hamid et al., 2019). The disease affects the intestine, liver, and bile duct of the animal and is associated with digestive disorders (Manjunatha et al., 2019). Coccidiosis is a highly contagious disease that has negative impacts on the domestic rabbit industry worldwide (Pakandl, 2009; Geru et al., 2017). It seriously impairs growth performance parameters, causes high morbidity and mortality rates (Abdel-Megeed et al., 2005), and reduces the carcass weight by more than 23% (Barriga and Arnoni, 1981). Coccidiosis causes annual losses in the rabbit industry in the USA amounting to 127 million dollars, and similar losses may happen worldwide (Chapman, 2009). The prevalence of coccidiosis in rabbits is varied and influenced by geographical location, season, as well as management factors like housing and rearing conditions, and the use of preventive coccidiostats (Chowdhury and Fraser, 2008). Furthermore, the prevalence of coccidiosis in weaned rabbits is higher than that in growing and adult ones (El-Ghoneimy and El-Shahawy, 2017). Duszynski and Couch (2013) stated that the rate of coccidial infection ranged from 64% to 100% all over the world.

The disease is caused by an intracellular ubiquitous protozoan parasite of the genus *Eimeria* (Jing et al., 2012; Mäkitaipale et al., 2017). Mixed infection with more than one *Eimeria* spp. is common (Jithendran and Bhat, 1996). There are 15 *Eimeria* spp. that affect the intestinal tract and one species (*E. stidea*) affecting the liver and bile duct (Li and Ooi, 2009). *E. intestinalis*, *E. magna*, *E. piriformis*, *E. perforans*, *E. media*, *E. agnotsa*, *E. exigua*, *E. flavescens*, *E. irrisidua*, *E. coecicola*, *E. vej dovskyi*, *E. roobroucki*, *E. oryctolagi*, *E. nagpurensis*, and *E. matsubayashi* invade the small intestine (Soulsby, 1968). The highly pathogenic *Eimeria* spp. in rabbits are *E. intestinalis* and *E. flavescens*, the moderately pathogenic spp. are *E. magna*, *E. irrisidua*, and *E. piriformis*, and the low pathogenic or nonpathogenic spp. are *E. exigua*, *E. media*, *E. coecicola*, and *E. perforans* (Jithendran, 1995). The rapid spread of infection, due to the direct and short life cycle of the parasite, is characteristic of coccidiosis (García-Rubio et al., 2017). The severity of coccidial infection depends on the number of ingested oocysts, age, and the immune status of the animal (Miller and Fowler, 2003). In the early stage of infection with coccidiosis, sudden diarrhea, or death may occur without any clinical signs.

Clinical infection is characterized by apathy, diarrhea, dehydration, reduced appetite, and weight loss resulting in death (Ogolla et al., 2018).

Prevention and control of coccidiosis in rabbits are achieved through careful management, enhancing rabbits' immunity, vaccination, and using synthetic anticoccidial drugs. Housing in large, dry, sunny, and disinfected rooms is very essential to fight coccidial infection (Pilarczyk et al., 2020). Although synthetic anticoccidials were effective, their hazardous use in the field resulted in some drawbacks. Therefore, there is a shift toward using novel approaches that pose a minimal risk to human or animal health. Different kinds of herbal extracts (Indrasanti et al., 2017; Sorour et al., 2018; Rivero-Perez et al., 2019) and acidifiers (Shkromada et al., 2019) showed successful preliminary results in treating coccidiosis in rabbits. Efficient vaccines are very important for the prevention of rabbit coccidiosis (Song et al., 2017), but so far, no vaccine is available for rabbits.

Given the above information, the aim of the present review is to shed light on coccidiosis in rabbits considering pathology, diagnosis, and control.

Parasite life cycle

Infection usually occurs through the ingestion of contaminated feed and water containing sporulated oocysts (i.e., infective oocysts consist of four sporocysts, each containing two sporozoites). Upon entering the gut, due to the effects of gastric and pancreatic juices, the oocysts walls rupture and the sporozoites invade the intestinal epithelial cells and then pass via the mesenteric lymph nodes and hepatic portal circulation to the liver where they enter the epithelial cells of the bile duct becoming trophozoites and then schizonts. The recognition and invasion processes may occur via the action of sugar residues in *Eimeria* sporozoites (John et al., 1999), followed by the release of merozoites (merogony stage) that form four generations in the asexual stage of the parasite. Lastly, male (micro) and female (macro) gametocytes combine sexually (gametogony stage) to form zygotes that develop into non-sporulated oocysts. The non-sporulated oocysts pass in the bile and are shed in the feces of the infected rabbits about 18 days after infection. Under favorable environmental conditions, sporulated oocysts are formed within three days (Gardiner et al., 1998).

Susceptibility

All breeds of domestic are highly susceptible to coccidiosis, and those of 1-4 months old more susceptible than adults (González-Redondo et al., 2008; Papeschi et al., 2013; Bachene et al., 2018). Suckling rabbits cannot be infected with coccidiosis before three weeks of age. It has been found that the production of *E. flavescens* and *E. intestinalis* oocysts in suckling rabbits increases with age (Pakandl and Hlášková, 2007). More susceptibility in young rabbits may be due to reduced immunity resulting from weaning stress, feeding and reproductive status (Drouet-viard et al., 1997a; Al-Mathal, 2008). Also, it was found that female rabbits were more infected than males (Faraj, 2017).

Types of rabbit coccidiosis

Hepatic coccidiosis

Hepatic coccidiosis involves the liver and bile ducts of rabbits causing cirrhosis and cholestasis (Singla et al., 2000). Hepatic coccidiosis is associated with severe economic losses in rabbitries and is caused by *E. stiedae* that is one of the most pathogenic species of *Eimeria* (Xin et al., 2016; Al-Tae and Al-Zubaidi, 2017). The oocysts of this species of *Eimeria* were first detected in the bile of infected rabbits in 1674 (Duszynski and Couch, 2013). The most adverse effects of hepatic coccidiosis are reduced growth and feed utilization and increased mortality rate in young rabbits (Hanptman et al., 2001). Like other *Eimeria* spp., *E. stiedae* penetrates the intestinal wall and migrates to the bile ducts where it reproduces (Kraus et al., 1984). Hepatic coccidiosis is mostly chronic and subclinical infection specifically in adults who are carriers and sources of infection (Barriga and Arnoni, 1981; Al-Mathal, 2008; Pakandl, 2009). Some of the affected animals displayed symptoms of anorexia, polydipsia, brown watery diarrhea, dehydration, icteric membranes, poor feed conversion, growth retardation, wasting of the back and hindquarters, coarse hair, abdominal distension, and even death especially in young rabbits with severe infection (Erdogmus and Eroksuz, 2006; Lakshmanan et al., 2011; Al-Saeed et al., 2017). High mortality results from high doses of oocysts, while morbidity results from diarrhea and reduction in body weight (Renaux et al., 2003). Several studies have described the post-mortem findings of hepatic coccidiosis. On gross examination, enlarged and cirrhotic liver with multiple distributed whitish nodules containing creamy thick exudate, and distended gall bladder has been reported. Moreover, histologic findings included cholangitis, bile duct hyperplasia, hemorrhage, dilatation and congestion of the central veins as well as necrosis of hepatocytes and hepatic fibrosis (Sanyal and Sharma, 1990; Cam et al., 2008; AL-Naimi et al., 2012; Sorour et al., 2018).

Intestinal coccidiosis

There are about 15 species of *Eimeria* that are known to induce pathology in the intestine of rabbits causing intestinal coccidiosis. The most common *Eimeria* spp. that cause intestinal coccidiosis in rabbits are *E. magna*, *E.*

irresidua, *E. media*, and *E. perforans* (Fox, 1984). The parasite colonizes distinct parts of the intestine and the mucosa at different depths (Pakandl, 2009). Intestinal coccidiosis often is observed in rabbits aged from six weeks to five months old. Older animals can acquire immunity after recovery and become carriers (Kulisic et al., 2006). This type of coccidial infection results in atrophy of the intestinal villi, malabsorption of nutrients, hypoproteinemia, electrolyte imbalance, dehydration, anemia (Dakshinkar and Dharmadhikari, 1985; Hana et al., 2011) and is manifested by diarrhea, weight loss, and mortality (Lebas et al., 1986). Affected rabbits suffer from weakness, gnashing teeth, dirty anus, weight loss, soft to watery hemorrhagic diarrhea, dehydration, and thirst (Fioramonti et al., 1982). Inflammation and edema in the ileum and jejunum associated with mucosal bleeding and ulcerations were observed (Coudert et al., 1995; Oncel et al., 2011). There are differences in pathogenicity among intestinal coccidial spp. (Jithendran, 1995). These differences in pathogenicity can be determined through the experimental infection of animals with different *Eimeria* species.

Laboratory diagnosis

Laboratory diagnosis of hepatic and intestinal coccidiosis depends on the analysis of feces of suspected rabbits. Microscopic identification of *Eimeria* spp. oocysts through the fecal analysis of suspected animals is very important (Pakandl et al., 2008). Developmental stages of *E. stiedae* have been detected in stained impression smears from the liver (Al-Rukibat et al., 2001; Sivajothi et al., 2016). Histopathological examination of the liver tissues, bile duct, or intestine is also used for the detection of different developmental stages of the parasite (Sivajothi et al., 2016). The oocyte detection site under a microscope is a guide to determining *Eimeria* species. Immuno-diagnosis of *E. stiedae* was investigated in previous studies (Zayed and Kutkat, 1998; Kandil et al., 2000). Serological diagnosis of *E. stiedae* using ELISA was reported (Abu-El-Ezz et al., 2010; Wei et al., 2020). Identification of *Eimeria* spp. using molecular assays such as multiplex PCR assay was reported (Oliveira et al., 2011; Yan et al., 2013). Hassan et al. (2015) detected *E. stiedae* schizonts using PCR 12 days after the experimental infection of rabbits, and this occurred prior to the development of lesions or shedding of the oocysts in feces.

It should be noted that the presence of oocysts in fecal samples does not confirm the presence of clinical disease. It has been reported that rabbits with high-intensity infection showed no clinical symptoms of coccidiosis (Pilarczyk et al., 2020). The induction of symptoms may depend on the virulence and pathogenicity of the infecting *Eimeria* spp. (Pakandl, 2009).

Prevention and treatment

Management practices

Coccidiosis in rabbits is aggravated by poor hygienic conditions and high stocking densities that encourage the spread of protozoa (González-Redondo et al., 2008). Rabbits raised in groups are more affected than those kept alone (Sharma et al., 2016). Accordingly, the first steps for preventing the occurrence and spread of coccidiosis in a rabbitry are proper hygiene and husbandry practices as well as strict biosecurity measures (Pakandl et al., 2008; Scholaut et al., 2013). Control of coccidial infection using common disinfectants is difficult as oocysts have a remarkable ability to survive under exogenous environmental conditions (Chapman et al., 2013).

Anticoccidial drugs

Prevention of coccidiosis in rabbits using coccidiostats is regulated by Regulation (EC) No 1831/2003 of the European Parliament and of the Council on additives for use in animal nutrition. Globally, synthetic anticoccidial drugs, either ionophores or synthetic chemicals, remain the mainstream pharmaceuticals that are used for the control of rabbit coccidiosis (Pakandl, 2009). Anticoccidial drugs should be broad-spectrum, highly effective with a good therapeutic index, and easily administered for short time. Coccidiostats in rabbits prevent the developmental stages (schizogony and gamogony stages) of the parasite inside the host. Coccidiostats are usually added to the feed of animals; however, coccidiocidal drugs are added to water. It is preferable to prevent coccidial infection before its occurrence as the treatment is usually not very successful when clinical signs of coccidiosis appear (Pakandl, 2009). A previous study by Peeters et al. (1981) showed that concordant infection with hepatic and intestinal *Eimeria* spp. could be treated with narasin, while hepatic coccidiosis could be prevented by clodolol/methylbenzoquate, robenidine, and salinomycin (Peeters et al., 1982). Several reports recommended the use of toltrazuril for either prevention or treatment of coccidiosis in rabbits (Mikhail et al., 1981; Redrobe et al., 2010; Qamar et al., 2013; El-Ghoneimy and El-Shahawy, 2017). In a study by Vereecken et al. (2012), treatment with diclazuril, salinomycin, and robenidine showed significant improvement in both growth performance and parasitological parameters in infected rabbits compared to non-treated animals. However, using 25 ppm toltrazuril/liter of drinking water for two days was successful in treating the clinical hepatic coccidiosis (Singla et al., 2000), increasing body weight and lowering the mortality rate in rabbits infected with mixed intestinal and hepatic *Eimeria* spp. compared to control animals (Balicka-Ramisz et al., 2014). Treatment with toltrazuril immediately reduced signs and oocysts shedding, allowing the development of immunity against reinfection (Peeters and

Greeroms, 1986). Treatment using diclazuril is recommended for eliminating oocyst shedding in rabbits worldwide (Vanparijs et al., 1989a, b; Peeters and Geeroms, 1989; Polozowski, 1993; Vereecken et al., 2012). Diclazuril and sulfachloropyridazine were efficacious in treating rabbit coccidiosis (Ogolla et al., 2018). Studies presented variable results following the prophylactic and therapeutic use of sulphonamides in treating coccidiosis (Joyner et al., 1983; Polozowski, 1993; Redrobe et al., 2010; Qamar et al., 2013). Kolabskii et al. (1973) and Ogolla et al. (2018) reported the effectiveness of sulphachloropyridazine in controlling clinical coccidiosis in rabbits. Successful control of hepatic coccidiosis using sulphaquinoxaline was also reported (Magray et al., 2010). Amprosol, bifuran, and sulpha-based drugs have been used for the prevention of rabbit coccidiosis (Bhat et al., 2010). Trimethoprim-sulfamethoxazole showed moderate to satisfactory efficacious results in the treatment of field infection (Ogolla et al., 2018). It has been documented that sulfonamides protected against experimental *E. magna* and *E. media* infections better than colistin and trimethoprim (Bachene et al., 2019). Amprolium could not treat intestinal and hepatic coccidiosis (Laha et al., 1999; Ogolla et al., 2018). On the other hand, superior effects have been demonstrated following the prophylactic use of amprolium in intestinal coccidiosis especially when applied concurrently with other anticoccidials (Qamar et al., 2013; Laha et al., 2015; El-Ghoneimy and El-Shahawy, 2017). Some anticoccidials such as sulphonamides, salinomycin, and robenidine were toxic for pregnant does and kids, while few others used for poultry were recommended for rabbits (Ogolla et al., 2018).

Alternatives to anticoccidial drugs

Anticoccidial drugs are relatively inexpensive and showed successful results. However, increase in consumer demand for the production of organic products, the potential development of resistant strains of parasites toward drugs (Pakandl, 2009), and the presence of antibiotic residues in meat created a potential need for searching for natural and safe alternatives to anticoccidial chemicals. Hence, several studies investigated the effects of natural alternatives such as sulfur and sulfates (including copper sulfate), tannic acid, bismuth compounds, thymol, camphor, alum, volatile oils, and garlic, oregano, sage, caraway, cinnamon, basil, and rosemary on rabbit coccidiosis (Kowalska et al., 2012). Oral prophylactic administration of garlic was effective in the amelioration of *E. stiedae* infection (Toulah and Al-Raw, 2007; Abu-Akkada et al., 2010; Indrasanti et al., 2017). Adding of a mixture of garlic oil and oregano to rabbit feed can also help in disease prevention (Kowalska et al., 2012; Nosal et al., 2014). Other plant extracts such as banana stem extract could decrease the number of *E. stiedae* oocysts *in vitro* (Indrasanti et al., 2015). However, Matekaire et al. (2005) stated that both banana root (*Musa paradisiaca*) and sulphadimidine sodium treatments caused a significant decrease in oocyst excretion in rabbits. Neem extract improved body weight and liver function in experimentally infected rabbits with *E. stiedae*, and these results were similar to those in the toltrazuril treated group (Ahmed et al., 2014). Cervantes-Valencia et al. (2015) demonstrated that the hydroalcoholic extract of *Curcuma longa* at doses of 25 and 40 mg/kg body weight reduced the fecal *Eimeria* sp. excretion in naturally infected rabbits. Artemisinin liquid extract, cinnamon, and clove essential oils could also be used for protection against *E. stiedae* infection in rabbits compared to toltrazuril (Sorour et al., 2018). Furthermore, it has been demonstrated that 25 and 50 mg *Salix babylonica* hydroalcoholic extract per kg of body weight successfully reduced the intestinal *Eimeria* oocyst count per gram of feces in rabbits (Rivero-Perez et al., 2019). The effects of herbal antioxidants (*Psidium guajava*) as inhibitors of *E. intestinalis*, *E. magna*, *E. flavescens*, and *E. stiedae* sporozoites were evaluated *in vitro*, and preliminary results showed that these herbs were the best substitutes to chemical anticoccidials (Cedric et al., 2017). A dose of 20 g *Calotropis procera* dried leaves powder per kg pelleted ration of rabbits reduced the number of oocysts in feces with the absence of alterations in the intestine and liver tissues of rabbits with mixed infections (Seddek et al., 2015). Aloe vera and liquid paraffin showed varied efficacy in the treatment of rabbit coccidiosis (Ogolla et al., 2017). Shkromada et al. (2019) proved that water treatment of rabbits with an acidifier concentrate (a mixture of benzoic, acetic, and propionic acids) reduced *Eimeria* invasion, increased body weight, and improved feed conversion. In addition, compared to anticoccidial drugs, this composition was not toxic, improved digestion processes, and inhibited the development of conventionally pathogenic microflora.

Vaccines

Vaccinations using oral or spray dispersion of precocious live lines of *E. magna* oocysts in the nest boxes (Drouet-Viard et al., 1997a, b; Licois, 2004), and vaccination against *E. magna* and *E. media* (Akpo et al., 2012) showed satisfactory preliminary results. Some efforts have been directed toward the production of recombinant vaccines against coccidiosis in rabbits (Hanada et al., 2003; Abdel-Megeed et al., 2005; Song et al., 2010). A recent trial for the preparation of a vaccine against hepatic coccidiosis was carried out. Using immuno-proteomic analysis of sporozoite proteins of *E. stiedae*, the immuno-reactive proteins were recognized by the sera of infected rabbits, which may be helpful for the production of a vaccine (Song et al., 2017). However, production of a vaccine on a commercial and field scale is still distant and limited, may be due to the high cost of production and the time required for the processes of

optimization, registration, safety assessment, and distribution to customers (Song et al., 2017). Vaccination is promising and still requires extensive research and development to be applied in the near future.

CONCLUSION

As coccidiosis is considered a very important parasitic disease in rabbits, future studies should focus on finding novel approaches for the prevention and control of such a significant threat.

DECLARATIONS

Competing interests

The author has no conflict of interest.

REFERENCES

- Abdel-Megeed KN, Abu El-Ezz NM and Abdel-Rahman EH (2005). Protective effect of *Eimeria stiedae* coproantigen against hepatic coccidiosis in rabbits. *Journal of the Egyptian Society of Parasitology*, 35: 581-595. Available at: <https://pubmed.ncbi.nlm.nih.gov/16083069/>
- Abu-Akkada SS, Samah SO and Ashmawy KI (2010). Garlic and hepatic coccidiosis: prophylaxis or treatment? *Tropical Animal Health and Production*, 4: 1337-1343. DOI: <https://www.doi.org/10.1007/s11250-010-9590-6>
- Abu-El-Ezz NMT, Megeed KNA, Mahdy OA and Hassan SE (2010). ELISA assessment in the diagnosis of hepatic coccidiosis in experimentally infected rabbits. *Global Veterinaria*, 9: 517-523. Available at: DOI: <https://www.dx.doi.org/10.5829/idosi.gv.2012.9.5.6620>
- Ahmed SE, Abdel Razek MSZ, Ramadan ME and Esmail EM (2014). Hepatic coccidiosis in rabbits and comparative study on treatment with herbal drug. *Zagazig Veterinary Journal*, 42: 41-50. DOI: <https://www.dx.doi.org/10.21608/zvjz.2014.59960>
- Akpo Y, Kpodekon MT, Djago Y, Licois D and Youssao IA (2012). Vaccination of rabbits against coccidiosis using precocious lines of *Eimeria magna* and *Eimeria media* in Benin. *Veterinary Parasitology*, 184: 73-76. DOI: <https://www.doi.org/10.1016/j.vetpar.2011.08.012>
- Al-Mathal IM (2008). Hepatic coccidiosis of the domestic rabbit *Oryctolagus cuniculus domesticus* L. in Saudi Arabia. *World Journal of Zoology*, 3: 30-35. Available at: <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.506.3784&rep=rep1&type=pdf>
- AL-Naimi RAS, Khalaf OH, Tano SY and Al-Tae EH (2012). Pathological study of hepatic coccidiosis in naturally infected rabbits. *AL-Qadisiya Journal of Veterinary Medical Science*, 11: 63-69. Available at: <https://www.iasj.net/iasj?func=fulltext&aid=80344>
- Al-Rukibat RK, Irizarry AR, Lacey JK, Kazacos KR, Storandt ST and DeNicola DB (2001). Impression smear of liver tissue from a rabbit. *Veterinary Clinical Pathology*, 30: 57-61. DOI: <https://www.doi.org/10.1111/j.1939-165X.2001.tb00259.x>
- Al-Saeed MH, Al Saeed AH and Jori MM (2017). Study of physiological and histological changes in rabbits induced with hepatic coccidiosis. *Journal University of Kerbala*, 15: 217-228. Available at: https://kj.uokerbala.edu.iq/article_123448.html
- Al-Tae MNK and Al-Zubaidi MTS (2017). Protection against *Eimeria stiedae* in rabbits by using sonicated sporulated oocyst vaccine. *Journal of Entomology and Zoology Studies*, 205: 579-585. Available at: <https://www.entomoljournal.com/archives/?year=2017&vol=5&issue=4&ArticleId=2110>
- Bachene MS, Temim S, Ainbaziz H and Bachene A (2019). Prevalence of rabbit coccidia in Medea Province, Algeria. *World's Veterinary Journal*, 9: 123-128. DOI: <https://www.dx.doi.org/10.36380/scil.2019.wvj16>
- Bachene MS, Temim S, Ainbaziz H, Bachene A and Suo X (2018). A vaccination trial with a precocious line of *Eimeria magna* in Algerian local rabbits *Oryctolagus cuniculus*. *Veterinary Parasitology*, 261: 73-76. DOI: <https://www.doi.org/10.1016/j.vetpar.2018.08.013>
- Balicka-Ramis A, Wróbel M and Adadyńska K (2014). Epidemiology and economic benefits of treating rabbits coccidiosis in small farms from West Pomerania province, Poland. *Annals of Parasitology*, 60: 247-251. PMID: 25706421. Available at: <https://pubmed.ncbi.nlm.nih.gov/25706421/>
- Barriga OO and Arnoni JV (1981). Pathophysiology of hepatic coccidiosis in rabbits. *Veterinary Parasitology*, 8: 201-210. DOI: [https://www.doi.org/10.1016/0304-4017\(81\)90051-0](https://www.doi.org/10.1016/0304-4017(81)90051-0)
- Bhat TK, Jithendran KP and Kurade NP (2010). Rabbit coccidiosis and its control: a review. *World Rabbit Science*, 4: 37-41. DOI: <https://www.doi.org/10.4995/wrs.1996.269>
- Cam Y, Alasever A, Eraslan G, Kibar M, Atalay O, Beyaz L, Inci A and Liman BC (2008). *Eimeria stiedae*: Experimental infection in rabbits and the effect of treatment with toltrazuril and ivermectin. *Experimental Parasitology*, 119: 164-171. DOI: <https://www.doi.org/10.1016/j.exppara.2008.01.005>
- Cedric Y, Payne VK, Nadia NAC, Kodjio N, Kollins E, Megwi L, Kuate JR and Mbida M (2017). *In vitro* anticoccidial, antioxidant activities and cytotoxicity of *Psidium guajava* extracts. *Journal of Parasitic Diseases: Diagnosis and Therapy*, 2: 14-24. Available at: <https://www.alliedacademies.org/articles/in-vitro-anticoccidial-antioxidant-activities-and-cytotoxicity-of-psidium-guajava-extracts-8865.html>
- Cervantes-Valencia ME, Alcalá-Canto Y, Salem AZM, Kholif AE, Ducoing-Watty AM, Bernad-Bernad MJ and Gutiérrez-Olvera C (2015). Influence of curcumin (*Curcuma longa*) as a natural anticoccidial alternative in adult rabbits: first results. *Italian Journal of Animal Science*, 14: 3838. DOI: <https://www.doi.org/10.4081/ijas.2015.3838>

- Chapman HD (2009). A landmark contribution to poultry science prophylactic control of coccidiosis in poultry. *Poultry Science*, 88: 813-815. DOI: <https://www.doi.org/10.3382/ps.2008-00316>
- Chapman HD, Barta JR, Blake D, Gruber A, Jenkins M, Smith NC, Suo X and Tomley FM (2013). A selective review of advances in coccidiosis research. *Advances of Parasitology*, 83: 93-171. DOI: <https://www.doi.org/10.1016/b978-0-12-407705-8.00002-1>
- Chowdhury AA and Fraser GC (2008). *Coccidia (Eimeria spp.) of domestic rabbits in New South Wales*. *Australian Veterinary Journal*, 86: 365-366. DOI: <https://www.doi.org/10.1111/j.1751-0813.2008.00340.x>
- Coudert P, Licois D, Provot F and Drouet-Viard F (1995). *Eimeria* species and strains of rabbits. In: Eckert J, Braun R, Shirley MW, Coudert P, Editors: Guidelines on techniques in coccidiosis research. COST 89/820: Biotechnology. European Commission, Luxembourg, 52-73.
- Dakshinkar NP and Dharmadhikari DN (1985). Haematological observations in intestinal coccidiosis during clinical outbreak. *Poultry Adviser*, 18: 55-56. Available at: <http://cehea.org/wp-content/uploads/2016/08/59-.pdf>
- Drouet-viard F, Coudert P, Licois P and Boivin M (1997a). Acquired protection of the rabbit (*Oryctolagus cuniculus*) against coccidiosis using a precocious line of *Eimeria magna*: effect of vaccine dose and age at vaccination. *Veterinary Parasitology*, 69: 197-201. DOI: [https://www.doi.org/10.1016/S0304-4017\(96\)01133-8](https://www.doi.org/10.1016/S0304-4017(96)01133-8)
- Drouet-viard F, Coudert P, Licois D and Boivin M (1997b). Vaccination against *Eimeria magna* coccidiosis using spray dispersion of precocious line oocysts in the nest box. *Veterinary Parasitology*, 70: 61-66. DOI: [https://www.doi.org/10.1016/S0304-4017\(96\)01134-X](https://www.doi.org/10.1016/S0304-4017(96)01134-X)
- Duszynski DW and Couch L (2013). *The Biology and Identification of the Coccidia (Apicomplexa) of Rabbits of the World*. San Diego, California, USA. Academic Press, pp. 4-5.
- El-Ghoneimy A and El-Shahawy I (2017). Evaluation of amprolium and toltrazuril efficacy in controlling natural intestinal rabbit coccidiosis. *Iranian Journal of Veterinary Research*, 18: 164-169. DOI: <https://www.ncbi.nlm.nih.gov/pubmed/29163644>
- El-Shahawi GA, El-Fayomi HM and Abdel-Haleem HM (2012). Coccidiosis of domestic rabbit (*Oryctolagus cuniculus*) in Egypt: Light microscopic study. *Parasitology Research*, 110: 251-258. DOI: <https://www.doi.org/10.1007/s00436-011-2479-0>
- Erdogmus ZS and Eroksuz Y (2006). DS hepatic coccidiosis in Angora rabbits. *Journal of Animal and Veterinary Advances*, 5: 462-463. DOI: <https://medwelljournals.com/abstract/?doi=javaa.2006.462.463>
- Faraj AA (2017). Molecular diagnosis of natural infection with *Eimeria stiedae* in domestic rabbits in Baghdad city-Iraq. *Journal of Entomology and Zoological Studies*, 5: 1438-1440. Available at: <https://www.entomoljournal.com/archives/?year=2017&vol=5&issue=3&ArticleId=1974>
- Fioramonti J, Sorraing JM, Licois D and Bueno L (1982). Intestinal motor and transit disturbances associated with experimental coccidiosis (*Eimeria magna*) in the rabbit. *Annals of Veterinary Research*, 12: 413-420. Available at: <https://pubmed.ncbi.nlm.nih.gov/7137853/>
- Fox JG (1984). *Laboratory Animal Medicine*. Academic Press Inc.: New York, 1984.
- García-Rubio V, Bautista-Gómez LG, Martínez-Castañeda JS and Romero-Núñez C (2017). Multicausal etiology of the enteric syndrome in rabbits from Mexico. *Revista Argentina de Microbiología*, 49: 132-138. DOI: <https://www.doi.org/10.1016/j.ram.2017.03.001>
- Gardiner GH, Fayer R and Dubey JP (1998). Apicomplexa. In: *An Atlas of Protozoan Parasites in Animal Tissues*, 2nd edition, Armed Forces Institute of Pathology, Washington, DC., 20-30.
- Geru T, Wang Y, Li C, Gu X, Cui P, Fang S, Suo X and Liu X (2017). High pathogenicity and strong immunogenicity of a Chinese isolate of *Eimeria magna* Pérard, 1925. *Parasitology International*, 66: 207-209. DOI: <https://www.doi.org/10.1016/j.parint.2017.01.014>
- González-Redondo P, Finzi A, Negretti P and Micci M (2008). Incidence of coccidiosis in different rabbit keeping systems. *Brazilian Archive of Veterinary Medicine and Animal Science*, 60: 1267-1270. DOI: <https://www.doi.org/10.1590/S0102-09352008000500034>
- Grès V, Voza T, Chabaud A and Landau I (2003). Coccidiosis of the wild rabbit (*Oryctolagus cuniculus*) in France. *Parasite*, 10: 51-57. DOI: <https://www.doi.org/10.1051/parasite/2003101p51>
- Hamid PH, Prastowo S and Kristianingrum Y (2019). Intestinal and hepatic coccidiosis among rabbits in Yogyakarta, Indonesia. *Veterinary World*, 12: 1256-1260. DOI: <https://www.ncbi.nlm.nih.gov/pubmed/31641305>
- Hana A, Salasia SIO, Mangkoewidjojo S and Kusindarto DL (2011). Blood profile of rabbits infected with *Eimeria magna*. *Animal Production*, 13: 185-190. Available at: <http://animalproduction.net/index.php/JAP/article/view/333>
- Hanada S, Umemoto Y, Omata Y, Koyama T, Nishiyama K, Kobayashi Y, Furuoka H, Matsui T, Maeda R and Saito A (2003). *Eimeria stiedae* merozoite 49-kDa soluble antigen induces protection against infection. *Journal of Parasitology*, 89: 613-617. DOI: [https://www.doi.org/10.1645/0022-3395\(2003\)089\[0613:esmksa\]2.0.co;2](https://www.doi.org/10.1645/0022-3395(2003)089[0613:esmksa]2.0.co;2)
- Hanptman K, Tichy F and Knotek Z (2001). Clinical diagnostics of hepatopathies in small mammals: Evaluation of importance of individual method. *Acta Veterinaria Brno*, 70: 297-311. DOI: <https://www.doi.org/10.2754/avb200170030297>
- Hassan KM, El-Askalany MA, Mousa WM, Shokier KAM, Arafah WM and Aboelhadid SM (2015). Molecular diagnosis of *Eimeria stiedae* in hepatic tissue of experimentally infected rabbits in comparison with traditional methods. *Egyptian Veterinary Medical Society of Parasitology Journal*, 11: 51-57. Available at: https://www.researchgate.net/deref/http%3A%2F%2Fdx.doi.org%2F10.21608%2Fevmspj.2015.37917?_sg%2F
- Indrasanti D, Indradji M, Hastuti S, Fatikha AE and Rosyadi KA (2017). The administration of garlic extract on *Eimeria stiedae* oocysts and the hematological profile of the coccidia infected rabbits. *Media Peternakan*, 40: 158-164. DOI: <https://www.doi.org/10.5398/medpet.2017.40.3.158>

- Indrasanti D, Indradji M, Hastuti S, Wihadmadyatami H and Ismoyowati A (2015). The efficacies of banana stem extract as a candidate of coccidiostat against rabbit *Eimeria stiedae* oocysts: An *in vitro* analysis. *Animal Production* 17: 161-168. DOI: <https://www.dx.doi.org/10.20884/1.anprod.2015.17.3.503>
- Jing F, Yin G, Liu X, Suo X and Qin Y (2012). Large-scale survey of the prevalence of *Eimeria* infections in domestic rabbits in China. *Parasitology Research*, 110: 1495-1500. DOI: <https://www.doi.org/10.1007/s00436-011-2653-4>
- Jithendran KP (1995). Clinical coccidiosis in Angora rabbits. *Veterinary Revista Kathmandu*, 10: 21-22.
- Jithendran KP and Bhat T (1996). Subclinical coccidiosis in angora rabbits, a field survey in Himachal Pradesh, India. *World Rabbit Science*, 4: 29-32. DOI: <https://www.doi.org/10.4995/wrs.1996.267>
- John NM, Zea ME, Kawano T, Omata Y, Saito A, Toyoda Y and Milkano T (1999). Identification of carbohydrates on *Eimeria stiedae* sporozoites and their role in the invasion of cultured cells *in vitro*. *Veterinary Parasitology*, 81: 99-105. DOI: [https://www.doi.org/10.1016/s0304-4017\(98\)00239-8](https://www.doi.org/10.1016/s0304-4017(98)00239-8)
- Joyner LP, Catchpole J and Berrett S (1983). *Eimeria stiedae* in rabbits: The demonstration of responses to Chemotherapy. *Research in Veterinary Science*, 34: 64-67. DOI: [https://www.doi.org/10.1016/S0034-5288\(18\)32285-9](https://www.doi.org/10.1016/S0034-5288(18)32285-9)
- Kandil OM, Mahdy OA, Mousa WM and Derbala AA (2000). Evaluation of two *E. stiedae* antigens for serodiagnosis of hepatic coccidiosis in experimentally infected rabbits. *Journal of Egyptian Veterinary Medical Association*, 60: 83-91. Available at: <https://scholar.cu.edu.eg/?q=drolfat/publications&sort=year&order=asc>
- Kolabskii NA, Dubovoi BL and Vergerenko LV (1973). Effectiveness of sulfachlorpyrazine in coccidiosis of rabbits. *Veterinariia*, 4: 71-73. Available at: <https://www.hindawi.com/journals/jvm/2018/5402469/>
- Kowalska D, Bielanski P, Nosel P and Kowal J (2012). Natural alternatives to coccidiostats in rabbit nutrition. *Annals of Animal Science*, 12: 561-574. DOI: <http://www.izoo.krakow.pl/en/periodicals>
- Kraus AL, Weisbroth SH, Flatt RE and Brewer N (1984). Biology and diseases of rabbits. In: laboratory animal medicine Fox, J.G. (ed.), Pp: 270-240. Academic Press, Orlando, USA.
- Kuliscic Z, Tambur Z, Malicevic Z, Aleksic-Bakrac N and Mistic Z (2006). White blood cell differential count in rabbits artificially infected with intestinal coccidia. *Journal of Protozoology Research*, 16: 42-50. DOI: https://www.doi.org/10.32268/jprotozoolres.16.3-4_42
- Laha R, Das M and Goswami A (2015). Coccidiosis in rabbits in a subtropical hilly region. *Indian Journal of Animal Research*, 49: 231-233. DOI: <https://arccjournals.com/journal/indian-journal-of-animal-research/B-2614>
- Laha R, Dey HA and Harbola PC (1999). Comparative efficacy of sulphadimidine and combination of amprolium, sulphaquinoxaline in the control of natural coccidial infection in rabbits. *Indian Veterinary Journal*, 76: 1013-1015. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6218750/>
- Lakshmanan B, Ravindran R, Vasudeven VN and Devada K (2011). Hepatic coccidiosis in rabbits in Kerala. *Journal of Indian Veterinary Association, Kerala*, 9: 56-57. DOI: <http://jivaonline.net/jivaonline/index.php>
- Lebas F, Coudert P, Rouvier R and De Rochambeau H (1986). The rabbit husbandry, health and production. *Animal Production and Health*, no. 21, FAO, Rome, Italy.
- Li M and Ooi H (2009). Fecal occult blood manifestation of intestinal *Eimeria* spp. infection in rabbit. *Veterinary Parasitology*, 161: 327-329. DOI: <https://www.doi.org/10.1016/j.vetpar.2009.01.009>
- Licois D (2004). Domestic rabbit enteropathies. Proc. 8th World Rabbit Congress, invited paper. Puebla, Mexico, pp. 385-403.
- Magray A, Khan AR and Rashid M (2010). Study on fatal hepatic coccidiosis in rabbits of Kashmir Valley. *Journal of Environment and Biological Sciences*, 24: 243-245. Available at: https://zvzjournals.ekb.eg/article_59960_2c1fcb70729e80575ebc5585fa4eaf1d.pdf
- Mäkitaipale J, Karvinen I, Virtala AMK and Näreaho A (2017). Prevalence of intestinal parasites and risk factor analysis for *Eimeria* infections in Finnish pet rabbits. *Veterinary Parasitology, Regional Studies and Reports*, 9: 34-40. DOI: <https://www.doi.org/10.1016/j.vprsr.2017.04.005>
- Manjunatha V, Rout M, Sujay CS, Jaisingh N, Salin N and Byregowda SM (2019). Clinico-pathologic observations of spontaneous hepatic coccidiosis in broiler rabbits maintained in Bannerghatta Biological Park in Karnataka state of India. *Indian Journal of Animal Research*, 53: 528-532. DOI: <https://www.doi.org/10.18805/ijar.B-3436>
- Matekaire T, Mupangwa JF and Kanyamura EF (2005). The efficacy of banana plant (*Musa paradisiaca*) as a coccidiostat in rabbits. *International Journal of Applied Research in Veterinary Medicine*, 3: 326-331. Available at: arvm.com/articles/Vol3Iss4/MATEKAIRE%20IJARVM%20V3N4W.pdf
- Mikhail EG, Sabet S, El-Boulaqi HA, Zaki IE and Gaber A (1981). Treatment of hepatic coccidiosis in rabbits by tinidazole. *Journal of the Egyptian Society of Parasitology*, 1: 389-397. PMID: 7299175. Available at: <https://europepmc.org/article/med/7299175>
- Miller RE and Fowler ME (2003). *Zoology and Wild Animal Science*. McGraw-Hill: London.
- Nistor E, Bampidis VA, Pacala N, Pentea M, Tozer J and Prundeanu A (2013). Nutrient content of rabbit meat as compared to chicken, beef and pork meat. *Journal of Animal Production Advances*, 3: 172-176. DOI: <https://www.doi.org/10.5455/JAPA.20130411110313>
- Nosal P, Kowalska D, Bielanski P, Korwal J and Kornas S (2014). Herbal formulations as feed additives in the course of rabbit subclinical coccidiosis. *Annals of Parasitology*, 60: 65-69. PMID: 24930248. Available at: <https://pubmed.ncbi.nlm.nih.gov/24930248/>
- Ogolla KO, Chebet J and Gathumbi PK (2017). Farmer practices that influence risk factors, prevalence and control strategies of rabbit coccidiosis in Central Kenya. *Livestock Research for Rural Development*, p. 29. DOI: <http://www.lrrd.org/lrrd29/7/koko29134.html>

- Ogolla KO, Gathumbi PK, Waruiru RM, Okumu PO, Kitale PM and Chebet J (2018). Efficacy of sulphachloropyrazine, amprolium hydrochloride, trimethoprim-sulphamethoxazole, and diclazuril against experimental and natural rabbit coccidiosis. *Journal of Veterinary Medicine*; 5402469. DOI: <https://www.dx.doi.org/10.1155%2F2018%2F5402469>
- Okumu PO, Gathumbi PK, Karanja DN, Mande JD, Wanyoike MM, Gachui CK, Kiarie N, Mwanza RN and Borter DK (2014). Prevalence, pathology and risk factors for coccidiosis in domestic rabbits (*Oryctolagus cuniculus*) in selected regions in Kenya. *Veterinary Quarterly*, 34: 205-210. DOI: <https://www.doi.org/10.1080/01652176.2014.978044>
- Oliveira UC, Fraga JS, Licois D, Pakandl M and Gruber A (2011). Development of molecular assays for the identification of the 11 *Eimeria* species of the domestic rabbit (*Oryctolagus cuniculus*). *Veterinary Parasitology*, 176: 275-280. DOI: <https://www.doi.org/10.1080/01652176.2014.978044>
- Oncel T, Gulegen E, Senlik B and Bakirci S (2011). Intestinal coccidiosis in Angora rabbits (*Oryctolagus cuniculus*) caused by *Eimeria intestinalis*, *Eimeria perforans* and *Eimeria coecicola*. *YYU Veteriner Fakultesi Dergisi*, 22: 27-29. Available at: [http://vfdergi.yyu.edu.tr/archive/2011/22-1/2011_22_\(1\)_27-29.pdf](http://vfdergi.yyu.edu.tr/archive/2011/22-1/2011_22_(1)_27-29.pdf)
- Pakandl M (2009). Coccidia of rabbit: A review. *Folia Parasitologica*, 56: 153-166. DOI: <https://www.doi.org/10.14411/fp.2009.019>
- Pakandl M and Hlášková L (2007). The reproduction of *Eimeria flavescens* and *Eimeria intestinalis* in suckling rabbits. *Parasitology Research*, 101: 1435-1437. DOI: <https://www.doi.org/10.1007/s00436-007-0646-0>
- Pakandl M, HlaskovaL, Poplstein M, Chroma M, Vodicka T, Salat J and Mucksova J (2008). Dependence of the immune response to coccidiosis on the age of rabbit suckling. *Parasitology Research*, 103: 1265-1271. DOI: <https://www.doi.org/10.1007/s00436-008-1123-0>
- Papeschi C, Fichi G and Perrucci S (2013). Oocyst excretion pattern of three intestinal *Eimeria* species in female rabbits. *World Rabbit Science*, 21: 77-83. DOI: <https://www.doi.org/10.4995/wrs.2013.1235>
- Peeters JE and Geeroms R (1986). Efficacy of toltrazuril against intestinal and hepatic coccidiosis in rabbits. *Veterinary Parasitology*, 22: 21-35. DOI: [https://www.doi.org/10.1016/0304-4017\(86\)90004-X](https://www.doi.org/10.1016/0304-4017(86)90004-X)
- Peeters JE and Geeroms R (1989). Efficacy of diclazuril against robenidine resistant *Eimeria magna* in rabbits. *The Veterinary Record*, 124: 589-590. DOI: <https://www.doi.org/10.1136/vr.124.22.589>
- Peeters JE, Geeroms R, Antoine O, Mammerickx M and Halen P (1981). Efficacy of narasin against hepatic and intestinal coccidiosis in rabbits. *Parasitology*, 83, 293-301. DOI: <https://www.doi.org/10.1017/s0031182000085309>
- Peeters JE, Geeroms R, Molderez J and Halen P (1982). Activity of clodidol/methylbenzoquate, robenidine and salinomycin against hepatic coccidiosis in rabbits. *Reihe B. Journal of Veterinary medicine. Series B*, 29: 207-218. DOI: <https://www.doi.org/10.1111/j.1439-0450.1982.tb01217.x>
- Pilarczyk B, Tomza-Marciniak A, Pilarczyk R, Januś E, Stanek P, Seremak B and Sablik P (2020). The effect of the sex, age, and breed of farmed rabbits and the choice of management system on the extensivity and intensity of *Eimeria* infection. *Veterinary World*, 13: 1654-1660. DOI: <https://www.dx.doi.org/10.14202%2Fvetworld.2020.1654-1660>
- Polozowski A (1993). Coccidiosis of rabbits and its control. *Wiadomosci parazytologiczne*, 39: 13-28. PMID: 8346656. Available at: <https://pubmed.ncbi.nlm.nih.gov/8346656/>
- Qamar F, Sharif R, Qamar MM and Basharat A (2013). Comparative efficacy of sulphadimidine sodium, toltrazuril and amprolium for coccidiosis in rabbits. *Science International (Lahore)*, 25: 295-298. Available at: https://www.researchgate.net/publication/316692292_Comparative_efficacy_of_sulphadimidine_sodium_toltrazuril_and_amprolium_for_Coccidiosis_in_Rabbits
- Redrobe SP, Gakos G, Elliot SC, Saunders R, Martin S and Morgan ER (2010). Comparison of toltrazuril and sulphadimethoxine in the treatment of intestinal coccidiosis in pet rabbits. *The Veterinary Record*, 167: 287- 290. DOI: <https://www.doi.org/10.1136/vr.c3453>
- Regulation (EC) No. 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition/ OJ L 268: 29-43. Available at: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32003R1831&rid=10>
- Renaux S, Quéré P, Buzoni-Gatel D, Sewald B, Le Vern Y, Coudert P and Drouetviard F (2003). Dynamics and responsiveness of T lymphocytes in secondary lymphoid organs of rabbits developing immunity to *Eimeria intestinalis*. *Veterinary Parasitology*, 110: 181-195. DOI: [https://www.doi.org/10.1016/S0304-4017\(02\)00305-9](https://www.doi.org/10.1016/S0304-4017(02)00305-9)
- Rivero-Perez N, Hernández-Alvarado JL, Valladares-Carranza B, Delgado-Ruiz L, Ojeda-Ramírez D, Sosa-Gutiérrez CG, Morales-Ubaldo AL, Vega-Sánchez V and Zaragoza-Bastida A (2019). *Salix babylonica* L. as a natural anticoccidial alternative in growing rabbits. *Evidence Based Complementary Alternative Medicine*, pp. 1-8. DOI: <https://www.doi.org/10.1155/2019/2107231>
- Sanyal PK and Sharma SC (1990). Clinicopathology of hepatic coccidiosis in rabbits. *Indian Journal of Animal Science*, 60: 924-928. Available at: <https://arccjournals.com/journal/indian-journal-of-animal-research/B-3436>
- Schlögl W, Hudson R and Rödel HG (2013). Impact of rearing management on health in domestic rabbits: A review. *World Rabbit Science*, 21: 145-159. DOI: <https://www.doi.org/10.4995/wrs.2013.1029>
- Seddek ASH, El-Ghoneimy AA, Shibat El-hamd DW and Mahmoud EG (2015). Effect of *Calotropis procera* on rabbits coccidiosis. *Egyptian Journal of Chemical and Environmental Health*, 1: 768-784. Available at: <http://cehea.org/wp-content/uploads/2016/08/59-.pdf>
- Sharma KG, Vidyarthi VK, Archana K and Zuyie R (2016). Probiotic supplementation in the diet of rabbits a review. *Livestock Research International*, 4: 1-10. Available at: http://jakraya.com/journal/pdf/11-lriArticle_1.pdf
- Shkromada O, Skliar O, Paliy A, Ulko L, Suprun Y, Naumenko O, Ishchenko K, Kysterina O, Musiienko O and Paliy A (2019). Development of preventing means for rabbits' coccidiosis. *Eureka: Health Sciences*, 3: 58-68. DOI: <https://www.dx.doi.org/10.21303/2504-5679.2019.00914>

- Singla LD, Juyal PD and Sandhu BS (2000). Pathology and therapy in naturally *Eimeria stiedae*-infected rabbits. *Journal of Protozoology Research*, 10: 185-191. Available at: <https://www.semanticscholar.org/paper/Pathology->
- Sivajothi S, Reddy BS and Rayulu VC (2016). Study on impression smears of hepatic coccidiosis in rabbits. *Journal of Parasitic Diseases*, 40: 906-909. DOI: <https://www.dx.doi.org/10.1007%2Fs12639-014-0602-8>
- Song H, Yan R, Xu L, Song X, Shah MAA, Zhu H and Li X (2010). Efficacy of DNA vaccines carrying *Eimeria acervulina* lactate dehydrogenase antigen gene against coccidiosis. *Experimental Parasitology*, 126: 224-231. DOI: <https://www.doi.org/10.1016/j.exppara.2010.05.015>
- Song H, Dong R, Qiu B, Jing J, Zhu S, Liu C, Jiang Y, Wu L, Wang S, Miao J and Shao Y (2017). Potential vaccine targets against rabbit coccidiosis by immunoproteomic analysis. *The Korean Journal of Parasitology*, 55: 15-20. DOI: <https://www.dx.doi.org/10.3347%2Fkjp.2017.55.1.15>
- Sorour SS, Abou-Asa S, Elhawary NM, Ghazy EW, Abd-El-Latif A, El-Abasy MA and Khalifa HO (2018). Anticoccidial and hepatoprotective effects of artemisinin liquid extract, cinnamon essential oil and clove essential oil against *Eimeria stiedae* infection in rabbits. *Tropical Biomedicine*, 35: 926-943. DOI: <http://msptm.org/files/Vol35No4/926-943-Hazim-O-Khalifa.pdf>
- Soulsby EJL (1968). *Helminths, Arthropods and Protozoa of Domesticated Animals*. 6th ed. Bailliere Tindall, London, 676-682.
- Toulah FH and Al-Raw MM (2007). Efficacy of garlic extract on hepatic coccidiosis in infected rabbits (*Oryctolagus cuniculus*): histological and biochemical studies. *Journal of Egyptian Society of Parasitology*, 37: 957-968. Available at: <https://pubmed.ncbi.nlm.nih.gov/18383795/>
- Vanparijs O, Desplenter L and Marsboom R (1989a). Efficacy of diclazuril in the control of intestinal coccidiosis in rabbits. *Veterinary Parasitology*, 34: 185-190. DOI: [https://www.doi.org/10.1016/0304-4017\(89\)90049-6](https://www.doi.org/10.1016/0304-4017(89)90049-6)
- Vanparijs O, Hermans L, Van Der Flaes L and Marsboom R (1989b). Efficacy of diclazuril in the prevention and cure of intestinal and hepatic coccidiosis in rabbits. *Veterinary Parasitology*, 32: 109-117. DOI: [https://www.doi.org/10.1016/0304-4017\(89\)90111-8](https://www.doi.org/10.1016/0304-4017(89)90111-8)
- Vereecken D, Lavazza A, De Gussem K, Chiari M, Tittarelli C, Zuffellato A and Maertens L (2012). Activity of diclazuril against coccidiosis in growing rabbits: experimental and field experiences. *World Rabbit Science*, 20: 223-230. DOI: <https://www.doi.org/10.4995/wrs.2012.1232>
- Wei W, Shen N, Xiao J, Tao Y, Luo Y, Angel C, Gu X, Xie Y, He R, Jing B, Peng X and Yang G (2020). Expression analysis and serodiagnostic potential of microneme proteins 1 and 3 in *Eimeria stiedae*. *Genes*, 11: 725. DOI: <https://www.doi.org/10.3390/genes11070725>
- Xin C, Wu B, Li J, Gong P, Yang J, Li H, Cai X and Zhang X (2016). Complete genome sequence and evolution analysis of *Eimeria stiedae* RNA virus 1, a novel member of the family Totiviridae. *Archive of Virology*, 161: 3571-3576. DOI: <https://www.doi.org/10.1007/s00705-016-3020-7>
- Yan W, Wang W, Wang T, Suo X, Qian W, Wang S and Fan D (2013). Simultaneous identification of three highly pathogenic *Eimeria* species in rabbits using a multiplex PCR diagnostic assay based on ITS1- 5.8S rRNA-ITS2 fragments. *Veterinary Parasitology*, 193: 284-288. DOI: <https://www.doi.org/10.1016/j.vetpar.2012.11.013>
- Yin G, Goraya MU, Huang J, Suo X, Huang Z and Liu X (2016). Survey of coccidial infection of rabbits in Sichuan Province, Southwest China. *Springer Plus*, 5: 870. DOI: <https://www.doi.org/10.1186/s40064-016-2586-6>
- Zayed AA and Kutkat MA (1998). SDS-Page and immunoelectrophoretic evaluation of sporulated oocysts antigens of *E. stiedae* infecting domestic rabbits. *Alexandria Journal of Science*, 14: 1-9. Available at: [https://www.idosi.org/gv/GV9\(5\)12/3.pdf](https://www.idosi.org/gv/GV9(5)12/3.pdf)