



Newcastle Disease Virus in Poultry: Current Status and Control Prospects

Ahamidou Moustapha^{1*}, Essodina Talaki^{1,2}, Adamou Akourki³, and Moumouni Ousseini⁴

¹University of Lomé (UL), Regional Center of Excellence on Avian Sciences (CERSA), 01 BP: 1515, Lomé, Togo

²University of Lomé (UL), Higher School of Agronomy (ESA), 01 BP: 1515, Lomé, Togo

³Dan Dicko Dankoulodo University of Maradi (UDDM), Faculty of Agronomy and Environmental Sciences (FASE), BP: 465, Maradi, Niger

⁴Bayero University Kano, Faculty of Agriculture, Department of Animal Science, PMB 3011, Kano, Nigeria

*Corresponding author's Email: ahmidoum1@gmail.com

ABSTRACT

Since its first appearance in Java Island, Indonesia, in 1926, Newcastle disease has significantly impacted the global poultry industry, leading to substantial economic losses. The disease has rapidly spread worldwide, becoming endemic in many countries where agriculture is the primary source of national income. The present study aimed to present a comprehensive review of the recent literature on the Newcastle disease virus to contribute to understanding the virus and its control measures in poultry and provide an updated perspective on current knowledge. All strains of the Newcastle disease virus are classified under a single serotype; however, they are grouped into two classes and have been found to have emerging genetic diversity. Although various molecular diagnostic protocols have been developed, many have limitations. Nucleotide variability in the *F* gene of the Newcastle disease virus seems to explain the false-negative results provided by different real-time reverse transcription polymerase chain reaction protocols. Vaccination combined with biosecurity measures has been shown to limit the devastating effect of the Newcastle disease virus. However, the current vaccines are not effective enough to prevent viral shedding and infection of vaccinated animals. The efficacy of the vaccine strains utilized for decades is being scrutinized, raising questions about their effectiveness over time. The development of reverse genetics offers promising prospects for exploring new generations of attenuated vaccines capable of protecting poultry against clinical diseases and infections, such as Newcastle disease.

Keywords: Diagnosis, Genotype, Newcastle disease, Pathogenicity, Poultry, Vaccination

INTRODUCTION

Newcastle disease, or fowl plague, is a highly contagious viral poultry disease affecting various avian species with varying degrees of susceptibility (Shakal et al., 2020). The virus is easily transmitted, leading to its rapid spread worldwide within three decades after its initial appearance in Indonesia in 1926. Newcastle disease is currently endemic in many countries worldwide (Azizah et al., 2021). Based on pathogenic studies, the virus strains can be classified into five pathotypes, namely asymptomatic enteric, lentogenic, mesogenic, viscerotropic, and neurotropic velogenic strains (Abdisa and Tagesu, 2017; Getabalew et al., 2019). Velogenic strains are capable of causing 100% mortality in unprotected poultry farms, leading to trade restrictions and embargoes in countries where the disease occurs (Dzogbema et al., 2021; Ali et al., 2022). The disease can harmfully affect developing countries where agriculture is the main source of household income and food (Fellahi and Boudouma, 2021). Newcastle disease is a notifiable disease by the World Organization for Animal Health (WOAH). It is defined as any infection caused by an isolate of *Avian Paramyxovirus type 1* with an intracerebral pathogenicity index (ICPI) in day-old chicks of 0.7 or greater and having several basic amino acids at the C-terminal part of the F2 protein and phenylalanine at residue 117 of the N-terminal part of the F1 protein (WOAH, 2022).

Despite the use of vaccines to limit the devastating effects of the virus, Newcastle disease continues to affect the poultry industry worldwide, causing epidemics and significant economic losses (Hu et al., 2022). Although current vaccines can prevent the outbreak of the disease and death from the disease when administered properly, the virus can infect vaccinated animals and replicate, leading to its spread via feces and saliva to other birds (Miller and Koch, 2013; Bello et al., 2018; Mahamud et al., 2022). Recent years have seen a renewed interest in research on the virus due to its pathogenic potential and its use as a vaccine vector in both human and animal health (Jamil et al., 2022; Naz et al., 2022). Advances in molecular biology and genomic sequencing have greatly expanded the understanding of the biology and replication of the Newcastle disease virus (NDV). Furthermore, the development of reverse genetics offers prospects for the construction of a new generation of an attenuated vaccine capable of controlling both the clinical disease and its infection with regard to the modular nature of transcription, low frequency of recombination, and the absence of DNA phase during replication (Fellahi and Boudouma, 2021).

REVIEW ARTICLE
pitt: S232245682300026-13
Received: 23 February 2023
Accepted: 13 April 2023

The objective of this review was to provide a synthesis of recently published studies in the field. This review included the original studies conducted on the NDV, its virulence and pathogenicity mechanisms, genetic diversity, clinical and lesional picture resulting from its infection, and new methods and tools for laboratory diagnosis. The investigated articles were limited to those searched in the Google Scholar search engine. This review also examined the prevention and control methods currently in use and prospect.

Taxonomy and structure of Newcastle disease virus

Newcastle disease is caused by virulent strains of *Avian orthoavulavirus 1* (AOAV-1), formerly known as *Avian avulavirus 1* (AAvV-1) and commonly known as *Avian paramyxovirus 1* (APMV-1) or NDV. Newcastle disease virus belongs to the genus *Orthoavulavirus*, subfamily *Avulavirinae*, family *Paramyxoviridae* and order *Mononegavirales*. Isolated avian paramyxoviruses are classified into 21 serotypes designated APMV-1 to APMV-21. Each virus belongs to a viral species that are dispersed among the genera *metaavulavirus*, *orthoavulavirus*, and *paraavulavirus* (Amarasinghe et al., 2019; Dimitrov et al., 2019).

Newcastle disease virus is a pleomorphic enveloped virus sized approximately 200-300 nm in diameter. Its envelope is derived from the plasma membrane of the infected cell, with an outer face, where 8-12 nm long spicules are inserted, corresponding to the HN glycoprotein and the F protein. The inner side of the envelope is lined with a matrix protein. The viral genome is a non-segmented, single-stranded RNA of negative polarity. The RNA genome of NDV has a molecular weight of 5.2 to 5.7 X 10⁶ Daltons. Genome sizes vary between 15186 and 15198 nucleotides (Ali et al., 2022). The nucleoprotein combines with the RNA to form a tubular nucleocapsid with a diameter of 18 nm, and the nucleoprotein, phosphoprotein, and large protein are tightly bound to the genomic RNA (Mao et al., 2022).

Viral proteins

The NDV genome consists of six genes, encoding six different proteins. The genes arranged in order of 3'-NP-P-M-F-HN-L-5' encode for nucleoprotein, phosphoprotein, matrix protein, fusion protein, hemagglutinin-neuraminidase protein, and large polymerase protein, respectively (Phale, 2018; Nurziah et al., 2022). The nucleoprotein has a length of 489 amino acids and a molecular weight of 55 kDaltons. It surrounds the genomic RNA and protects it from nucleases, while phosphoprotein is made up of 395 amino acids and forms multiple bands with a molecular weight ranging from 50 to 55 kDaltons, which is essential in viral replication and transcription. Matrix protein (M) is a non-glycosylated protein associated with the inner surface of the envelope. It consists of 364 amino acids and has a molecular weight of about 40 kDaltons. It plays an important role in viral assembly and envelope stabilization. Glycoprotein (F) consists of 553 amino acids and fuses the viral envelope with the membrane of the target cell. Glycoprotein (HN) is a 577 amino acid long polypeptide with a molecular weight of 74 kDaltons. It is a protein with many functions related to the attachment of the virion to the target cell receptor, enzymatic cleavage of sialic acid, and promotion of viral fusion. The large protein is the largest in the viral genome consisting of 2204 amino acids and has a molecular weight of 250 kDaltons. This protein synthesizes viral mRNA and assists in the replication of genomic RNA (Ganar et al., 2014; Phale, 2018; Fellahi and Boudouma, 2021). Accessory proteins V and W are proteins derived from the P gene translated from alternative mRNAs produced by RNA editing during P gene transcription. The V protein consists of 239 amino acids with a molecular weight of 36 kDaltons. It is an interferon antagonist and plays an important role in NDV virulence. Finally, the role of the W protein in the NDV replication cycle has not yet been established (Ganar et al., 2014; Nurziah et al., 2022).

Viral replication

The spread of NDV in the infected organism occurs in several steps. It begins with the penetration of the virus into the host cell via surface glycoproteins (HN and F). The HN protein binds to sialic acid-containing cell receptors on the host cell surface, while the F protein mediates viral entry into the host cell. After the fusion of the viral envelope with the host cell membrane, the viral nucleocapsid is released into the cytoplasm, where the transcription and translation phase begin. During this phase, the negative-sense RNA genome is transcribed into positive-sense mRNA, which is then translated into viral proteins. The mRNA of individual genes is synthesized and used as a template for the synthesis of negative-sense genomic RNA. Transcription of genomic RNA into messenger RNA is carried out by the viral transcriptase (P and L) associated with the nucleocapsid. After the transcription and translation phase, the newly formed viral particles undergo budding and release. The M protein of NDV is essential for the assembly and budding of viral particles (Mao et al., 2022). Its nuclear localization could promote viral replication, increase viral RNA synthesis, and inhibit host cell replication (Duan et al., 2019). The F protein synthesized as a non-functional precursor F₀ is cleaved into F₁ and F₂ by host cell proteases. Viral proteins synthesized in the host cell are transported and incorporated at the cell membrane. Once the nucleocapsids are incorporated into the modified regions of the cell membrane, new viral particles bud at the cell surface, taking with them part of the cytoplasmic membrane. Finally, the HN protein catalyzes the cleavage of sialic acid in the cell receptor and allows the release of the virus (Ganar et al., 2014; Mao et al., 2022).

Infection can also occur by receptor-mediated endocytosis (Sánchez-Felipe et al., 2014). Cellular cholesterol may be required for optimal cell entry into the NDV infection cycle (Martín et al., 2012).

Resistance, virulence, and pathogenicity

The NDV can survive for long periods at room temperature (18 to 23.5°C), especially in feces (Getabalew et al., 2019; Dzogbema et al., 2021). However, it is inactivated at an elevated temperature of 56°C for 3 hours or 60°C for half an hour. It is also inactivated by an acidic pH ≤ 2 . Finally, it is sensitive to detergents, lipid solvents, formaldehyde, and oxidizing agents (Getabalew et al., 2019).

The pathogenicity of NDV differs depending on the virus strain. It is well known that activation of the fusion protein (F) by host cell proteases is the primary determinant of virulence (Ganar et al., 2014). An inactive precursor glycoprotein (F0) is produced during replication. It must be cleaved into F1 and F2 polypeptides to exhibit virulence and become infectious (Worku and Teshome, 2020). The cleavability of the F0 molecule is directly related to the virulence of the virus. However, it can be influenced by the nature of proteases present in host tissues and organs. Molecular studies of the F0 glycoprotein site have revealed that the sequence of the F cleavage site between positions 112 and 116 primarily determines the virulence of NDV isolates (Puro and Sen, 2022). Virulent viruses are cleaved by furin-like proteases found in many tissues and organs, allowing them to cause lethal systemic infections. In contrast, trypsin-like proteases can only cleave low-virulence viruses in limited areas, such as the respiratory and intestinal tracts (Phale, 2018). According to Heiden et al. (2014), regions of the F protein modulate virulence other than the polybasic cleavage site.

The development of reverse genetics over the past two decades has allowed researchers to modify NDV genetically and study the individual contribution of genes and genome regions concerning its virulence (Dortmans et al., 2011). According to Samal et al. (2011), the conserved glutamine residue at position 114 in the cleavage site of the F protein is important in NDV replication and pathogenicity. In addition, the replacement of isoleucine at position 118 with valine around the fusion cleavage site reduces the pathogenicity of NDV. Additionally, mutation of the F protein glycosylation site can lead to a hyper fusogenic virus that could increase the virulence of NDV strains (Samal et al., 2012). Mutation of the cytoplasmic domain of the F protein can lead to the creation of a new hyperfusogenic virus that may increase viral replication and pathogenicity (Samal et al., 2013). Other studies have found that the amino acid residue located on site 430 of the HN protein of NDV can influence viral fusion capacity by promoting F protein cleavage (Chen et al., 2021). In addition, ubiquitination on lysine 247 of the NDV matrix protein enhances viral replication and virulence by promoting nuclear-cytoplasmic trafficking (Peng et al., 2022).

Genetic diversity and geographic distribution

Several classification schemes have been developed to identify and differentiate strains of NDV. Initially, these schemes were based on the biological properties of the virus (Diel et al., 2012). However, a new classification system has been proposed by Dimitrov et al. (2019), which is based on phylogenetic topology, genetic distances, branch support, and epidemiological independence of the virus. This system maintains two classes (I and II) of NDV, identifies three new class II genotypes, and reduces the number of subgenotypes. The class I NDV isolates are all grouped into a single genotype and three subgenotypes, as they exhibit strong genetic relatedness. These isolates have been identified in wild and domestic birds from Africa, Asia, Europe, and America (Rauw et al., 2009). It is worth noting that NDV isolates belonging to class I are generally considered to be of low virulence to chicken, with a few exceptions, such as an isolate that caused a severe epidemic in Northern Ireland in the early 1990s (Bello et al., 2018), as well as the JS10-A10 and 9a5b strains generated by consecutive experimental passages through chicken (Rehman et al., 2018).

Class II NDV isolates include viruses of varying susceptibilities, including lentogenic, mesogenic, and velogenic strains, resulting in nearly twenty genotypes due to ongoing virus emergence and evolution (Bello et al., 2018). The most prevalent genotype among all class II viruses in waterfowl is genotype I, which is apathogenic like nearly all class I viruses. Genotype I is further divided into three subgenotypes, namely 1a, 1b, and 1c (Rauw et al., 2009). Genotype II is neurotropic but includes viruses of different pathogenicity (lentogenic, mesogenic, and velogenic). Some viruses belonging to this genotype, including B1, LaSota, VG/GA, and strains not commonly isolated, are used as vaccines (Miller et al., 2010). Genotype III has been isolated mainly in Southeast Asia, Australia, Japan, the United Kingdom, Zimbabwe, Singapore, and China. All NDV isolates belonging to genotype III are virulent, and their isolates have been found in chicken and domestic waterfowl (Dimitrov et al., 2016). Genotype IV is associated with Newcastle disease epizootics in Europe after World War II (Wehmann et al., 2003). Genotype V includes four subgenotypes (Va-Vd) and was responsible for an epizootic wave that started in Western Europe and spread to Yugoslavia during the 1970s (Wehmann et al., 2003). This genotype has also been isolated in Central and North America and Africa (Denis et al., 2015). Genotype VI has 11 subgenotypes (VIa-VIi) and has been described in pigeons in Asia, Africa, Europe, and South America (Dimitrov et al., 2016; Bello et al., 2018). Genotype VII is commonly found in Asia and Africa (Xue et al., 2017; Naguib et al., 2021) and has been associated with the most recent Newcastle disease epizootics

worldwide (Miller et al., 2015). This genotype has complex genetic diversity since it can be subdivided into 9 subgenotypes (VIIa-VIIi; Bello et al., 2018). Genotype VIII has been isolated from South Africa, South Asia, and western China (Cao et al., 2013; Denis et al., 2015; Megahed et al., 2020). Virulent and low-virulent isolates of genotype IX have been described in several asymptomatic wild bird species in China (Duan et al., 2014). Virulent strains of genotype X have been identified in Taiwan, Argentina, and the USA. Genotype XI is also virulent, but its distribution is largely restricted to the island of Madagascar (Maminaiina et al., 2010). Genotype XII has been isolated in South America and China (Liu et al., 2013; Chumbe et al., 2017), while XIII genotype includes three subgenotypes (XIIIa-XIIIc) and has been identified in Asia, Europe, and Africa (Denis et al., 2015; Das and Kumar, 2016; Ana et al., 2020). Genotype XIV includes two subgenotypes and is highly virulent, and has only been found in domestic birds in Africa (Snoeck et al., 2013; Samuel et al., 2013). Genotype XV was isolated in China and included virulent and vaccine strains (Bello et al., 2018). Genotype XVI is strongly related to genotype IV and has been isolated in Europe, Africa, and Asia (Bello et al., 2018). Finally, genotypes XVII and XVIII have two subgenotypes, and they are highly virulent and are found in West and Central Africa (Snoeck et al., 2013; Bello et al., 2018; Souley et al., 2021).

Host species

Newcastle disease virus is capable of infecting over 200 different bird species, but the disease infection consequence varies greatly depending on the host species and the virus strain (Rauw et al., 2009). Newcastle disease affects chickens more severely than turkeys, which typically show few clinical signs (WOAH, 2022). Pheasants, partridges, quail, and guinea fowl are also susceptible (Getabalew et al., 2019). Pigeons (Colombiformes) can also be infected with pigeon paramyxovirus (PPMV-1, Ramsubeik et al., 2023). Ducks and geese (Anseriformes) may be infected but show little to no clinical signs (Rahman et al., 2018). Teal, swans, wild geese, double-crested cormorants, white pelicans, and gulls are susceptible, but they are generally resistant and show little clinical signs of apathy and anorexia (Rasamoelina et al., 2016). Parrots, ravens, sparrows, and kingbirds are susceptible (Rahman et al., 2018). Generally, wild birds and waterfowl are considered reservoir hosts (Getabalew et al., 2019). Newcastle disease can also infect a range of non-avian species. Evidence of Newcastle disease infection and detection have been reported in cattle, sheep, mink, hamsters, mice, rabbits, camels, pigs, monkeys, and humans (UI-Rahman and Shabbir, 2019; Shabbir et al., 2021; UI-Rahman et al., 2022). Workers in poultry production and processing and vaccine production laboratories are at the greatest risk (Shabbir et al., 2021).

Mode of dissemination

Transmission of NDV can occur through inhalation or ingestion of virus particles. In the case of natural infection, the virus multiplies in the respiratory and/or digestive tract of infected poultry, which then excretes the virus through airborne or fecal routes (Brown and Bevins, 2017). These released virus particles are inhaled by healthy birds or affect their mucous membranes. Viruses released in feces may also contaminate feed and drinking water and thus be ingested by other birds in the poultry house (Rauw et al., 2009). Non-bird species, such as cats, dogs, foxes, and rodents, may play a role in virus transmission because they may spread the virus for up to 72 hours in their feces after ingesting contaminated poultry (Rasamoelina et al., 2016). Similar to insects, rodents, and reptiles may also be potential vectors of NDV, as their susceptibility to infection has been reported (Rasamoelina et al., 2016). Semi-captive or free-living wild birds may act as natural reservoir hosts for NDV and play a considerable role in the spread of the virus (Rahman et al., 2018).

Clinical signs and lesions

Pathogenicity of the virus strain, host species, host age, secondary infections, stress, environmental conditions, host immune status, viral dose, and route of exposure may play a role in determining incubation time and disease severity (Getabalew et al., 2019; WOAH, 2022). Infection with lentogenic NDV virus strains can range from unapparent respiratory or gastrointestinal illness to mild disease in adult chickens. Infection with mesogenic isolates can lead to the development of nonfatal respiratory disease, decreased egg production, and rare nervous signs, but mortality is generally low (Abdisa and Tagesu, 2017). Infection with highly virulent velogenic viruses can cause mortality in highly susceptible flocks of up to 100% (WOAH, 2022). Viscerotropic velogenic strains cause acute infection of the gastrointestinal mucosa and result in hemorrhagic lesions and death. Clinical signs resulting from chicken infection include general ill health, respiratory disturbances, greenish or whitish diarrhea, muscle tremors, and paralysis of the extremities. Edema may also be seen around the eyes (Caroline, 2022). Neurotropic velogenic strains of NDV primarily cause respiratory distress followed by neurological disease and decreased egg production (Abdisa and Tagesu, 2017). Mortality in chickens infected with velogenic NDVs can reach 100% (Alazawy and Al Ajeeli, 2020).

Newcastle disease virus lesions are primarily characterized by hemorrhages and ulcerations in the proventriculus and intestine (Rauw et al., 2009). Infection with viscerotropic velogenic NDV can result in severe clinical signs in chickens, including dehydration and emaciation of the chicken carcass, hemorrhagic lesions, and necrotic ulcerations at

the tips of the proventricular glands, in the intestine, caecal tonsils, and spleen parenchyma. Additionally, the lungs may show congestive, edematous, and hemorrhagic lesions (Mariappan et al., 2018; Sonkusale et al., 2023). In the respiratory form of the disease in chicken, lesions are marked initially by congestive or hemorrhagic tracheitis that becomes complicated after a few days of consecutive colibacillosis evolution into fibrinous tracheitis, fibrinous pneumonia, aerosacculitis, pericarditis and perihepatitis (Fellahi and Boudouma. 2021). The involvement of the central nervous system in NDV infection can be observed through anatomical manifestations, such as progressive encephalitis, neuronal necrosis, neuronal phagocytosis, the presence of clusters of cells of microglial morphology in the gray matter of the brain, cerebellum and spinal cord, axonal degeneration, and demyelination lesions (Ecco et al., 2011).

Diagnose

Observed clinical signs cannot provide a reliable basis for diagnosis, but can be used as an element of suspicion to guide the diagnosis (Abdisa and Tagesu, 2017). Lesions may also be confused with those of other diseases, including Avian Influenza (Ganar et al., 2014). However, a definitive diagnosis of Newcastle disease is based on the direct detection of viral antigens, which can be achieved by virus isolation from swabs of live animals or organs taken from cadavers. Virus culture is readily performed on 9-11-day-old embryonated chicken eggs or a wide range of cells, including chicken embryo fibroblast cells, chicken embryo hepatocytes, African vervet monkey kidney cells, and chicken embryo reticulum cells (WOAH, 2022). After incubation at 37°C for 4-7 days, the inoculated embryonated chicken eggs are refrigerated at +4°C. The allantoic-amniotic fluid from the eggs is then tested for NDV using the hemagglutination test. Confirmation of NDV infection can be done by performing a hemagglutination inhibition test or by using molecular methods (WOAH, 2022). Agar gel immunodiffusion technique, fluorescent antibody test, hemolysis test, or identification of viral particle morphology by electron microscopy can also be used to identify NDV, but none of them provide information on the pathotype of ND (Cattoli et al., 2011).

Serological diagnosis provides indirect evidence of NDV through antibodies that demonstrate infection. The hemagglutination inhibition (HI) test is the most widely used reference and confirmatory test for Newcastle disease serology (WOAH, 2022). The test is based on the agglutination of chicken red blood cells caused by the hemagglutinin of the viral envelope and the inhibition of hemagglutination by specific antibodies. Enzyme-linked immunosorbent assays (ELISAs) are also commonly used for the detection and quantification of antibodies. The antibody detection technique consists of binding antibodies to viral antigens on a microtiter plate. The viral antigens are captured by other detection antibodies produced in another species against those of the chicken and coupled to an enzyme that catalyzes the reaction to cause a color change. The plate is then read on a spectrophotometer (Dzogbema et al., 2021). The HI test correlates well with the ELISA test (Getabalew et al., 2019; WOA, 2022). An intracellular cytokine labeling technique was developed using Pinette et al. (2014) to quantify the immune response mediated by T cells. There are other serological tests, such as the neutralizing antibody test, the immunofluorescence test, and the colloidal gold immune technique. However, the limitation of serological tests lies in the cross-reactivity between NDV and other homologous avian paramyxoviruses, which can give false positive results (Mao et al., 2022).

Techniques based on molecular biology methods have been developed for the isolation and identification of NDV strains (Dzogbema et al., 2021; Moa et al., 2022). Several laboratory protocols have been developed, including gel-based reverse transcription polymerase chain reaction (RT-PCR), real-time RT-PCR, restriction enzyme-based methods, and rapid sequencing. In addition, nested PCR, fluorogenic probe-based real-time RT-PCR, ligase chain reaction (LCR), intercalated DNA (SYBR Green), or light-extended fluorogenic primer (LUX) should also be mentioned. All of these assays have shown promising results, but most of them have some limitations and their ability to detect different virus types needs validation (Cattoli et al., 2011; Mao et al., 2022). Since the NDV *F* gene is the primary determinant of NDV pathogenicity, RT-PCR protocols to pathotype NDV generally target this gene (Putri et al., 2017; Abd Elfatah et al., 2021; Mao et al., 2022). The growth of the gene sequence database over the last decade has revealed the high degree of nucleotide variability in the NDV *F* gene, which is mainly responsible for mismatches between oligonucleotides (primers and/or probes) and cDNA templates. This may well explain the false negative results provided by different real-time RT-PCR protocols (Cattoli et al., 2011). Another real-time reverse transcription isothermal loop amplification (RT-LAMP) method has recently been developed. This assay is faster and more sensitive than real-time RT-PCR but had specificity only for NDV genotype VII (Selim et al., 2022).

Pathogenicity testing

Based on calculations of the intracerebral pathogenicity index (ICPI) in day-old chicks, the intravenous pathogenicity index (IVPI) in six-week-old chickens, or the mean time to death in embryos. The NDV isolates can be characterized as virulent (velogenic), moderately virulent (mesogenic), or less virulent (lentogenic). However, ICPI appears to be the more accurate and sensitive method compared to the other two methods (Dortmans et al., 2011). The calculation of ICPI is done after intracerebral infection of day-old chicks with a score ranging from 0 to 2, where 0 is assigned to a normal chick, 1 to a sick chick, and 2 to a dead chick. This score is assigned daily to each chick for eight

days. Finally, the virulence of the virus strain is determined by the average of the scores. Strains are velogenic when the ICPI value is greater than 1.5. They are mesogenic when the ICPI value is between 0.7 and 1.5. Finally, they are lentogenic when the ICPI value is less than 0.7. The IVPI is calculated in the same way as the ICPI, but the six-week-old birds are injected intravenously. The presence of a velogenic strain is indicated by an IVPI greater than 2.5 (Al-Shammari et al., 2020). Mean time to death refers to the average time in hours required for all inoculated embryos to die. If the embryos die within 60 hours, it is an indication of the presence of velogenic strains, while survival of the embryos beyond 90 hours indicates the presence of lentogenic strains (Cattoli et al., 2011). *In vitro*, NDV induces plaque formation in embryonic fibroblast culture, the size and morphology of which varies according to the virulence of the viral strain. The amino acid sequence of the F protein cleavage site can also be used in differentiating the virulence of NDV isolates (Awad et al., 2020).

Control

Control of Newcastle disease must include strict biosecurity that will prevent velogenic virus strains from coming into contact with poultry and adequate administration of effective vaccines (Dimitrov et al., 2017). In addition, serological monitoring of the immunological status of vaccinated flocks should be performed to assess the antibody response to the administered vaccines (Ahmed and Odisho, 2018). Traditional vaccines (live and inactivated vaccines) have been widely used for several decades (Dimitrov et al., 2017). However, another generation of vaccines, such as recombinant vaccines and antigen-matching vaccines, have recently been adopted in some countries, and other vaccine approaches are in the experimental phase (Dimitrov et al., 2017; Nurzajah et al., 2022). The vaccine strains used are obtained from the naturally apathogenic, low pathogenic, or medium pathogenic strains (Dimitrov et al., 2017; Hu et al., 2022). These vaccine strains belong to genotype I (Ulster and V4), genotype II (Lasota and B1), genotype III (Mukteshwar), genotype IV (Herts/33, UK), genotype V (Anhinga, USA), genotype VIII (AF22440), and genotype XV (vaccine strains, China, Bello et al., 2018).

With the development of genetic engineering, NDV vaccine strains have emerged as promising vectors for developing effective multipurpose vaccines against pathogens that infect both animals and humans (Kim and Samal, 2016; Choi, 2017). For example, the use of an NDV virus-like particle-based vaccine expressing NDV F protein and influenza virus matrix 1 (M1) protein could protect chickens from a lethal challenge, while significantly reducing viral shedding and can also be used as a DIVA strategy to differentiate infected from vaccinated animals (Park et al., 2014). An experiment conducted by Izquierdo-Lara et al. (2019) indicated that vaccine strains paired with wild type XII strains conferred better protection to vaccinated animals and significantly reduced viral shedding. Similarly, Miller et al. (2013) in an experimental setting showed that by increasing the homology of the vaccine to a wild-type virus, humoral antibody efficacy levels could be increased. As the level of humoral antibodies increases in vaccinated birds, the number of infected birds and the amount of virus shed decreases. Results of an experiment conducted by Pandarangga et al. (2022) also revealed that viral shedding in chickens is reduced when vaccinated with strains homologous to the infectious challenge strain compared to the vaccine using the heterologous strain (LaSota).

Furthermore, in an intranasal vaccination trial in chickens, Zhao et al. (2016) found that a DNA vaccine containing the NDV *F* gene encapsulated in hollow nanoparticles induced higher serum antibody (IgA) titers. Results of a comparative study by Zhao et al. (2012) also showed that a live vaccine (LaSota strain) encapsulated in chitosan nanoparticles (NDV-CS-NP) induced better protection to immunized specific pathogen-free chickens, compared to the live LaSota strain vaccine and the inactivated NDV vaccine. An encapsulated inactivated NDV vaccine in chitosan nanoparticles has also been successfully developed (Mohammadi et al., 2021). These studies lay the foundation for the future development of mucosal vaccines encapsulated in nanoparticles. Plant-derived vaccines are in development and offer potential advantages over traditional vaccines, including a significant reduction in viral shedding and the ability to differentiate infected birds from vaccinated birds (Nurzajah et al., 2022; Smith et al., 2023).

CONCLUSION

Newcastle disease is a health threat to the global poultry industry. It is endemic in many developing countries. In disease-free countries, NDV outbreaks are occasional, but when they occur, they result in significant trade restrictions and embargoes. Newcastle disease is sometimes misdiagnosed because serological and molecular diagnostic tests for NDV have certain limitations as a result of cross-reactivity of NDV with other homologous avian paramyxoviruses and variability in the nucleotide sequence of the NDV *F* gene. To prevent ND, only sanitary and medical prophylactic measures can be used effectively to enhance the immune capacities of birds and reduce ambient viral pressure. The international reports show that vaccinated poultry can be infected with wild viruses. The NDV genotypes are emerging, which can make control strategies complex and often difficult, hence the need to use high-quality and up-to-date vaccines. Research priorities should therefore focus on improving diagnostic tools and developing better vaccines.

DECLARATIONS

Acknowledgments

This study was supported by the Regional Centre of Excellence on Avian Science of the University of Lomé, in Togo. The authors wish to express their warm gratitude to World Bank IDA 5424, which is the main sponsor of CERSA.

Authors' contributions

This work was carried out with the contribution of all authors. Ahamidou Moustapha drafted the manuscript. Essodina Talaki, Akourki Adamou, and Moumouni Ousseini revised the manuscript. All authors read and approved the final version of the manuscript.

Competing interests

The authors have not declared any conflict of interest.

Ethical consideration

The authors of the current study checked for ethical issues, including plagiarism, consent to publish, misconduct, double publication and/or submission, and redundancy.

Funding

Not applicable.

Data availability and materials

Data from the study are available according to a reasonable request.

REFERENCES

- Abd Elfatah KS, Elabasy MA, El-khyate F, Elmahallawy EK, Mosad SM, El-Gohary FA, Abdo W, Al-Brakati A, Seadawy MG, Tahoon AE et al. (2021). Molecular characterization of velogenic Newcastle disease virus (sub-genotype VII.1.1) from wild birds, with assessment of its pathogenicity in susceptible chickens. *Animals*, 11: 505. DOI: <https://www.doi.org/10.3390/ani11020505>
- Abdisa T and Tagesu T (2017). Review on Newcastle disease of poultry and its public health importance. *Journal of Veterinary Science and Technology*, 8(3): 1000441. Available at: <https://www.hilarispublisher.com/open-access/review-on-newcastle-disease-of-poultry-and-its-public-health-importance-2157-7579-1000441.pdf>
- Ahmed AI and Odisho ShM (2018). Isolation identification and pathotyping of Newcastle disease viruses form naturally infected chickens in Iraqi Kurdistan region. *The Iraqi Journal of Agricultural Sciences*, 49(1): 132-141. Available at: <https://archive-jcoagri.uobaghdad.edu.iq/wp-content/uploads/sites/8/uploads/Latest%20Issue/2018/2018-1/17.pdf>
- Alazawy AK and Al Ajeeli KS (2020). Isolation and molecular identification of wild Newcastle disease virus isolated from broiler farms of Diyala Province, Iraq. *Veterinary World*, 13(1): 33-39. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7020111/pdf/Vetworld-13-33.pdf>
- Ali AAH, Abdallah FM, Farag GK, and Fatehi S (2022). An overview on virological pathogenesis, clinical manifestation, and molecular studies on newcastle disease virus. *European Journal of Molecular & Clinical Medicine*, 9(4): 2807-2824. Available at: https://ejmcm.com/pdf_19147_252376f6f087e9fbf06385528c6f4108.html
- Al-Shammari AM, Hamad MA, AL-Mudhafar MA, Raad K, and Ahmed A (2020). Clinical, molecular and cytopathological characterization of a Newcastle disease virus from an outbreak in Baghdad, Iraq. *Veterinary Medicine and Science*, 6: 477-484. Available at: <https://onlinelibrary.wiley.com/doi/full/10.1002/vms3.262>
- Amarasinghe GK, Ayllón MA, Bào Y, Basler CF, Bavari S, Blasdel KR, Briese T, Brown PA, Bukreyev A, Balkema-Buschmann A et al. (2019). Taxonomy of the order Mononegavirales: Update 2019. *Archives of Virology*, 164: 1967-1980. DOI: <https://www.doi.org/10.1007/s00705-019-04247-4>
- Ana PS, Emily JA, Gaspar HC, Ashley B, Amandus PM, Boniface BK, Terra K, Huaijun Z, and Rodrigo AG (2020). Molecular characterization of Newcastle disease viruses isolated from chickens in Tanzania and Ghana. *Viruses*, 12(9): 916. DOI: <https://www.doi.org/10.3390/v12090916>
- Awad MM, Mosad SM, and El-Kenawy AA (2020). Molecular differentiation between velogenic isolates and lentogenic LaSota strain of Newcastle disease virus. *Mansoura Veterinary Medical Journal*, 21(4): 208-216. Available at: https://mvmj.journals.ekb.eg/article_146340.html
- Azizah AN, Anwar Ch, and Rahardjo AP (2021). Isolation and identification of Newcastle disease virus from ducks sold at traditional livestock market center in Indonesia. *Journal of World's Poultry Research*, 11(1): 96-100. DOI: <https://www.doi.org/10.36380/jwpr.2021.12>
- Bello MB, Yusoff KM, Ideris A, Hair-Bejo M, Peeters BPH, Jibril AH, Tambuwal FM, and Omar AR (2018). Genotype diversity of Newcastle disease virus in Nigeria: Disease control challenges and future outlook. *Advances in Virology*, 2018: 6097291. DOI: <https://www.doi.org/10.1155/2018/6097291>
- Brown VR and Bevins SN (2017). A review of virulent Newcastle disease viruses in the United States and the role of wild birds in viral persistence and spread. *Veterinary Research*, 48: 68. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5659000/pdf/13567_2017_Article_475.pdf
- Cao Y, Gu M, Zhang X, Liu W, and Liu X (2013). Complete genome sequences of two Newcastle disease virus strains of genotype VIII. *Genome Announcements Journal*, 1(1): e00180-12. DOI: <https://www.doi.org/10.1128/genomeA.00180-12>
- Caroline VS (2022). Newcastle disease virus (NDV) in poultry birds. *African Journal of Poultry Farming*, 10(2): 1-2. Available at: <https://www.internationalscholarsjournals.com/articles/newcastle-disease-virus-ndv-in-poultry-birds.pdf>
- Cattoli G, Susta L, Terregino C, and Brown C (2011). Newcastle disease: A review of field recognition and current methods of laboratory detection. *Journal of Veterinary Diagnostic Investigation*, 23(4): 637-656. DOI: <https://www.doi.org/10.1177/1040638711407887>

- Chen X, Jia Y, Wei N, Ye C, Hao H, Xiao S, Wang X, Liu H, and Yang Z (2021). Identification of a new amino acid mutation in the HN protein of NDV involved in pathogenicity. *Veterinary Research*, 52: 147. DOI: <https://www.doi.org/10.1186/s13567-021-01019-4>
- Choi KS (2017). Newcastle disease virus vectored vaccines as bivalent or antigen delivery vaccines. *Clinical and Experimental Vaccine Research*, 6(2): 72-82. DOI: <https://www.doi.org/10.7774/cevr.2017.6.2.72>
- Chumbe A, Izquierdo-Lara R, Tataje L, Gonzalez R, Cribillero G, González AE, Fernández-Díaz M, and Icochea E (2017). Pathotyping and phylogenetic characterization of Newcastle disease viruses isolated in Peru. *Avian Diseases*, 61(1): 16-24. Available at: <https://www.jstor.org/stable/26432216>
- Das M and Kumar S (2016). Evidence of independent evolution of genotype XIII Newcastle disease viruses in India. *Archives of Virology*, 162: 997-1007. DOI: <https://www.doi.org/10.1007/s00705-016-3182-3>
- Denis KB, Kizito KM, John BO, Martin O, Agnes W, Maxwell OO, Halid K, Jessika LN, Angélique T, Mathilde P et al. (2015). Virulence et faible évolution génétique de virus de la maladie de Newcastle vélogènes isolés en Ouganda [Virulence and low genetic evolution of velogenic Newcastle disease viruses isolated in Uganda]. 11èmes Journées de la Recherches Avicole, Tours, France. Available at: <https://hal.inrae.fr/hal-02739493/document>
- Diel DG, da Silva LHA, Liu H, Wang Z, Miller PJ, and Afonso CL (2012). Genetic diversity of avian paramyxovirus type 1: Proposal for a unified nomenclature and classification system of Newcastle disease virus genotypes. *Infection, Genetics and Evolution*, 12(8): 1770-1779. DOI: <https://www.doi.org/10.1016/j.meegid.2012.07.012>
- Dimitrov KM, Abolnik C, Afonso CL, Albina E, Bahl J, Berg M, Briand FX, Brown IH, Choi KS, Chvala I et al (2019). Updated unified phylogenetic classification system and revised nomenclature for Newcastle disease virus. *Infection, Genetics and Evolution*, 74: 103917. DOI: <https://www.doi.org/10.1016/j.meegid.2019.103917>
- Dimitrov KM, Afonso CL, Yu Q, and Miller PJ (2017). Newcastle disease vaccines-A solved problem or a continuous challenge?. *Veterinary Microbiology*, 206: 126-136. DOI: <http://www.doi.org/10.1016/j.vetmic.2016.12.019>
- Dimitrov KM, Ramey AM, Qiu X, Bahl J, and Afonso CL (2016). Temporal, geographic, and host distribution of avian paramyxovirus 1 (Newcastle disease virus). *Infection, Genetics and Evolution*, 39: 22-34. DOI: <https://www.doi.org/10.1016/j.meegid.2016.01.008>
- Dortmans JCFM, Koch G, Rottier PJM, and Peeters BPH (2011). Virulence of Newcastle disease virus: What is known so far?. *Veterinary Research*, 42: 122. Available at: <http://www.veterinaryresearch.org/content/42/1/122>
- Duan X, Zhang P, Ma J, Chen S, Hao H, Liu H, Fu X, Wu P, Zhang D, Zhang W et al. (2014). Characterization of genotype IX Newcastle disease virus strains isolated from wild birds in the Northern Qinling Mountains, China. *Virus Genes*, 48: 48-55. DOI: <https://www.doi.org/10.1007/s11262-013-0987-y>
- Duan Z, Deng S, Ji X, Zhao J, Yuan C, and Gao H (2019). Nuclear localization of Newcastle disease virus matrix protein promotes virus replication by affecting viral RNA synthesis and transcription and inhibiting host cell transcription. *Veterinary Research*, 50(22): 1-9. DOI: <https://www.doi.org/10.1186/s13567-019-0640-4>
- Dzozbema KFX, Talaki E, Batawui KB, and Dao BB (2021). Review on Newcastle disease in poultry. *International Journal of Biological and Chemical Sciences*, 15(2): 773-789. DOI: <https://www.doi.org/10.4314/ijbcs.v15i2.29>
- Ecco R, Susta L, Afonso CL, Miller PJ, and Brown C (2011). Neurological lesions in chickens experimentally infected with virulent Newcastle disease virus isolates. *Avian Pathology*, 40(2): 145-152. DOI: <https://www.doi.org/10.1080/03079457.2010.544289>
- Fellahi S and Boudouma F (2021). Virus de la maladie de Newcastle chez la volaille: Perspectives actuelles et émergentes [Newcastle disease virus in poultry: Current and emerging perspectives]. *Revue Marocaine des Sciences Agronomiques et Vétérinaires*, 9(3): 453-460. Available at: https://www.agromaroc.com/index.php/Actes_IAPH2/article/view/1021
- Ganar K, Das M, Sinha S, and Kumar S (2014). Newcastle disease virus: Current status and our understanding. *Virus Research*, 184: 71-81. DOI: <http://www.doi.org/10.1016/j.virusres.2014.02.016>
- Getabalew M, Alemneh T, Akeberegn D, Getahun D, and Zewdie D (2019). Epidemiology, diagnosis & prevention of Newcastle disease in poultry. *American Journal of Biomedical Science and Research*, 3(1): 50-59. DOI: <https://www.doi.org/10.34297/AJBSR.2019.03.000632>
- Heiden S, Grund C, Röder A, Granzow H, Kühnel D, Mettenleiter TC, and Römer-Oberdörfer A (2014). Different regions of the Newcastle disease virus fusion protein modulate pathogenicity. *PLoS ONE*, 9(12): e113344. DOI: <https://www.doi.org/10.1371/journal.pone.0113344>
- Hu Z, He X, Deng J, Hu J, and Liu X (2022). Current situation and future direction of Newcastle disease vaccines. *Veterinary Research*, 53: 99. DOI: <https://www.doi.org/10.1186/s13567-022-01118-w>
- Izquierdo-Lara R, Chumbe A, Calderón K, Fernández-Díaz M, and Vakharia VN (2019). Genotype matched Newcastle disease virus vaccine confers improved protection against genotype XII challenge: The importance of cytoplasmic tails in viral replication and vaccine design. *PLoS ONE*, 14(11): e0209539. Available at: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0209539>
- Jamil F, Aslam L, Laraib, Ali H, Shoukat K, Rasheed MA, Raza S, and Ibrahim M (2022). An *in silico* study of derivative of Newcastle disease virus epitopes based vaccine against hemagglutinin neuraminidase protein. *Journal of Animal Science*, 101: skac375, DOI: <https://www.doi.org/10.1093/jas/skac375>
- Kim SH and Samal SK (2016). Newcastle disease virus as a vaccine vector for development of human and veterinary vaccines. *Viruses*, 8(7): 183. DOI: <https://www.doi.org/10.3390/v8070183>
- Liu H, Lv Y, Afonso CL, Ge S, Zheng D, Zhao Y, and Wang Z (2013). Complete genome sequences of new emerging Newcastle disease virus strains isolated from China. *Genome Announc*, 1(1): e00129-12. Available at: <https://journals.asm.org/doi/pdf/10.1128/genomeA.00129-12>
- Mahamud SNA, Bello MB, Ideris A, and Omar AR (2022). Efficacy of genotype-matched Newcastle disease virus vaccine formulated in carboxymethyl sago starch acid hydrogel in chickens vaccinated via different routes. *Journal of Veterinary Science*, 23(4): e25. DOI: <https://www.doi.org/10.4142/jvs.21242>
- Maminaiina OF, Gil P, Briand FX, Albina E, Keita D, Andriamanivo HR, Chevalier V, Lancelot R, Martinez D, Rakotondravao R et al. (2010). Newcastle disease virus in Madagascar: Identification of an original genotype possibly deriving from a died out ancestor of genotype IV. *PLoS ONE*, 5(11): e13987. DOI: <https://www.doi.org/10.1371/journal.pone.0013987>
- Mao Q, Ma S, Schrickel PL, Zhao P, Wang J, Zhang Y, Li S, and Wang C (2022). Review detection of Newcastle disease virus. *Frontiers in Veterinary Science*, 9: 936251. DOI: <https://www.doi.org/10.3389/fvets.2022.936251>
- Mariappan AK, Munusamy P, Kumar D, Latheef SK, Singh SD, Singh R, and Dhama K (2018). Pathological and molecular investigation of velogenic viscerotropic Newcastle disease outbreak in a vaccinated chicken flocks. *Virus Disease*, 29(2): 180-191. DOI: <https://www.doi.org/10.1007/s13337-018-0445-5>
- Martín JJ, Holguera J, Sánchez-Felipe L, Villar E, and Muñoz-Barroso I (2012). Cholesterol dependence of Newcastle disease virus entry. *Biochimica et Biophysica Acta*, 1818(3): 753-761. DOI: <https://www.doi.org/10.1016/j.bbame.2011.12.004>
- Megahed M, Mohamed W, and Hassanin O (2020). A complex genetic diversity of newcastle disease virus (ndv) in Africa continent: An updated review. *Journal of Animal Health and Production*, 9(s1): 97-109. DOI: <http://www.doi.org/10.17582/journal.jahp/2020/9.s1.97.109>

- Miller PJ and Koch G (2013). Newcastle disease. In: D. E. Swayne, J. R. Glisson, L. R. McDougald, L. K. Nolan, D. L. Suarez, V. L. Nair (Editors), *Diseases of Poultry*, 13th Edition. Wiley-Blackwell, Ames, IA, pp. 89-107, 120-130. Available at: <https://www.ars.usda.gov/research/publications/publication/?seqNo115=286542>
- Miller PJ, Afonso CL, El Attache J, Dorsey KM, Courtney SC, Guo Z, and Kapczynski DR (2013). Effects of Newcastle disease virus vaccine antibodies on the shedding and transmission of challenge viruses. *Developmental and Comparative Immunology*, 41(4): 505-513. DOI: <http://www.doi.org/10.1016/j.dci.2013.06.007>
- Miller PJ, Decanini EL, and Afonso CL (2010). Newcastle disease: Evolution of genotypes and the related diagnostic challenges. *Infection, Genetics and Evolution*, 10(1): 26-35. DOI: <https://www.doi.org/10.1016/j.meegid.2009.09.012>
- Miller PJ, Haddas R, Simanov L, Lublin A, Rehmani SF, Wajid A, Bibi T, Khan TA, Yaqub T, Setiyaningsih S et al. (2015). Identification of new sub-genotypes of virulent Newcastle disease virus with potential panzootic features. *Infection, Genetics and Evolution*, 29: 216-229. DOI: <https://www.doi.org/10.1016/j.meegid.2014.10.032>
- Mohammadi AR, Zamani MA, and Shahsavandi S (2021). Encapsulation of inactivated Newcastle disease virus onto the chitosan nanoparticles for use in mucosal immunity. *Iranian Journal of Virology*, 15(1): 32-39. Available at: <https://journal.isv.org.ir/article-1-417-fa.pdf>
- Naguib MM, Höper D, Elkady MF, Afifi MA, Erfan A, Abozeid HH, Hasan WM, Arafa AS, Shahein M, Beer M et al. (2021). Comparison of genomic and antigenic properties of Newcastle disease virus genotypes II, XXI and VII from Egypt do not point to antigenic drift as selection marker. *Transboundary and Emerging Diseases*, 69(2): 849-863. DOI: <https://www.doi.org/10.1111/tbed.14121>
- Naz D, Rahman S, Aslam MA, and Muhammad F (2022). Newcastle disease virus in poultry with an interface as a human vector. *Veterinary Vaccine*, 1(1): 100003. DOI: <https://www.doi.org/10.1016/j.vetvac.2022.100003>
- Nurzijah I, Elbohy OA, Kanyuka K, Daly JM, and Dunham S (2022). Development of plant-based vaccines for prevention of avian influenza and Newcastle disease in poultry. *Vaccines*, 10(3): 478. DOI: <https://www.doi.org/10.3390/vaccines10030478>
- Pandarangga P, McAllister MM, Peaston AE, Ngai YT, Cahyono MI, and Hemmatzadeh F (2022). Performance comparison of homologous and heterologous Newcastle disease virus in vaccines and antibody tests. *Research in Veterinary Science*, 149: 82-89. DOI: <https://www.doi.org/10.1016/j.rvsc.2022.06.014>
- Park JK, Lee DH, Yuk SS, Tseren-Ochir EO, Kwon JH, Noh JY, Kim BY, Choi SW, Kang SM, Lee JB et al. (2014). Virus-like particle vaccine confers protection against a lethal Newcastle disease virus challenge in chickens and allows a strategy of differentiating infected from vaccinated animals. *Clinical and Vaccine Immunology*, 21(3): 360-365. DOI: <https://www.doi.org/10.1128/CVI.00636-13>
- Peng T, Qiu X, Tan L, Yu S, Yang B, Dai J, Liu X, Sun Y, Song C, Liu W et al. (2022). Ubiquitination on lysine 247 of Newcastle disease virus matrix protein enhances viral replication and virulence by driving nuclear-cytoplasmic trafficking. *Journal of Virology*, 96(2): e01629-21. Available at: <https://journals.asm.org/doi/epub/10.1128/JVI.01629-21>
- Phale S (2018). Newcastle disease virus: Structural and molecular basis of pathogenicity. *Medicinal Chemistry*, 8(8): 202-204. Available at: <https://www.hilarispublisher.com/open-access/newcastle-disease-virus-structural-and-molecular-basis-of-pathogenicity-2161-0444-1000514.pdf>
- Pinette MM, Rodriguez-Lecompte JC, Pasick J, Ojkie D, Leith M, Suderman M, and Berhane YJ (2014). Development of a duplex fluorescent microsphere immunoassay (FMIA) for the detection of antibody responses to influenza A and newcastle disease viruses. *Journal of Immunological Methods*, 405: 167-177. DOI: <https://www.doi.org/10.1016/j.jim.2014.02.004>
- Puro K and Sen A (2022). Newcastle disease in backyard poultry rearing in the Northeastern States of India: Challenges and control strategies. *Frontiers in Veterinary Science*, 9: 799813. Available at: <https://www.frontiersin.org/articles/10.3389/fvets.2022.799813/full>
- Putri DD, Handharyani E, Soejoedono RD, Setiyono A, Mayasari NI, and Poetri ON (2017). Pathotypic characterization of Newcastle disease virus isolated from vaccinated chicken in West Java, Indonesia. *Veterinary World*, 10(4): 438-444. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5422248/pdf/VetWorld-10-438.pdf>
- Rahman A, Habib M, and Shabbir MZ (2018). Adaptation of Newcastle disease virus (NDV) in feral birds and their potential role in interspecies transmission. *The Open Virology Journal*, 12: 52-68. DOI: <https://www.doi.org/10.2174%2F1874357901812010052>
- Ramsubeik S, Crossley B, Jerry C, Bland M, Rejmanek D, Shivaprasad HL, and Stoute S (2023). Molecular and retrospective analysis of pigeon paramyxovirus type1 infections in confinement-reared pigeons (Columbia livia); 2010-2020. *Journal of Applied Poultry Research*, 32(2): 100343. DOI: <https://www.doi.org/10.1016/j.japr.2023.100343>
- Rasamoelina AH, Molia S, Razafindraibe NP, Andria-Mananjara DE, Rakotomanana OD, and Chevalier V (2016). The Newcastle disease. In: H. Duchaufour, T. Razafimbelo, J. Rakotoarisoa, B. Ramamonjisoa, and Rakotondravao (Editors), *Recherche interdisciplinaire pour le développement durable et la biodiversité des espaces ruraux malgaches. Application à différentes thématiques de territoire [Interdisciplinary research for sustainable development]*. Actes du projet FSP Parrur. Available at: <http://www.apdra.org/IMG/file/Collectif%20Parrur%20web.pdf>
- Rauw F, Gardin Y, van den Berg T, and Lambrecht B (2009). La vaccination contre la maladie de Newcastle chez le poulet (*Gallus gallus*) [Vaccination against Newcastle disease in chickens (*Gallus gallus*)]. *Biotechnologie, Agronomie, Société et Environnement*, 13(4): 587-596. Available at: <https://popups.uliege.be/1780-4507/index.php?id=4758>
- Rehman ZU, Meng C, Sun Y, Mahrose KM, Umar S, Ding C, and Munir M (2018). Pathobiology of avian avulavirus 1: Special focus on waterfowl. *Veterinary Research*, 49: 94. Available at: <https://veterinaryresearch.biomedcentral.com/articles/10.1186/s13567-018-0587-x>
- Samal S, Khattar SK, Kumar S, Collins PL, and Samal SK (2012). Coordinate deletion of N-glycans from the heptad repeats of the fusion F protein of Newcastle disease virus yields a hyperfusogenic virus with increased replication, virulence, and immunogenicity. *Journal of Virology*, 86(5): 2501-2511. DOI: <https://www.doi.org/10.1128/JVI.06380-11>
- Samal S, Khattar SK, Paldurai A, Palaniyandi S, Zhu X, Collins PL, and Samal SK (2013). Mutations in the cytoplasmic domain of the Newcastle disease virus fusion protein confer hyperfusogenic phenotypes modulating viral replication and pathogenicity. *Journal of Virology*, 87(18): 10083-10093. DOI: <https://www.doi.org/10.1128/jvi.01446-13>
- Samal S, Kumar S, Khattar SK, and Samal SK (2011). A single amino acid change, Q114R, in the cleavage-site sequence of Newcastle disease virus fusion protein attenuates viral replication and pathogenicity. *Journal of General Virology*, 92: 2333-2338. DOI: <https://www.doi.org/10.1099/vir.0.033399-0>
- Samuel A, Nayak B, Paldurai A, Xiao S, Aplogan GL, Awoume KA, Webby RJ, Ducatez MF, Collins PL, and Samal SK (2013). Phylogenetic and pathotypic characterization of Newcastle disease viruses circulating in West Africa and efficacy of a current vaccine. *Journal of Clinical Microbiology*, 51(3): 771-781. DOI: <https://www.doi.org/10.1128%2FJCM.02750-12>
- Sánchez-Felipe L, Villar E, and Muñoz-Barroso I (2014). Entry of Newcastle disease virus into the host cell: Role of acidic pH and endocytosis. *Biochimica et Biophysica Acta*, 1838(1): 300-309. DOI: <http://www.doi.org/10.1016/j.bbame.2013.08.008>
- Selim K, Adel A, Eid S, and Shahein M (2022). Development of real time reverse transcription loop-mediated isothermal amplification assay for rapid detection of genotype VII of Newcastle disease viruses. *British Poultry Science*, DOI: <https://www.doi.org/10.1080/00071668.2022.2094219>

- Shabbir MZ, Nissly RH, Ahad A, Rabbani M, Lim L, Chothe SK, Subbiah M, Sebastian A, Albert I, Ul-Rahman A et al. (2021). Genomic characterization of velogenic avian orthoavulavirus 1 isolates from poultry workers: Implications to emergence and its zoonotic potential towards public health. *Asian Pacific Journal of Tropical Medicine*, 14: 64-72. Available at: <https://www.apjtm.org/article.asp?issn=19957645;year=2021;volume=14;issue=2;spage=64;epage=72;aulast=Shabbir>
- Shakal M, Maher M, Metwally AS, AbdelSabour MA, Madbbouly YM, and Safwat G (2020). Molecular identification of a velogenic Newcastle disease virus strain isolated from Egypt. *Journal of World's Poultry Research*, 10(2S): 195-202. DOI: <https://www.doi.org/10.36380/jwpr.2020.25>
- Smith T, O'Kennedy MM, Ross CS, Lewis NS, and Abolnik C (2023). The production of Newcastle disease virus-like particles in *Nicotiana benthamiana* as potential vaccines. *Frontiers in Plant Science*, 14: 1130910. DOI: <https://www.doi.org/10.3389/fpls.2023.1130910>
- Snoeck CJ, Owoade AA, Couacy-Hymann E, Alkali BR, Okwen MP, Adeyanju AT, Komoyo GF, Nakoune E, Le Faou A, and Muller CP (2013). High genetic diversity of Newcastle disease virus in poultry in West and Central Africa: Cocirculation of genotype XIV and newly defined genotypes XVII and XVIII. *Journal of Clinical Microbiology*, 51(7): 2250-2260. DOI: <https://www.doi.org/10.1128/JCM.00684-13>
- Sonkusale PM, Ghosh RC, Giri DK, Kumar V, Shukla N, Gumasta P, and Bisen S (2023). Pathology and molecular diagnosis of velogenic viscerotropic Newcastle disease virus in Desi chickens of Chhattisgarh. *The Indian Journal of Veterinary Sciences and Biotechnology*, 19(1): 120-123. Available at: <https://acspublisher.com/journals/index.php/jvsbt/article/view/3440/3837>
- Souley MM, Ibrahim AI, Bachir SK, Alassane A, Rahila I, Bachir Y, Hadiza A, Hama H, Hassane A, Nourou A et al. (2021). Co-circulation of genotypes XIV.2 and XVIII.2 of avian paramyxovirus-1 (Newcastle disease virus) in backyard poultry in Niger. *Virus Genes*, 57: 100-105. DOI: <https://www.doi.org/10.1007/s11262-020-01804-x>
- Ul-Rahman A and Shabbir MZA (2019). Comparative phylogenomic analysis of avian avulavirus 1 isolated from non-avian hosts: Conquering new frontiers of zoonotic potential among species. *Archives of Virology*, 164: 1771-1780. DOI: <https://www.doi.org/10.1007/s00705-019-04276-z>
- Ul-Rahman A, Ishaq HM, Raza MA, and Shabbir MZ (2022). Zoonotic potential of Newcastle disease virus: Old and novel perspectives related to public health. *Reviews in Medical Virology*, 32(1): e2246. DOI: <https://www.doi.org/10.1002/rmv.2246>
- Wehmann E, Ujvári D, Mazija H, Velhner M, Ciglar-Grozđanić I, Savić V, Jermolenko G, Čač Ž, Pukner-Radovčić E, and Lomniczi B (2003). Genetic analysis of Newcastle disease virus strains isolated in Bosnia-Herzegovina, Croatia, Slovenia and Yugoslavia, reveals the presence of only a single genotype, V, between 1979 and 2002. *Veterinary Microbiology*, 94(4): 269-281. DOI: [https://www.doi.org/10.1016/S0378-1135\(03\)00133-0](https://www.doi.org/10.1016/S0378-1135(03)00133-0)
- Worku T and Teshome I (2020). Review on the role of viral structural proteins on the pathogenicity of Newcastle disease virus in chickens. *American Journal of Zoology*, 3(2): 40-46. Available at: <https://www.sciencepublishinggroup.com/journal/paperinfo?journalid=657&doi=10.11648/j.ajz.20200302.12>
- World organisation of animal health (WOAH) (2022). Manual of diagnostic tests and vaccines for terrestrial animals. Manual of diagnostic tests and vaccines for terrestrial animals 2022. Available at: <https://www.woah.org/en/what-we-do/standards/codes-and-manuals/terrestrial-manual-online-access/>
- Xue C, Cong Y, Yin R, Sun Y, Ding C, Yu S, Liu X, Hu S, Qian J, Yuan Q et al. (2017). Genetic diversity of the genotype VII Newcastle disease virus: identification of a novel VIIj sub-genotype. *Virus Genes*, 53: 63-70. DOI: <https://www.doi.org/10.1007/s11262-016-1404-0>
- Zhao K, Chen G, Shi Xm, Gao TT, Li W, Zhao Y, Zhang FQ, Wu J, Cui X, and Wang YF (2012). Preparation and efficacy of a live Newcastle disease virus vaccine encapsulated in chitosan nanoparticles. *PLoS ONE*, 7(12): e53314. DOI: <https://www.doi.org/10.1371/journal.pone.0053314>
- Zhao K, Rong G, Hao Y, Yu L, Kang H, Wang X, Wang X, Jin Z, Ren Z, and Li Z (2016). IgA response and protection following nasal vaccination of chickens with Newcastle disease virus DNA vaccine nanoencapsulated with Ag@SiO₂ hollow nanoparticles. *Scientific Reports*, 6: 25720. DOI: <https://www.doi.org/10.1038/srep25720>

Publisher's note: [Scienceline Publication](#) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023