



Potential Benefits of Propolis in Large and Small Animal Practices: A Narrative Review of the Literature

Ashraf M. Abu-Seida 

Department of Surgery, Anesthesiology and Radiology, Faculty of Veterinary Medicine, Cairo University, Giza, PO: 12211, Egypt

*Corresponding author's Email: ashrafseida@cu.edu.eg

REVIEW ARTICLE
pII: S232245682300048-13
Received: 20 June 2023
Accepted: 08 August 2023

ABSTRACT

Propolis is a resinous substance from a mixture of different plant parts and molecules bees compose. This narrative review article explored the application of propolis in large and small animal practices in PubMed, Scopus, and Google Scholar databases. Propolis is applied in different pharmaceutical forms. Due to its numerous biological actions, such as antimicrobial, anti-inflammatory, antioxidant, antiparasitic, antiulcer, antitumor, and immunomodulatory, propolis can improve animal health and production. Propolis could be used as an alternative treatment for many diseases, such as mastitis, lumpy skin disease, foot and mouth disease, reproductive disorders, and diarrhea in cattle. Moreover, it could improve weight gain in cattle. In equine, propolis has been used as a local anesthetic and for treating dermatomycosis, chronic bronchitis, and skin wounds. In pigs, propolis has been used to treat enzootic pneumonia and as a prophylaxis for gastrointestinal and respiratory diseases in weak pigs. Propolis has been applied to treat caseous lymphadenitis and parasitic diseases in sheep and goats. Furthermore, it improves the immune status of kids and the health status of late pregnant ewes. In dogs and cats, propolis has been applied to treat otitis externa, eye diseases, Cushing's syndrome, and dermatophytosis. In dogs, propolis can treat transmissible venereal tumors. Moreover, propolis positively affects animal production, average daily gain and milk yield in sheep, growth of calves, lambs, and piglets, and cow's milk nutritional quality. On the other hand, the addition of propolis to the diet of feedlot bulls and pigs has no effect on their feed intake, hematological, biochemical, and immunological parameters, nutrient digestibility, microbial synthesis, and carcass characteristics. Based on the available clinical studies, propolis has potential benefits for animal health in cattle, equine, sheep, goats, pigs, dogs, and cats. According to the available literature, propolis is a natural promising agent that can alternate conventional pharmaceuticals, particularly antibiotics. It improves animal health and production with no adverse effects and low cost. Most conducted studies on the efficacy of propolis on animal health and production are *in vitro*. Due to its scarcity, further controlled clinical trials are recommended to evaluate the exact usefulness of propolis in veterinary medicine and to obtain reliable conclusions on the benefits of propolis in animal health and production.

Keywords: Cattle, Dog, Horse, Pig, Propolis, Sheep

INTRODUCTION

The term "propolis" originates from the Greek language, where "pro" signifies "in front" and "polis" means "city." This combination conveys the concept of "in front of the city," which aptly characterizes the protective role of propolis within the bee colony. Propolis serves to safeguard the hive by sealing openings and crevices (Bogdanov, 2017). Moreover, propolis also serves as a protective shield for the bee colony due to its antibacterial and antifungal properties, which contribute to shielding the colony from diseases (Bogdanov, 2017; Özdemir et al., 2022).

Propolis has other common names, such as propóleos, bee glue, bee propolis, hive dross, propolis balsam, propolis cera, propolis wax, propolis resin, Russian penicillin, and synthetic beeswax (Farooqui and Farooqui, 2010; Righi et al., 2011; Bogdanov, 2017; Özdemir et al., 2022).

Propolis consists of various quantities of beeswax and resins collected by the honeybee from different plants' flowers, and leaf buds (Bogdanov, 2017). The chemical composition of propolis varies according to its geographic and botanical origin. Nevertheless, the pharmacological effects of different types of propolis are the same (Marcucci, 1995). Therefore, it is used in alternative medicine like wound and burn dressing and as a nutritional supplement due to its unique biological actions (Elshater et al., 2017; Elkhenany et al., 2019).

Propolis has antibacterial, antifungal, antiviral, antiparasitic, anti-inflammatory, immuno-stimulatory, anti-tumoral, local anesthetic, and antioxidant activities (Betancourt et al., 2015; Bogdanov, 2017; González-Búrquez et al., 2017). Therefore, propolis represents a promising medicine and natural supplement in the diet for supporting body activities without adverse effects on either animals or the environment. Moreover, propolis is considered a natural alternative to many pharmaceutical drugs to overcome the drug residue problem in food, such as antibiotics residue (Banskota et al., 2001; Özdemir et al., 2022).

This narrative review aimed to describe the different clinical applications of propolis in large and small animal practices and its potential benefits in animal health and production.

DATA COLLECTION

This review relied on an extensive literature search conducted in January 2023, encompassing the utilization of propolis in both large and small animals. The search specifically targeted relevant articles published in the English language. Literature pertinent to propolis in veterinary medicine during the last 40 years (1982-2022) was explored in PubMed, Scopus, and Google Scholar databases. The relevant literature was reviewed and critically appraised in this review. The search terms included “propolis”, “horse”, “donkey”, “equine” “cattle”, “buffaloes”, “bovine”, “sheep”, “goats”, “dogs”, “cats”, and “pigs”.

PHYSICOCHEMICAL PROPERTIES OF PROPOLIS

Propolis has various colors and types with no standard chemical contents due to the variation in botanical origin, season, and types of bees (Marcucci, 1995; Özdemir et al., 2022). Its color ranges from transparent or yellow to dark brown according to the source of resin. Ethanol (ethyl alcohol) ether, glycol, and water are commonly used to extract the propolis (Abdulkhani et al., 2017; Anjum et al., 2019).

Propolis has more than 300 compounds with various compositions and isomers. Among these compounds, vitamins (C, B, B1, B2, A, and E), acids (organic acid, gallic acid, isoferulic acid, ferulic acid, and phenolic acids), flavonoids (flavones, flavonol, flavonones, and flavononol), cafeique, pectolinarigenine, chrysin, acacetin, coumarins, vaniline, pinocembrin, tectochrysin, galangin, izalpinin, kaempferol, quercetin, rhamnocitrin, pinostrobil, sakuranetin, pinobanksin, isovaniline, P-coumarin, cinnamin, scopoletin, and terpene, are the most common bioactive chemical agents present in all kinds of propolis (Boukraa, 2013; Bankova et al., 2014; Bogdanov, 2017; Anjum et al., 2019; Özdemir et al., 2022).

Propolis is composed of 50% resin compounds and balsams, 40% beeswax, 5% aromatic oils, and 5% bee pollen (Figure 1). Moreover, propolis is rich in albumin, calcium, magnesium, iron, zinc, silica, potassium, manganese, cobalt, copper, sodium, aluminum, nickel, chromium, and cadmium (Abdulkhani et al., 2017; Özdemir et al., 2022). Nevertheless, the various chemical compositions of propolis may induce a problem with its clinical application and quality control (Özdemir et al., 2022).




Figure 1. Chemical composition of raw propolis (Sources: Abdulkhani et al., 2017; Özdemir et al., 2022).

THERAPEUTIC ACTIONS OF PROPOLIS IN LARGE AND SMALL ANIMAL PRACTICES

Propolis is commonly applied in veterinary medicine for its antibacterial (El-Tayeb et al., 2019; Przybyłek and Karpiński, 2019), antifungal (de Mendonça et al., 2015), antiviral (Alp, 2018), antiparasitic (Anjum et al., 2019), antioxidant (Torres et al., 2018), anti-inflammatory (Machado et al., 2016), antitumor (Doğan et al., 2020), immunomodulatory (Shvarzbejn and Huleihel, 2011), antiulcer (da Silva et al., 2018), local anesthetic (Özdemir et al., 2022), and cytotoxic (Bonamigo et al., 2017) actions as shown in Figure 2. Moreover, propolis enhances wound healing (Abu-Seida, 2015), dentinogenesis after pulp capping (Saleh et al., 2016; Abo El-Mal et al., 2021), and hard-tissue deposition as well as soft-tissue formation inside the necrotic dental pulp in dogs (El-Tayeb et al., 2019; Abdelsalam et al., 2020; Abo EL Wafa et al., 2021; Mohamed et al., 2023).




Figure 2. Therapeutic actions of propolis (Sources: Machado et al., 2016; Torres et al., 2018; Anjum et al., 2019; Doğan et al., 2020; Özdemir et al., 2022).

Antibacterial action

Propolis has potent bacteriostatic and bactericidal effects since it inhibits protein production of bacterial growth by inhibiting cell division (Havsteen, 2002). In addition, some active ingredients of propolis induce disorganization of the cytoplasmic membrane and cell wall of the bacteria and cause partial bacteriolysis (Havsteen, 2002). Polyphenols, flavonoids, terpenes, pinocembrin, galangin, and ferulic acid are the common components of propolis responsible for the antibacterial action (Banskota et al., 2001; Özdemir et al., 2022). Propolis affects biofilm formation in its different diluents and has a marked antimicrobial activity of mastitis due to *Staphylococcus spp.* in goats (Dos Santos et al., 2019). In an *in vitro* study, propolis efficiently inhibits the *Pythium insidiosum* causing pythiosis in horses and dogs (Araújo et al., 2016). In addition, the antimicrobial activity of an alcoholic extract of Italian propolis, particularly on *Listeria monocytogenes*, could be used in ready-to-eat refrigerated dairy products, such as sterile skim milk, pasteurized cow's milk, as well as cow's and goat's whey cheese (Pedonese et al., 2019). Fouad et al. (2021) treated sheep experimentally infected with *Clostridium novyi* type B by administering 50 mg propolis extract/kg body weight daily for 15 days. In this study, the authors documented the antibacterial action of the propolis. Considering the inexpensive nature, widespread availability, natural source, and effectiveness of propolis as an antibacterial substance based on the *in vitro* investigations, it is imperative to conduct further *in vivo* studies to establish its antibacterial. Additionally, it is crucial to conduct experiments to evaluate the potentially toxic effects of various propolis extracts.

Antiviral action

Propolis has antiviral action due to its ability to inhibit the body's enzymes from removing the viral protein coating and changing the virus to an inactive form (Bogdanov, 2017). The antiviral activity of propolis is related to its content of polyphenols, flavonoids, caffeic acid, and quercetin (Farooqui and Farooqui, 2010; Shvarzbeyn and Huleihel, 2011; Özdemir et al., 2022). *In vitro* studies indicated that propolis has antiviral action against the herpes virus, rotavirus, pseudorabies virus, feline calicivirus, canine adenovirus type 2, bovine viral diarrhea virus, bovine respiratory syncytial virus and coronavirus strains (Cueto et al., 2011; Affonso et al., 2012; González-Búrquez et al., 2017; Gamil Zeedan and Abdalhamed, 2021). On the other hand, Tsuchiya et al. (2018) mention that Brazilian green propolis does not prevent acute equine encephalitis caused by herpes virus-9 despite its immune-stimulating action.

Antifungal action

Pinocembrin, galangin, benzoic acid, salicylic acid, and vanillin are the most common antifungal components of propolis (Özdemir et al., 2022). Propolis can inhibit *Malassezia pachydermatis*, *Dermatophytes*, *Trichophyton*, and *Microsporum* in dogs (Cardoso et al., 2010; Cruz Sánchez et al., 2014; Betancourt et al., 2015).

Antiparasitic action

The propolis contains compounds such as chrysin, quercetin, and galangin, which exhibit antiparasitic properties (Morsy et al., 2013; Dos Santos Araujo and Levistk, 2019; Linécio et al., 2022). Propolis reduces the intensity of infection with giardiasis and trypanosomiasis in experimentally infected mice (Abdel-Fattah and Nada, 2007) and treats rats infected with *Trypanosoma evansi* after oral administration of 100- 400 mg/kg daily for ten days (Gressler et al., 2012). Additionally, Fouad et al. (2021) successfully treated sheep experimentally infected with *Fasciola gigantica* by oral administration of 50 mg propolis extract/kg body weight daily for 15 days.

Anti-inflammatory action

Polyphenols, flavonoids, caffeic acid phenethyl ester, acacetin, etheric oils, and polyprenylated benzophenones are the main components of propolis responsible for its anti-inflammatory action (de Almeida and Menezes, 2002; Ramos and Miranda, 2007; Özdemir et al., 2022). Propolis inhibits myeloperoxidase activity, NADPH-oxidase ornithine decarboxylase, tyrosine-protein-kinase, and hyaluronidase from guinea pig mast cells (de Almeida and Menezes, 2002). Moreover, propolis inhibits the lipoxygenase pathway of arachidonic acid metabolism during inflammation (Mirzoeva and Calder, 1996). Future studies are necessary to obtain reliable conclusions on the effectiveness and underlying mechanisms through which propolis functions as an anti-inflammatory agent.

Immunomodulatory action

Polyphenols, flavonoids, and caffeic acid phenethyl ester are linked to the immunomodulatory action of propolis (Farooqui and Farooqui, 2010; Shvarzbeyn and Huleihel, 2011). Propolis has immune-boosting activity due to its antioxidant and broad-spectrum antibiotic actions. It inhibits the growth of fungi, viruses, bacteria, and other microbes. Moreover, propolis increases antibody production, activates B and T lymphocytes, enhances phagocytosis, stimulates the thymus gland, and thus boosts the thyroid gland to improve immunity (Boukraa, 2013; Gao et al., 2014). Oral use of alginate-propolis nanoparticles improves the immune status of Egyptian-Nubian newborn kids because it increases the immunoglobulins, IgG, and IgA and reduces the pro-inflammatory cytokines (Hegazi et al., 2021). Moreover, ethanolic extract of propolis has been used as a vaccine's adjuvant for some viral vaccines like canine parvovirus (CPV), and enhanced the production of antibodies against CPV in rats (El Ashry and Ahmad, 2012; Ferreira et al., 2012).

Antioxidant action

The antioxidant effect of propolis is related to its content's polyphenols, flavonoids, caffeic acid phenethyl ester, polyprenylated benzophenones, artemillin C, and Prenylated flavanones (Banskota et al., 2001; Farooqui and Farooqui, 2010). Propolis prevents tissue destruction from oxidative stress due to its ability to decrease the overproduction of superoxide anion and to restore the respiratory control ration in mitochondrial tissue. Propolis extract has a dose-dependent free radical scavenging action and inhibits xanthine oxidase activity (Banskota et al., 2000; Bogdanov, 2017, Özdemir et al., 2022).

Anti-tumoral action

Chrysin, quercetin, artemillin C and caffeic acid phenethyl ester are responsible for the anti-carcinogenic effect of propolis (Özdemir et al., 2022). Brazilian propolis is reported to have Artemillin C, which has a significant cytotoxic activity against transmissible venereal tumor cells from dogs. This cytotoxicity is due to DNA fragmentation and apoptosis induction (Bassani-Silva et al., 2007). Propolis can eliminate the canine osteosarcoma (OSA) cells obtained from naturally affected dogs (Costa Cinegaglia et al., 2013).

Wound healing

Propolis's antimicrobial, anti-inflammatory, and antioxidant activities enhance wound healing (Abu-Seida, 2015). Histologically, propolis-treated skin wounds exhibited moderate to complete thick vascular granulation tissue, more fibroblasts and collagen deposition, mild inflammatory cell infiltration, and complete epithelialization (Abu-Ahmed et al., 2013). Propolis also ameliorates the healing of burn scars in guinea pigs treated with daily topical application of 100 mg propolis extract/kg body weight (Elshater et al., 2017).

CLINICAL APPLICATIONS OF PROPOLIS IN LARGE AND SMALL ANIMAL PRACTICES

Due to its biological actions, Propolis has been used successfully in large and small animal practices. The clinical applications, dose, and pharmaceutical form of propolis are indicated in Table 1.

In veterinary medicine, propolis has been applied clinically to treat subclinical mastitis and mastitis (Fiordalisi et al., 2016; Šuran et al., 2020), lumpy skin disease (Farg et al., 2020), diarrhea and gynecological disorders in cattle (Gubicza and Molnar, 1987). It treated enzootic pneumonia in pigs and prevented gastrointestinal and respiratory diseases (Boukraa, 2013; Bogdanov, 2017). While in sheep, propolis treated the parasitic infestations such as

gastrointestinal nematodes and trematodes (Morsy et al., 2013; Kalil et al., 2019; Linécio et al., 2022), stimulated the immunity in Nubian breed kids (Hegazi et al., 2021), enhanced diet digestibility, and improved health status of ewes (Morsy et al., 2021). Otitis externa (Lozina et al., 2010; Betancourt et al., 2015), dermatophytosis (Cruz Sánchez et al., 2014), and Cushing's syndrome (Boukraa, 2013; Betancourt et al., 2015) could be treated with propolis in dogs. In equine practice, propolis could be used for local anesthesia (Bogdanov, 2017), enhancing wound healing (Abdel-Wahed et al., 2011; Abu-Ahmed et al., 2013) and treatment of chronic bronchitis as well as dermatomycosis (Flores Rodríguez et al., 2016; Zoja et al., 2019).

Table 1. Clinical applications, form, and dose of propolis in different animal species

Clinical application	Form/Dose of propolis	References
Cattle		
Mastitis	Propolis liniment	Fiordalisi et al. (2016)
Subclinical mastitis	- 1% intramammary propolis formulation - Propolis non-alcoholic extract given as intra-mammary infusion three times at 12 h intervals	Bacic et al. (2016) Šuran et al. (2020)
Gynecological diseases	Propolis candles	Bogdanov (2017)
Improvement of weight gain in calves	Feeding with 5 ml of 20% ethanol extract	Gubicza and Molnar (1987)
Anti-diarrhea in calves	- Feeding with 5 ml of 20% ethanol extract. - 4 mL/d of 30% red propolis ethanolic extract in whole milk	Gubicza and Molnar (1987) Slanzon et al. (2019)
Lumpy skin disease	Oral administration of Alginate-Propolis nanoparticles in a dose of 300 µl/ animal/ 3 days	Farag et al. (2020)
Foot and mouth disease	Propolis extract spray	Bogdanov (2017)
Pigs		
Enzootic pneumonia	Feeding with 0.5% propolis extract in milk	Bogdanov (2017)
Feeding of weak pigs	Feeding with 0.5% propolis extract in milk	Bogdanov (2017)
Prophylaxis of gastrointestinal diseases	Feeding with 0.5% propolis extract in milk	Bogdanov (2017)
Prophylaxis of respiratory diseases	Feeding with 0.5 % propolis extract in milk	Bogdanov (2017)
Sheep and goats		
Caseous lymphadenitis	Filling of the cavity with green propolis-based ointment once after surgery	Kalil et al. (2019)
Antiparasitic action and improvement of the health status of ewes during the flushing period	Oral administration of propolis ethanolic extract at a dose of 3 g/ewe/day for 21 days	Morsy et al. (2013)
Respiratory diseases	Oral administration of 0.06 ml Alginate-propolis nanoparticles (twice/week)	Hegazi et al. (2021)
Enhance diet digestibility, rumen microbial biosynthesis, mitigating methane formation and health status of late pregnant ewes	3 g red propolis extract/ ewe/ day	Morsy et al. (2021)
Dogs and cats		
Dermatophytosis	Weekly baths with a commercial soap made with propolis for 3 to 8 weeks and application of a topical ointment daily for three weeks	Cruz Sánchez et al. (2014)
Cushing's syndrome	Propolis in water at a dose of 0.4 - 0.5 g crude propolis per kg body weight every 12 hours for three months	Betancourt et al. (2015)
Otitis externa	Ear drops using 2.5% Propolis ethanolic extract in a mixture of glycerin-propylenglycol (1:1)/ twice daily for 14 days	Lozina et al. (2010)
Equine		
Dermatomycosis	A weekly bath with propolis-based shampoo and application of propolis-based ointment on the lesions 2 to 3 times a week for 4 weeks	Flores Rodríguez et al. (2016)
Chronic bronchitis	Ethanol extract of propolis and honey inhalation 3 times per day for 7 days	Zoja et al. (2019)
Local anesthetic	1-10% propolis extract preparation	Bogdanov (2017)
Wound healing	5% propolis in fish oil	Bogdanov (2017)
Wound healing	Dressing with propolis powder or propolis with honey proved	Abdel-Wahed et al. (2011)

Infectious diseases

Due to its antimicrobial, anti-inflammatory, and local antioxidant activities in the udder, propolis can control mastitis in cattle (Wang et al., 2016; Šuran et al., 2020). Intramammary infusion with 1% propolis formulation induced satisfactory antibacterial and antioxidant effects in dairy cows (Bacic et al., 2016; Šuran et al., 2020). Therefore, propolis may alternate the conventional antimicrobial drugs used to prevent and control subclinical mastitis in dairy cattle.

Kalil et al. (2019) used green propolis as a promising post-operative dressing agent for ovine caseous lymphadenitis. Propolis enhances wound healing and hair recovery, as well as inhibits wound infection. Inhalation with ethanolic extract propolis in combination with honey for 7 days showed antibacterial and antifungal actions, low heart and respiratory rates, and mucolytic action in the lower respiratory tract in horses with chronic bronchitis (Zoja et al., 2019). In addition, Farag et al. (2020) treated lumpy skin disease in cattle with oral alginate-propolis nanoparticles at a dose of 300 µl/ animal for three successive days and topical dressing and eye drops of propolis in some cases.

Dermatophytosis is a superficial skin infection caused by pathogenic dermatophytes, *Trichophyton*, and *Microsporum* (Betancourt et al., 2015). These pathogens have high zoonotic potential. Commercial propolis-based soap was used successfully to treat three dogs with dermatophytosis in 3-8 baths at one-week intervals together with using a topical propolis-based ointment for three weeks (Cruz Sánchez et al., 2014). Flores Rodríguez et al. (2016) treated dermatomycosis in horses with four baths at one-week intervals using propolis-based shampoo and topical dressing with propolis-based ointment two to three times a week for 28 days. *Staphylococcus aureus* is the commonly isolated bacteria from dogs with otitis externa (OE). Topical propolis ear drops treated the OE in dogs due to propolis's wide antimicrobial spectrum, anti-inflammatory, and antimycotic effects (Cardoso et al., 2010).

Dos Santos Araujo and Levistk (2019) found that propolis is efficient in controlling gastrointestinal helminths in sheep and recommended its use. However, it is still necessary to conduct more research on the ideal concentrations of propolis and its mode of action and residual effects. A single dose of 10 mL of 30% propolis alcoholic extract given orally demonstrated an antiparasitic effect in sheep and can be used in the control of endoparasites in sheep (Linécio et al., 2022).

Diarrhea

Red propolis supplementation improved health and reduced the incidence of diarrhea in calves and piglets (Bogdanov, 2017; Slanzon et al., 2019). This anti-diarrheal action could be attributed to the various biological actions of the propolis, such as antimicrobial, anti-inflammatory, and antioxidant actions.

Wound healing

From ancient times to recently, propolis was used to treat full-thickness skin wounds in horses and donkeys. Propolis treats wounds that exhibited better healing than wounds dressed with honey or saline (Abdel-Wahed et al., 2011; Abu-Ahmed et al., 2013).

Ocular diseases

Propolis has been used successfully in cats and dogs with different eye diseases. These diseases include blepharitis, infectious conjunctivitis, corneal edema, tear duct obstruction, keratoconjunctivitis sicca, corneal ulcers, and glaucoma. Propolis drops have been used for 5-7 days in acute cases and 10-15 days in chronic cases (Betancourt et al., 2015). Due to the numerous aforementioned medicinal actions, propolis is a promising eye therapeutic in animals.

Cushing's syndrome

A few studies indicated positive effects of propolis in treating Cushing's syndrome in dogs. In this regard, propolis was given at a dosage of 0.4-0.5 g/kg body weight twice daily for 3 months (Boukraa, 2013; Betancourt et al., 2015).

Reproductive performance

Prepartum using antibiotic growth promoters as feed additives in ruminant diets helped the transition from pregnancy to lactation and resulted in measurable health benefits (Morsy et al., 2016). Nevertheless, their use in animals is controversial due to the risk of residues transfer into meat and milk with increasing the development of resistant strains of bacteria (Mirzaei et al., 2022). Therefore, there is a continuous search for alternative natural feed additives to improve the reproductive performances of ruminants (de Aguiar et al., 2014). Propolis supplementation in ewes' diet increased the average daily gain and milk yield. However, it had no effect on lamb birth and weaning weights. The prepartum supplementation of propolis positively impacted the transition of ewes from pregnancy to lactation with good health of both ewes and lambs (Morsy et al., 2016). In contrast, propolis had no effect in maintaining sperm integrity and viability after thawing and was toxic to spermatozoa at concentrations of 0.25 and 0.5% (de Castilho et al., 2009). Moreover, the dietary addition of 3g of propolis/animal/day had no effect on bull mating performance or feed efficiency (Valero et al., 2016).

POTENTIAL BENEFITS OF PROPOLIS IN ANIMAL PRODUCTION

Antibiotics are effective feed supplements, particularly the ionophores (monensin, lasalocid, and salinomycin). However, these agents have recently faced reduced social acceptance in many countries due to the risk of residue in the milk and meat that develops resistant strains of bacteria (Morsy et al., 2013; Mirzaei et al., 2022). Therefore, several natural alternative feed additives have been tested, such as propolis.

Propolis has received increased attention in the last decade as a potential animal growth promoter due to its potent bacteriostatic and bactericidal properties (Righi et al., 2011). It had beneficial actions in animal production, such as increasing the average daily gain of lambs and milk conversion ratio in ewes (Morsy et al., 2016). Propolis also promoted the growth in calves, lambs, and piglets, improved the reproductive performance and increased the nutritional quality of the cows' milk (Gubicza and Molnar, 1987; Morsy et al., 2016; Cottica et al., 2019).

Propolis has better features than antibiotics as a feed additive due to its antimycotic and antiprotozoal actions, natural origin, availability, low cost, and safety for humans and animals (do Prado et al., 2010; da Silva et al., 2015). Considering these characteristics, it is suggested that a diet supplemented with propolis may inhibit the proteolytic bacteria with protein deamination, proteolysis, and production of gases. Consequently, propolis enhances the function of the gastrointestinal tract and feed digestibility (Morsy et al., 2013; da Silva et al., 2015).

Addition of propolis to the animal's diet induced more rapid muscular growth than animals fed a standard diet. Daily intake of propolis ethanol extract improved the daily weight gain in feedlot calves, lambs, and piglets and increased animals' productivity and meat quality in cattle (Ítavo et al., 2011; Bogdanov, 2017).

Moreover, adding 5g propolis/kg to the diet improved milk production, milk composition, and the antioxidants in Barki ewes. In addition, supplementation of propolis improved lambs' immune functions, growth performance, and antioxidant status in arid environments (Shedeed et al., 2019). Nevertheless, the addition of propolis extract into the sheep diet (40-50, and 60-50 forage/concentrate ratio) had no effects on their feed intake hematological, biochemical, immunological, and nutrient digestibility features (Prado-Calixto et al., 2017; da Silva et al., 2018). Moreover, adding propolis to the lambs' diet did not influence carcass characteristics (Ítavo et al., 2009).

Soybean oil interacts with ethanolic extract of propolis when added to the diet of dairy goats; therefore, soybean oil decreases the intake of dry matter, organic matter, and neutral detergent fiber only in the presence of propolis and increases the intake of crude protein in the absence of propolis (de Paula Lana et al., 2005).

Using ethanolic propolis extract as a feed additive for dairy cows increased milk protection against lipid oxidation responsible for a rancid smell (Cottica et al., 2019). Therefore, the antioxidant capacity of the milk increased, and consequently, propolis improved the milk quality when added to the diet of dairy cows from this aspect (Cottica et al., 2019). However, adding propolis to the diets of dairy cows had no effects on dry matter intake, milk production, feed conversion efficiency, milk solid concentrations, or somatic cell score (Aguilar et al., 2014).

On the other hand, the addition of propolis to the forage-based diet had a negative effect on the concentration and intake of digestible energy of roughage-based diets for growing steers (do Prado et al., 2010). In addition, adding propolis extract to the diet of water buffaloes reduced the population of ciliate protozoa in the rumen (Ríspoli et al., 2009). Moreover, the addition of propolis to the diet of feedlot bulls had no effect on microbial synthesis and carcass characteristics. These characteristics include conformation, carcass length, leg length, cushion thickness, *Longissimus* muscle area, *Longissimus* muscle area/100 kg of live weight, fat thickness, color, texture, and marbling (Zawadzki et al., 2011; Valero et al., 2014; 2015). Propolis did not significantly affect weanling pigs' live weight gain, feed consumption, or feed conversion ratio (Dierckx and Funari, 1999).

According to the available literature, the performed clinical studies on the efficacy of propolis as a feed additive reveal controversial results. Therefore, future *in vivo* studies on this topic are highly recommended.

CONCLUSION

Propolis is a promising natural agent that can alternate conventional pharmaceuticals, particularly antibiotics. It improves animal health and production with no adverse effects and at a low cost. Propolis can prevent and treat several animal diseases like mastitis, lumpy skin disease, foot and mouth disease, reproductive disorders, and diarrhea in cattle. Furthermore, propolis could improve weight gain in cattle. In equine, propolis has been used as a local anesthetic for treating dermatomycosis, chronic bronchitis, and skin wounds. In pigs, propolis has been used to treat enzootic pneumonia, feeding weak pigs, and acts as a prophylaxis for gastrointestinal and respiratory diseases. Regarding sheep and goats, propolis has been applied to treat caseous lymphadenitis and parasitic diseases and improve the immune status of kids and the health status of late pregnant ewes. In dogs and cats, propolis has been applied to treat otitis externa, eye diseases, Cushing's syndrome, and dermatophytosis. In addition, propolis is used to treat transmissible venereal tumor in dogs. Moreover, propolis positively affects animal production, by increasing the average daily gain and milk yield in sheep, the growth of calves, lambs, and piglets, the reproductive performance, and cow's milk nutritional quality. However, further extensive clinical studies are recommended to declare the usefulness of propolis in veterinary medicine and to obtain reliable conclusions on its potential benefits in animal health and production.

DECLARATIONS

Availability of data and materials

All data are presented in the published manuscript.

Funding

This manuscript received no financial assistance from any agency.

Author's contributions

Ashraf M. Abu-Seida collected and analyzed the data as well as wrote and revised the manuscript.

Conflict of interests

The author declares no conflicts of interest.

Ethical consideration

The author checked plagiarism, misconduct, data fabrication and/or falsification.

REFERENCES

- Abdel-Fattah NS and Nada OH (2007). Effect of propolis versus metronidazole and their combined use in treatment of acute experimental Giardiasis. *Journal of Egyptian Society of Parasitology*, 37(Supple 2): 691-710. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17926808>
- Abdelsalam N, Abu-Seida AM, Fayyad D, and Tawfik H (2020). Radiographic and histopathologic outcomes of immature dog teeth with apical periodontitis after revascularization using propolis. An *in vivo* study. *Saudi Endodontic Journal*, 10(3): 199-207. DOI: http://www.doi.org/10.4103/sej.sej_178_19
- Abdulkhani A, Hosseinzadeh J, Ashori A, and Esmaeeli H (2017). Evaluation of the antibacterial activity of cellulose nanofibers/polylactic acid composites coated with ethanolic extract of propolis. *Polymer Composites*, 38(1): 13-19. DOI: <http://www.doi.org/10.1002/pc.23554>
- Abdel-Wahed RE, El-Kammar MH, Abu-Ahmed H, and Abo-Bakr E (2011). Propolis (bee glue). A new dressing for equine wounds. *Alexandria Journal of Veterinary Sciences*, 34(1): 21-36. Available at: <https://www.cabdirect.org/cabdirect/abstract/20133120860>
- Abo El-Mal EO, Abu-Seida AM, and El Ashry SH (2021). Biological evaluation of hesperidin for direct pulp capping in dogs' teeth. *International Journal of Experimental Pathology*, 102(1): 32-44. <http://www.doi.org/10.1111/iep.12385>
- Abo EL Wafa MR, Niazy MA, Abo Hagar EA, and Abu-Seida AM (2021). Biological pulp response of pulpine, polyamidoamine dendrimer and their combination in dogs and their remineralizing effect on carious affected human dentin: A randomized clinical trial. *Al-Azhar Journal of Dentistry*, 8(4): 8. DOI: <https://www.doi.org/10.21608/ADJG.2021.50064.1321>
- Abu-Ahmed H, Abdel-Wahed RE, El-Kammar MH, and El-Neweshy MS (2013). Evaluation of the effectiveness of propolis compared with honey on second intention wound healing in the equine. *Middle East Journal of Scientific Research*, 14(10): 1292-1298. DOI: <http://www.idosi.org/mejsr/mejsr14%2810%2913/6.pdf>
- Abu-Seida AM (2015). Effect of propolis on experimental cutaneous wound healing in dogs. *Veterinary Medicine International*, 2015: 672643. DOI: <https://www.doi.org/10.1155/2015/672643>
- Affonso IB, Gatti SP, de Medeiros ASR, Buzinaro MG, and Samara SI (2012). Bovine respiratory syncytial virus (BRSV) activity in dairy calves. *Ciência Animal Brasileira*, 13(2): 253-258. Available at: <https://www.cabdirect.org/cabdirect/abstract/20123237545>
- Aguiar SC, Coticab SM, Boeingb JS, Samensaria RP, Santosa GT, Visentainerb JV, and Zeoulaa LM (2014). Effect of feeding phenolic compounds from propolis extracts to dairy cows on milk production, milk fatty acid composition, and the antioxidant capacity of milk. *Animal Feed Science and Technology*, 193: 148-154. DOI: <https://www.doi.org/10.1016/j.anifeedsci.2014.04.006>
- Alp H (2018). Effects of propolis on immune system. *ANADOLU Ege Tarımsal Araştırma Enstitüsü Dergisi*, 28(2): 99-104. Available at: <https://dergipark.org.tr/en/pub/anadolu/issue/41816/504495>
- Anjum SI, Ullah A, Khan KA, Attaullah M, Khan H, Ali H, Bashir MA, Tahir M, Ansari MJ, Ghramh HA et al. (2019). Composition and functional properties of propolis (bee glue): A review. *Saudi Journal of Biological Sciences*, 26(7): 1695-1703. DOI: <https://www.doi.org/10.1016/j.sjbs.2018.08.013>
- Araújo MJAM, Bosco SDMG, and Sforcin JM (2016). *Pythium insidiosum*: Inhibitory effects of propolis and geopropolis on hyphal growth. *Brazilian Journal of Microbiology*, 47(4): 863-869. DOI: <https://www.doi.org/10.1016/j.bjm.2016.06.008>
- Bacic G, Macesic N, Radin L, Aladrovic J, Matanovic K, Masek T, Brozic D, Benic M, Radic B, Bacic I et al. (2016). Intramammary propolis formulation for subclinical mastitis prevention and treatment in dairy cows. *Journal of Dairy, Veterinary & Animal Research*, 3(5): 159. DOI: <https://www.doi.org/10.15406/jdvar.2016.03.00091>
- Bankova V, Popova M, and Trusheva B (2014). Propolis volatile compounds: Chemical diversity and biological activity: A review. *Chemistry Central Journal*, 8: 28. DOI: <https://www.doi.org/10.1186/1752-153X-8-28>
- Banskota AH, Tezuka Y, Adnyana IK, Midorikawa K, Matsushige K, Message D, Huertas AAG, and Kadota S (2000). Cytotoxic, hepatoprotective and free radical scavenging effects of propolis from Brazil, Peru, the Netherlands and China. *Journal of Ethnopharmacology*, 72(1-2): 239-246. DOI: [https://www.doi.org/10.1016/S0378-8741\(00\)00252-X](https://www.doi.org/10.1016/S0378-8741(00)00252-X)
- Banskota AH, Tezuka Y, and Kadota S (2001). Recent progress in pharmacological research of propolis. *Phytotherapy Research*, 15(7): 561-571. DOI: <https://www.doi.org/10.1002/ptr.1029>
- Bassani-Silva S, Sforcin JM, Amaral AS, Gazpar Luiz FJ, and Rocha Noeme S (2007). Propolis effect *in vitro* on canine transmissible venereal tumor cells. *Revista Portuguesa Ciências Veterinárias*, 102(563-564): 261-265. Available at: <https://spcv.pt/wp-content/uploads/2023/06/Vol102-n563-564.pdf>
- Betancourt NT, García-Contreras L, and Cruz Sánchez TA (2015). Propolis in dogs: Clinical experiences and perspectives (A brief review). *Open Journal of Veterinary Medicine*, 5(1): 53251. DOI: <https://www.doi.org/10.4236/ojvm.2015.51002>
- Bogdanov S (2017). Propolis: Biological properties and medical applications. *The propolis book*, Chapter 2, pp. 1-41. <https://s66a5b07cb261a6cc.jimcontent.com/download/version/1609255034/module/11112716973/name/PropolisBook2.pdf>

- Bonamigo T, Campos JF, Alfredo TM, Balestieri JBP, Cardoso CAL, Paredes-Gamero EJ, Picoli Souza K, and Dos Santos EL (2017). Antioxidant cytotoxic, and toxic activities of propolis from two native bees in Brazil: *Scaptotrigona depilis* and *Melipona quadrifasciata anthidioides*. *Oxidative Medicine and Cellular Longevity*, 2017: 1038153. DOI: <https://www.doi.org/10.1155/2017/1038153>
- Boukraa L (2013). Honey and bee products for animal health and wellness. Honey in traditional and modern medicine, 1st Edition. CRC Press., Boca Raton, Chapter 13, pp. 263-278. DOI: <https://www.doi.org/10.1201/b15608>
- Cardoso RL, Maboni F, Machado G, Hartz AS, and Castagna de Vargas A (2010). Antimicrobial activity of propolis extract against *Staphylococcus coagulase positive* and *Malassezia pachydermatis* of canine otitis. *Veterinary Microbiology*, 142(3-4): 432-434. DOI: <https://www.doi.org/10.1016/j.vetmic.2009.09.070>
- Costa Cinegaglia N, Bersano PRO, Mendes Araújo MJA, Búfalo MC, and Sforcin JM (2013). Anticancer effects of geopropolis produced by sting less bees on canine osteosarcoma cells *in vitro*. *Evidence-Based Complementary and Alternative Medicine*, 2013: 737386. DOI: <https://www.doi.org/10.1155/2013/737386>
- Cottica SM, Amado DAV, de Aguiar SC, Boeing JS, Franco SL, Zeoula LM, and Visentainer JV (2019). Antioxidant activity and lipid oxidation in milk from cows with soybean oil and propolis extract added to their feed. *Food Science and Technology*, 39(2): 467-474. DOI: <https://www.doi.org/10.1590/fst.33817>
- Cruz Sánchez TA, Estrada García PA, López Zamora CI, Martínez MA, Valencia VP, and Orozco AL (2014). Use of propolis for topical treatment of dermatophytosis in dog. *Open Journal of Veterinary Medicine*, 4(10): 239-245. DOI: <https://www.doi.org/10.4236/ojvm.2014.410028>
- Cueto AP, Alves SH, Pilau M, Weiblen R, Kubiça TF, and Lovato LT (2011). Antiviral activity of propolis extracts against feline calicivirus, canine adenovirus 2, and bovine viral diarrhoea virus. *Ciencia Rural*, 41(10): 1800-1806. DOI: <https://www.doi.org/10.1590/S0103-84782011001000021>
- da Silva FGB, Yamamoto SM, da Silva EMS, Queiroz MAA, Gordiano LA, and Formiga MA (2015). Propolis extract and sodium monensin on ruminal fermentation and hematological parameters in sheep. *Acta Scientiarum Animal Sciences*, 37(3): 273-280. DOI: <https://www.doi.org/10.4025/actascianimsci.v37i3.25725>
- da Silva LM, de Souza P, Al Jaouni SK, Harakeh S, Golbabapour S, and de Andrade SF (2018). Propolis and its potential to treat gastrointestinal disorders. *Evidence- Based Complementary and Alternative Medicine*, 2018: 2035820. DOI: <https://www.doi.org/10.1155/2018/2035820>
- de Aguiar SC, de Paula EM, Yoshimura EH, dos Santos WBR, Machado E, Valero MV, dos Santos GT, and Zeoula LM (2014). Effects of phenolic compounds in propolis on digestive and ruminal parameters in dairy cows. *Revista Brasileira de Zootecnia*, 43(4): 197-206. DOI: <https://www.doi.org/10.1590/S1516-35982014000400006>
- de Almeida EC and Menezes H (2002). Anti-inflammatory activity of propolis extracts: A review. *Journal of Venomous Animals and Toxins*, 8(2): 191-212. DOI: <https://www.doi.org/10.1590/S0104-79302002000200002>
- de Castilho EF, Guimarães JD, Martins LF, Pinho RO, Guimarães SEF, and Espescht CJB (2009). Use of propolis and ascorbic acid on goat semen cryopreservation. *Revista Brasileira de Zootecnia*, 38(12): 2335-2345. DOI: <https://www.doi.org/10.1590/S1516-35982009001200007>
- de Mendonça LS, de Mendonça FMR, de Araújo YLFM, de Araújo ED, Ramalho SA, Narain N, Jain S, Orellana SC, Padilha FF, and Cardoso JC (2015). Chemical markers and antifungal activity of red propolis from Sergipe. *Food Science and Technology*, 35(2): 291-298. DOI: <https://www.doi.org/10.1590/1678-457X.6554>
- de Paula Lana R, Camardelli MML, de Queiroz AC, Rodrigues MT, Eifert EDC, Miranda EN, and Almeida ICC (2005). Soybean oil and propolis in the diets of dairy goats. *Revista Brasileira de Zootecnia*, 34(2): 650-658. DOI: <https://www.doi.org/10.1590/S1516-35982005000200035>
- Dierckx SM and Funari SRC (1999). Use of propolis in the feeding of weanling pigs as an additive and for prevention of diarrhoea. *Latin American Archives of Animal Production*, 7(2): 109-116. Available at: https://ojs.alpa.uy/index.php/ojs_files/article/view/518
- Doğan H, Silici S, and Ozcimen AA (2020). Biological effects of propolis on cancer. *Turkish Journal of Agriculture- Food Science and Technology*, 8(3): 573-579. DOI: <https://www.doi.org/10.24925/turjaf.v8i3.573-579.2939>
- do Prado OPO, Zeoula LM, de Moura LP, Franco LS, do Prado IN, and Gomes HC (2010). Digestibility and ruminal parameters of diet based on forage with the addition of propolis and sodium monensin for steers. *Revista Brasileira de Zootecnia*, 39(6): 1336-1345. DOI: <https://www.doi.org/10.1590/S1516-35982010000600024>
- Dos Santos HC, Vieira DS, Yamamoto SM, Costa MM, Sá MCA, Silva EMS, and Silva TMS (2019). Antimicrobial activity of propolis extract fractions against *Staphylococcus spp.* isolated from goat mastitis. *Pesquisa Veterinária Brasileira*, 39(12): 954-960. DOI: <https://www.doi.org/10.1590/1678-5150-PVB-5940>
- Dos Santos Araujo J and Levistk IC (2019). Propolis in the control of helminths in sheep. *International Journal of Agronomy and Agricultural Research*, 14(6): 69-74. Available at: <https://www.innspub.net/wp-content/uploads/2022/04/IJAAR-V14-No6-p69-74.pdf>
- El Ashry ESH and Ahmad TA (2012). Review the use of Propolis as vaccine's adjuvant. *Vaccine*, 31(1): 31-39. DOI: <http://www.doi.org/10.1016/j.vaccine.2012.10.095>
- Elkhenany H, El-Badri N, and Dhar M (2019). Green propolis extract promotes *in vitro* proliferation, differentiation, and migration of bone marrow stromal cells. *Biomedicine & Pharmacotherapy*, 115: 108861. DOI: <https://www.doi.org/10.1016/j.biopha.2019.108861>
- Elshater AA, Salman MMA, and Abd-Elmegeed SAA (2017). Ameliorative effects of propolis extract on some biochemical and hematological parameters of burnt skin of male guinea pigs. *Egyptian Academic Journal of Biological Sciences. C, Physiology and Molecular Biology*, 9(1): 47-57. DOI: <https://www.doi.org/10.21608/EAJBSC.2017.13672>
- El-Tayeb MM, Abu-Seida AM, El Ashry SH, and El-Hady SA (2019). Evaluation of antibacterial activity of propolis on regenerative potential of necrotic immature permanent teeth in dogs. *BMC Oral Health*, 19: 174. DOI: <https://www.doi.org/10.1186/s12903-019-0835-0>
- Farag TK, El-Houssiny AS, Abdel-Rahman EH, and Hegazi AG (2020). A new approach to the treatment of lumpy skin disease infection in cattle by using propolis encapsulated within ALG NPs. *Advances in Animal & Veterinary Sciences*, 8(12): 1346-1355. Available at: http://researcherslinks.com/nexus/uploads/files/AAVS_8_12_1346-1355.pdf
- Farooqui T and Farooqui AA (2010). Molecular mechanism underlying the therapeutic activities of propolis: A critical review. *Current Nutrition & Food Science*, 6(3): 188-199. DOI: <https://www.doi.org/10.2174/157340110792389136>
- Ferreira LN, Finger FP, Caetano de Castro C, Siedler BS, Munhoz LS, Vargas GDA, Fischer G, and Hubner SO (2012). Effect of the etanolic extract from green propolis on production of antibodies after immunization against Canine Parvovirus (CPV) and Canine Coronavirus (CCoV). *Brazilian Journal of Veterinary Research and Animal Science*, 49(2): 116-122. Available at: <https://pesquisa.bvsalud.org/portal/resource/pt/lil-687583>
- Fiordalisi SAL, Honorato LA, Loiko MR, Avancini CAM, Veleirinho MBR, Filho LCPM, and Kuhnen S (2016). The effects of Brazilian propolis on etiological agents of mastitis and the viability of bovine mammary gland explants. *Journal of Dairy Science*, 99(3): 2308-2318. DOI: <https://www.doi.org/10.3168/jds.2015-9777>
- Flores Rodríguez IS, Monteagudo MM, Orozco AL, and Sánchez TAC (2016). Use of Mexican propolis for the topical treatment of dermatomycosis in horses. *Open Journal of Veterinary Medicine*, 6(1): 62770. DOI: <https://www.doi.org/10.4236/ojvm.2016.61001>

- Fouad EA, Toaleb NI, Hassan SE, El Shanawany EE, Keshta HG, Abdel-Rahman EH, and Hegazi AG (2021). Evaluation of the therapeutic effect of propolis on *Fasciola gigantica* and *Clostridium novyi* type B infections in sheep. *Tropical Biomedicine*, 38(2): 102-110. DOI: <https://www.doi.org/10.47665/tb.38.2.041>
- Gamil Zeedan GS and Abdalhamed AM (2021). Antiviral effects of plant extracts used in the treatment of important animal viral diseases. *World's Veterinary Journal*, 11(4): 521-533. DOI: <https://www.doi.org/10.54203/scil.2021.wvj67>
- Gao W, Wu J, Wei J, Pu L, Guo C, Yang J, Yang M, and Luo H (2014). Brazilian green propolis improves immune function in aged mice. *Journal of Clinical Biochemistry and Nutrition*, 55(1): 7-10. DOI: <https://www.doi.org/10.3164/jcfn.13-70>
- González-Búrquez MD, Cruz-Sánchez TA, Soto-zárate CI, Carrillo-Miranda L, and Fonseca-Coronado S (2017). *In vitro* activity of propolis on domestic animal viruses: A review. *Interciencia*, 42(5): 272-276. Available at: <https://www.redalyc.org/journal/339/33952810002/html/>
- Gressler LT, Da Silva AS, Machado G, Dalla Rosa L, Dorneles F, Gressler, TL, Oliveira MS, Zanette RA, Agueda de Vargas CP, and Monteiro SG (2012). Susceptibility of *Trypanosoma evansi* to propolis extract *in vitro* and in experimentally infected rats. *Research Veterinary Science*, 93(3): 1314-1317. DOI: <http://www.doi.org/10.1016/j.rvsc.2012.02.007>
- Gubicza A and Molnar P (1987). Propolis in the rearing of calves. *Magyar Mezogazd*, 42(17): 14-30.
- Havsteen BH (2002). The biochemistry and medical significance of the flavonoids. *Pharmacology & Therapeutics*, 96(2-3): 67-202. DOI: [https://www.doi.org/10.1016/s0163-7258\(02\)00298-x](https://www.doi.org/10.1016/s0163-7258(02)00298-x)
- Hegazi AG, El-Houssiny AS, Sadek WMA, Al-Guthami FM, Al-Gethami AFM, Sadik AMA, and Farag TK (2021). Egyptian propolis 13: Influence of propolis and alginate-propolis NPs on Egyptian Nubian goats serum immunoglobulins and cytokines level. *Advances in Animal and Veterinary Sciences*, 9(2): 280-288. DOI: <https://www.doi.org/10.17582/journal.aavs/2021/9.2.280.288>
- Ítavo CCBF, Morais MG, Costa C, Ítavo VCL, de Macedo FAF, and Tomich TR (2009). Carcass characteristics, non-components and yield of retail products from lambs in feedlot system receiving diets with propolis or sodic monensin as additive. *Revista Brasileira de Zootecnia*, 38(5): 898-905. DOI: <https://www.doi.org/10.1590/S1516-35982009000500017>
- Ítavo CCBF, Morais MG, Costa C, Ítavo LCV, Franco GL, da Silva JA, and Reis FA (2011). Addition of propolis or monensin in the diet: Behavior and productivity of lambs in feedlot. *Animal Feed Science and Technology*, 165(3-4): 161-166. DOI: <https://www.doi.org/10.1016/j.anifeeds.2011.02.020>
- Kalil MA, Santos LM, Barral TD, Rodrigues DM, Pereira NP, Aquino Sá MC, Umsza-Guez MA, Machado BAS, Meyer R, and Portel RW (2019). Brazilian green propolis as a therapeutic agent for the post-surgical treatment of caseous lymphadenitis in sheep. *Frontiers in Veterinary Science*, 6: 399. DOI: <https://www.doi.org/10.3389/fvets.2019.00399>
- Linécio M, Garcia RC, Galhardo D, de Oliveira TE, and de Mello Peixoto ECT (2022). Alcoholic extract of propolis for worms' control in sheep. *Research Society Development*, 11(1): e1711120617. DOI: <https://www.doi.org/10.33448/rsd-v11i1.20617>
- Lozina LA, Peichoto ME, Boehringer SI, Koscienczuk P, Granero GE, and Acosta OC (2010). Efficacy of Argentine propolis formulation for topical treatment of canine otitis extern. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 62(6): 1359-1366. DOI: <https://www.doi.org/10.1590/S0102-09352010000600010>
- Machado BS, Pulcino TN, Silva AL, Tadeu Melo D, Gomes Silva R, and Mendonça IG (2016). Propolis as an alternative in prevention and control of dental cavity. *Journal of Apitherapeutics*, 1(2): 47-50. DOI: <https://www.doi.org/10.5455/ja.20160726115117>
- Marcucci MC (1995). Propolis: Chemical composition, biological properties and therapeutic activity. *Apidologie*, 26(2): 83-99. DOI: <https://www.doi.org/10.1051/apido:19950202>
- Mirzaei A, Razavi SA, Babazadeh D, Laven R, and Saeed M (2022). Roles of probiotics in farm animals: A review. *Farm Animal Health and Nutrition*, 1(1): 17-25. DOI: <https://www.doi.org/10.58803/fahn.v1i1.8>
- Mirzoeva OK and Calder PC (1996). The effect of propolis and its components on eicosanoid production during the inflammatory response. *Prostaglandins, Leukotrienes, and Essential Fatty Acids*, 55(6): 441-449. DOI: [https://www.doi.org/10.1016/s0952-3278\(96\)90129-5](https://www.doi.org/10.1016/s0952-3278(96)90129-5)
- Mohamed M, Hashem AA, Obeid MF, Abu-Seida AM (2023). Histopathological and immunohistochemical profiles of pulp tissues in immature dogs' teeth to two recently introduced pulpotomy materials. *Clinical Oral Investigations*, 27: 3095-3103. DOI: <https://www.doi.org/10.1007/s00784-023-04915-5>
- Morsy AS, Abdalla AL, Soltan YA, Sallam SMA, El-Azrak KEM, Louvandini H, and Alencar SM (2013). Effect of Brazilian red propolis administration on hematological, biochemical variables and parasitic response of Santa Inês ewes during and after flushing period. *Tropical Animal Health and Production*, 45: 1609-1618. DOI: <https://www.doi.org/10.1007/s11250-013-0406-3>
- Morsy AS, Soltan YA, Sallam SMA, Alencar SM, and Abdalla AL (2016). Impact of Brazilian red propolis extract on blood metabolites, milk production, and lamb performance of Santa Inês ewes. *Tropical Animal Health and Production*, 48: 1043-1050. DOI: <https://www.doi.org/10.1007/s11250-016-1054-1>
- Morsy AS, SoltanYA, El-Zaiat HM, Alencarc SM, and Abdallad AL (2021). Bee propolis extract as a phytogetic feed additive to enhance diet digestibility, rumen microbial biosynthesis, mitigating methane formation and health status of late pregnant ewes. *Animal Feed Science and Technology*, 273: 114834. DOI: <https://www.doi.org/10.1016/j.anifeeds.2021.114834>
- Özdemir V, Yanar M, and Koçyiğit R (2022). General properties of propolis and its usage in ruminants. *Journal of the Hellenic Veterinary Medicine Society*, 73(2): 3905-3912. DOI: <https://www.doi.org/10.12681/jhvms.26334>
- Pedonese F, Verani G, Torracca B, Turchi B, Felicioli A, and Nuvoloni R (2019). Effect of an Italian propolis on the growth of *Listeria monocytogenes*, *Staphylococcus aureus* and *Bacillus cereus* in milk and whey cheese. *Italian Journal of Food Safety*, 8(4): 8036. DOI: <https://www.doi.org/10.4081/ijfs.2019.8036>
- Prado-Calixto OP, Mizubuti IY, Ribeiro ELA, Pereira ES, Silva RT, Corletto NL, Peixoto ELT, Carvalho N, Nihei AK, Massaro Júnior FL et al. (2017). Ingestive behavior and blood profile in sheep fed with diets containing powdered propolis extract additives. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 69(2): 381-390. DOI: <https://www.doi.org/10.1590/1678-4162-9065>
- Przybyłek I and Karpiński TM (2019). Antibacterial properties of propolis. *Molecules*, 24(11): 2047. DOI: <https://www.doi.org/10.3390/molecules24112047>
- Ramos AFN and Miranda JL (2007). Propolis: A review of its anti-inflammatory and healing actions. *Journal of Venomous Animals and Toxins Including Tropical Diseases*, 13(4): 697-710. DOI: <https://www.doi.org/10.1590/S1678-91992007000400002>
- Righi AA, Alves TR, Negri G, Marques LM, Breyer H, and Salatino A (2011). Brazilian red propolis: Unreported substances, antioxidant and antimicrobial activities. *Journal of the Science of Food and Agriculture*, 91(13): 2363-2370. DOI: <https://www.doi.org/10.1002/jsfa.4468>
- Ríspoli TB, Rodrigues IL, Neto RGM, Kazama R, Prado OPP, Zeoula LM, and Arcuri PB (2009). Ruminal ciliate protozoa of cattle and buffalo fed on diet supplemented with monensin or extracts from propolis. *Pesquisa Agropecuária Brasileira*, 44(1): 92-97. DOI: <https://www.doi.org/10.1590/S0100-204X2009000100013>

- Saleh RS, Nagi SM, Khallaf ME, Abd El-Alim SH, Zaazou MH, Abu- Seida AM, and Ibrahim MN (2016). *In-Vivo* assessment of dentin bridge formation after using MTA and experimental propolis paste as direct pulp capping material. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7(4): 1244-1250. Available at: [https://www.rjpbcs.com/pdf/2016_7\(4\)/1581.pdf](https://www.rjpbcs.com/pdf/2016_7(4)/1581.pdf)
- Shedeed HA, Farrag B, Elwakeel EA, Abd El-Hamid IS, and El-Rayes MA (2019). Propolis supplementation improved productivity, oxidative status, and immune response of Barki ewes and lambs. *Veterinary World*, 12(6): 834-843. DOI: <https://www.doi.org/10.14202/vetworld.2019.834-843>
- Shvarzbejn J and Huleihel M (2011). Effect of propolis and caffeic acid phenethyl ester (CAPE) on NFκB activation by HTLV-1 Tax. *Antivirus Research*, 90(3): 108-115. DOI: <https://www.doi.org/10.1016/j.antiviral.2011.03.177>
- Slanzon GS, Toledo AF, Silva AB, Coelho MG, da Silva MD, Cezar AM, and Bittar CM (2019). Red propolis as an additive for preweaned dairy calves: Effect on growth performance, health, and selected blood parameters. *Journal of Dairy Science*, 102(10): 8952-8962. DOI: <https://www.doi.org/10.3168/jds.2019-16646>
- Šuran J, Aladrović J, Ljubić BB, Vlamić J, Mamić M, Radić B, Bačić G, Mačešić N, Benić M, Kostelić A et al. (2020). The antioxidant effect of the novel bee-product based intramammary formulation Apimast® in dairy cattle. *Veterinarski Arhiv*, 90(3): 225-233. DOI: <https://www.doi.org/10.24099/vet.arhiv.0865>
- Torres AR, Sandjo L, Friedemann MT, Tomazzoli MM, Maraschin M, Mello CF, and Santos ARS (2018). Chemical characterization, antioxidant and antimicrobial activity of propolis obtained from *Melipona quadrifasciata* and *Tetragonisca angustula* stingless bees. *Brazilian Journal of Medical and Biological Research*, 51(6): e7118. DOI: <https://www.doi.org/10.1590/1414-431x20187118>
- Tsuchiya Y, Abd-Ellatieff HA, Abou Rawash AA, and Yanai T (2018). Evaluation of preventive effects of Brazilian green propolis on equine Herpes virus 9-induced acute encephalitis. *Journal of Veterinary Science and Medical Diagnosis*, 7(2): 1-5. DOI: <https://www.doi.org/10.4172/2325-9590.1000252>
- Valero MV, Torrecilhas JA, Zawadzki F, Bonafé EG, Madrona GS, do Prado RM, Passetti RAC, Rivaroli DC, Visentainer JV, and do PradoVN (2014). Propolis or cashew and castor oils effects on composition of *Longissimus* muscle of crossbred bulls finished in feedlot. *Chilean Journal of Agricultural Research*, 74(4): 445-451. DOI: <https://www.doi.org/10.4067/S0718-58392014000400011>
- Valero MV, Zeoula LM, Moura LPP, Júnior JBGC, Sestari BB, and Prado IN (2015). Propolis extract in the diet of crossbred (½ Angus vs. ½ Nelore) bulls finished in feedlot: Animal performance, feed efficiency and carcass characteristics. *Semina: Ciências Agrárias Londrina*, 36(2): 1067-1078. DOI: <https://www.doi.org/10.5433/1679-0359.2015v36n2p1067>
- Valero MV, Farias MS, Zawadzki F, Prado RM, Fugita CA, Rivaroli DC, Ornaghi MG, and do Prado IN (2016). Feeding propolis or essential oils (cashew and castor) to bulls: Performance, digestibility, and blood cell counts. *Revista Colombiana de Ciencias Pecuarias*, 29(1): 33-42. DOI: <https://www.doi.org/10.17533/udea.rccp.v29n1a04>
- Wang L, Jin XL, Shen XG, Sun LP, Wu LM, Wei JQ, Marcucci MC, Hu FL, and Liu JX (2016). Effects of Chinese propolis in protecting bovine mammary epithelial cells against mastitis pathogens-induced cell damage. *Mediators of Inflammation*, 2016: 8028291. DOI: <https://www.doi.org/10.1155/2016/8028291>
- Zawadzki F, Prado IN, Marques JA, Zeoula LM, Rotta PP, Sestari BB, Valero MV, and Rivaroli DC (2011). Sodium monensin or propolis extract in the diets of feedlot-finished bulls: effects on animal performance and carcass characteristics. *Journal of Animal Feed Science*, 20(1): 16-25. DOI: <https://www.doi.org/10.22358/jafs/66153/2011>
- Zoja M, Sonata S, Rita S, Elvina S, and Karolis S (2019). Influence of the inhalation with ethanol extract of propolis and honey on physiological parameters and inflammation in horses with chronic bronchitis. 16th World Equine Veterinary Association Congress, Verona, Italy, pp. 357. Available at: <https://ismu.lt/cris/handle/20.500.12512/22313>

Publisher's note: [Scienceline Publication](https://www.scienceopen.com/) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023