



# Common Infectious and Parasitic Diseases in Goats of Tropical Africa and their Impacts on Production Performance: A Review

Kétomon Pierre Challaton<sup>1\*</sup>, Kadoéito Cyrille Boko<sup>2</sup>, Coovi Guérolé Akouedegni<sup>1</sup>, Goué Géorcelin Alowanou<sup>1,3</sup>, Aboudou Habirou Kifouly<sup>1</sup>, and Mawulé Sylvie Hounzangbé-Adoté<sup>1</sup>

<sup>1</sup>Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, Cotonou, Benin

<sup>2</sup>Communicable Diseases Research Unit, Applied Biology Research Laboratory, Polytechnic School of Abomey-Calavi, University of Abomey-Calavi, Cotonou, Benin

<sup>3</sup>High School of Technical Education, National University of Sciences, Technologies, Engineering and Mathematics, Abomey, Benin

\*Corresponding author's Email: [pierrechallaton00@gmail.com](mailto:pierrechallaton00@gmail.com)

## ABSTRACT

Available scientific studies on goat diseases in tropical Africa are limited to specific regions or specific diseases. This study aimed to review scientific research findings on goat diseases in tropical Africa, focusing on their prevalence and impacts on production performance. All main diseases, such as parasitic, viral, and bacterial diseases, are included in the present study. Studies conducted in different countries have revealed high prevalence rates of gastrointestinal parasites exceeding 95%. These parasites resulted in growth retardation and reduced carcass weight at slaughter. Management of mites could decrease production and reproductive function. Trypanosomiasis led to decreased hematocrit levels, abortions, low birth weight, and high kid mortality. The prevalence of trypanosomiasis has been different across regions, ranging from 2.95% to 57.1%. Peste des Petits Ruminants has been reported in many African countries, causing significant outbreaks with seroprevalence rates ranging from 30% to 55%. Rift Valley fever was characterized by high mortality in adult goats (20-30%) and numerous abortions 2 weeks after infection, with seroprevalence rates ranging up to 25.8%. Contagious Caprine Pleuropneumonia indicated high morbidity (approximately 100%) and high mortality (80% to 100%), with prevalence ranging from 22% to 39% in abattoirs and from 35% to 52% in farms. Brucellosis did not affect the weight of infected animals but reduced litter size in goats and disrupts vital organs. This review highlights the extent of goat diseases in tropical Africa to determine the most appropriate prevention and control strategies.

**Keywords:** Control strategy, Goat diseases, Prevalence, Production performance, Tropical Africa

## INTRODUCTION

Like other agricultural activities, livestock production in tropical areas faces major sustainability challenges (Dedieu et al., 2011). Despite its numerous potential advantages and multifunctionality in household security, goat farming also confronts these very challenges (Dubeuf, 2011). In many African countries with tropical climates, ruminants, especially goats, play a crucial role in agricultural systems (Nair et al., 2021).

Goats could adapt to difficult climatic conditions and limited resources, especially in arid and semi-arid regions where crop production is uncertain in the context of climate change (Monau et al., 2020). Goats play a vital role in maintaining food security and the economic livelihoods of smallholder farmers in rural areas (Monau et al., 2020). A study by Dubeuf (2011) emphasizes the importance of supporting agricultural research, particularly in areas where goat farming has development potential. However, infectious and parasitic diseases pose a major challenge for small ruminant production, as they are one of the main causes of animal mortality on farms (Missohou et al., 2016; Armson et al., 2020). These diseases, including infectious and parasitic diseases, hurt the economy due to livestock mortality and high treatment costs (Adeyemo et al., 2022). They also pose a risk to public health, as some of these diseases can be transmitted from animals to humans (Lancelot et al., 2011). Several research studies have been conducted on these diseases in several tropical climate countries. These research studies have identified the pathogens responsible for goat diseases, whether bacterial, viral, parasitic, or fungal. Furthermore, the prevalences, clinical manifestations, and effects of these diseases on goat production have been described. However, available studies are often fragmented and limited to specific regions or particular diseases (Misinzo et al., 2015; Arsevska et al., 2016; Baron et al., 2016). Therefore, a comprehensive synthesis of research on this topic is necessary to assess the extent of goat diseases in tropical Africa and determine the most appropriate prevention and control strategies. An in-depth review of goat diseases and control and

REVIEW ARTICLE  
pII: S232245682300047-13  
Received: 26 June 2023  
Accepted: 08 August 2023

prevention strategies in Africa is crucial. It can help maintain the availability of goat products, thus contributing to both food security and livelihoods. It can also reduce economic losses and ensure the economic viability of breeders. In addition, such a review can identify potential risks to human health, paving the way for implementing appropriate measures to mitigate them. Developing prevention and control strategies specifically adapted to goat diseases in tropical Africa would also be possible. The present study summarizes the diseases encountered in tropical African goat farming by focusing on their prevalence and their effects on goat production performance.

## PARASITIC DISEASES

### Gastrointestinal parasites of goats in tropical Africa

#### *Impact of gastrointestinal parasites on the production performance of goats*

In tropical countries, gastrointestinal parasites are highly involved in helminth infections in small ruminants (Githiori et al., 2004; Mpfu et al., 2022). Economically, gastrointestinal strongylosis, especially haemonchosis, is widely acknowledged as a leading cause of health problems in ruminant animals, resulting in substantial economic losses in the livestock industry due to reduced production (Arsenopoulos et al., 2021). The presence of worms in livestock farms influences both the quantity and quality of production (Devadharshini et al., 2022). Gastrointestinal parasites are responsible for growth retardation in kid goats, resulting in reduced carcass weights at slaughter (Arsenopoulos et al., 2021). In dairy females, infestations by digestive *strongyles* have been consistently associated with decreased milk production (Mpfu et al., 2022). In Benin, a study on helminth infections in sheep and goats showed that a moderate worm burden of *haemonchus contortus* induced detectable anemia during the dry season (Attindehou et al., 2012). Naturally resistant animals show an increased immune response against these parasites, reducing the need for drug treatments. This resistance can result from genetic changes, including factors such as carriers and metabolism. Genetic differences play a significant role in susceptibility to gastrointestinal parasites, both across various breeds and species, as well as within genes that are vital for the immune response (Mpfu et al., 2022).

#### *Methods of diagnosing gastrointestinal parasites in goats*

Clinical diagnosis of strongylosis is based on clinical signs, such as weight loss, diarrhea, anemia, and rough hair coat (Zajac and Garza, 2020). These symptoms are generally common and nonspecific (Lotfalizadeh et al., 2022). Therefore, it is necessary to identify the responsible parasites using appropriate parasitological diagnostics to confirm the clinical diagnosis and establish an appropriate prevention plan. There are direct methods focused on coproscopic techniques for diagnosing strongylosis (Alowanou et al., 2021). The classical quantitative technique McMaster is the most popular in the field of veterinary parasitology (Alowanou et al., 2021). In addition, it is advocated by the World Association for the Advancement of Veterinary Parasitology in its guidelines for evaluating the efficacy of anthelmintic drugs in ruminants, based on the work of Cringoli et al. (2010). However, there are newer quantitative methods that are more sensitive, precise, and accurate, such as the flotation technique (FLOTAC) and Mini-FLOTAC, developed in the last decade (Cringoli et al., 2017). The FLOTAC technique utilizes a specific centrifuge and is highly sensitive, while Mini-FLOTAC is a variant that does not require a centrifuge and is based on passive flotation with fewer preparation steps (Cringoli et al., 2017). The Mini-FLOTAC method consists of a passive fluctuation of parasitic structures, which makes it possible to diagnose helminth eggs/larvae as well as protozoan oocysts or cysts at the same time (Cringoli et al., 2017). In their study on the assessment of the Mini-FLOTAC and McMaster methods for the detection of gastrointestinal parasites in West African sheep, dwarf goats, and crossbred rabbits, Alowanou et al. (2021) concluded that the Mini-FLOTAC method showcased enhanced diagnostic performance in terms of prevalence. As a result of this finding, they suggested that the Mini-FLOTAC method held greater reliability and could potentially serve as a more dependable alternative for veterinary clinics. Other research has also confirmed that the Mini-FLOTAC method has higher sensitivity and efficiency than the McMaster technique in identifying parasites in human and other animal samples (Barda et al., 2013; Dias de Castro et al., 2017; Noel et al., 2017). The method frequently employed in laboratory evaluations to ascertain worm types in living animals involves the identification of larvae in freshly expelled feces (mainly lungworm larvae) or those maturing in fecal cultures (gastrointestinal nematodes, Van Wyk and Mayhew, 2013). However, only an individual with expertise can accurately distinguish the larvae (Van Wyk and Mayhew, 2013). For numerous genera of nematodes, discerning features, such as the morphology of the anterior end (head) of the larvae, are nearly identical (Van Wyk and Mayhew, 2013). In order to address this issue, Van Wyk and Mayhew (2013) devised an innovative method that entails determining the portion of the larval sheath tail extension containing a delicate, whip-like filament at the tip.

#### *Prevalence of gastrointestinal parasites in goats across some tropical African countries*

It was reported that more than 95% of small ruminants in tropical regions suffer from gastrointestinal parasite infestations (Terefe et al., 2012). In the context of Benin, parasitological surveys indicated that goats primarily indicated infestations of *coccidia* (92.24%) and *strongyles* (83.91%), followed by *Strongyloides* spp. (73.25%, Challaton et al., 2023). Additionally, *Moniezia* spp. infestations were observed at a rate of 21.8%, along with other gastrointestinal parasites, such as *Trichostrongylus axei* spp. (0.94%) and *Toxocara* spp. (0.28%, Challaton et al., 2023). A gastrointestinal parasitic

investigation conducted by [Faihun et al. \(2017\)](#) on small ruminants (sheep and goats) and wild ruminant species (harnessed bushbuck) revealed the presence of six different parasites in sheep and goats (including *strongyles*, *Strongyloides*, coccidian, cestodes, *Capillaria*, and trematodes) with prevalence rates ranging from 60% to 100%. In a study conducted by [Ipungu et al. \(2017\)](#) on goats raised in the municipality of Lubumbashi city in Congo, an infestation rate of 89.6% was observed during the dry season and 91.2% during the rainy season, caused by gastrointestinal *strongyles* as well as a lungworm (*Dictyocaulus* spp). The identified parasites included *Dictyocaulus* spp. (88.6%), *Trichuris ovis* (60%), *Capillaria* spp. (29.5%), *Ostertagia* spp (20%), *Trichostrongylus* spp (30.5%), *Charbetia ovina* (8.6%), and *Strongyloides papillosus* (1%, [Ipungu et al., 2017](#)). In an epidemiological study on gastrointestinal helminths of goats in Middle Guinea, [Barry et al. \(2002\)](#) observed prevalence rates of 94% for *strongyles*, 25% for *Trichuris*, 20% for *Strongyloides*, and 12% for *Moniezia*. The results of a study conducted in Cameroon by [Malla et al. \(2021\)](#) revealed a high prevalence of 89.0% for gastrointestinal parasites in goats, including *Trichostrongylus* spp. and *Haemonchus* spp., *Strongyloides* spp., and Protozoa. In Ethiopia, a study conducted by [Chalchisa et al. \(2015\)](#) revealed a prevalence of 90.1% for mixed strongyle infestation in goats. The most common genera of *strongyles* in this study were *Haemonchus* (76.1%), *Trichostrongylus* (56.3%), *Oesophagostomum* (33.1%), *Bunostomum* (20.1%), and *Chabertia* (5.2%).

#### **Methods of controlling gastrointestinal parasites in goats**

The treatment of digestive parasitoses in small ruminants in tropical Africa is generally carried out using synthetic chemical products, such as benzimidazoles, imidazothiazoles, and avermectins ([Arsenopoulos et al., 2021](#)). However, the use of these molecules is limited by several factors, including parasite resistance, low vaccine coverage, inaccessibility to modern veterinary products for rural farmers, high cost of products and treatments, and unapproved products ([Sargison and Scott, 2003](#); [Fajimi and Taiwo, 2005](#)). Complementary approaches, such as fodder or bioactive plants/ethnoveterinary products, biological control using fungi, as well as grazing and nutritional management strategies are employed to control these parasites ([Mpofu et al., 2022](#)). Many farmers return to medicinal herbs as a natural alternative to synthetic chemical products ([Hounzangbe-Adote., 2001](#)). These herbs have been used for a long time to treat digestive parasitoses in small ruminants ([Sadr et al., 2022](#)). The study by [Tchetan et al. \(2021\)](#) listed several medicinal plants whose efficacy against gastrointestinal parasites has been confirmed both by farmers and traditional healers, as well as through scientific results. These plants, rich in bioactive molecules, such as flavonoids, tannins, and alkaloids, are easily accessible to farmers ([Tchetan et al., 2021](#)).

#### **Acarioses (mite agents of scabies)**

##### **Impact of scabies on the production performance of goats**

Mites infesting animals have significant economic implications due to factors such as slowed growth, decreased daily weight gain, treatment, skin and leather damage, and labor costs, as well as mortality losses ([Arul et al., 2023](#)). In Ethiopia, ectoparasites were responsible for 35% of sheep skin waste and 56% of goat skin waste in the market ([Kebede, 2013](#)).

##### **Methods of diagnosing scabies in goats**

Methods for diagnosing scabies in goats include careful clinical examination for characteristic signs such as itching, crusting and hair loss, papules, erythema, excoriations, desquamation, and thick wrinkled skin ([Chuka et al., 2020](#)). For direct microscopic examination, skin lesion samples are taken and dissolved in a 10% potassium hydroxide solution, centrifuged, and then observed under the microscope for the presence of mites or their structures ([Benti et al., 2020](#); [Chuka et al., 2020](#)). These skin lesion samples can be fixed in 10% neutral buffered formalin, followed by processing, staining with hematoxylin and eosin, and finally, microscopic observation to identify specific features of scabies, such as epidermal hyperplasia, hyperkeratosis, as well as the presence of mast cells, eosinophils or lymphocytes in the dermis ([Chuka et al., 2020](#)).

##### **Prevalence of scabies in goats across some tropical African countries**

A study carried out in southern Benin by [Salifou et al. \(2013\)](#) regarding the prevalence of scabies in small ruminants, along with affected owners suffering from the same disease, identified the species *Sarcoptes scabiei* with a prevalence rate of 28.33% and 9.5% in animals and small-scale farmers (human cases), respectively. Infestations were notably more frequent among goats (39.6%). These findings imply a significant correlation between suspected human cases and these animals ([Salifou et al., 2013](#)). The same species (*Sarcoptes scabiei*) had been identified in the northern region of Benin by [Tassou \(2009\)](#). Another research showed the existence of another variety of mites of the genus *Psoroptes* in the northern region of Couffo in Benin ([Koudandé, 2006](#)). The species *Psoroptes comunis* and *Sarcoptes scabiei* were encountered in goats with a prevalence of 58.1% in a study conducted by [Davou et al. \(2017\)](#) in north-central Nigeria. In southwestern Nigeria, [Ogundiya et al. \(2012\)](#) reported a prevalence of 0.53% of scabies in goats caused by *Sarcoptes scabiei aprae*. In Ethiopia, the combined prevalence of *Sarcoptes scabiei* in sheep and goats in a meta-analysis was estimated at 4.4%. The study further revealed that mites of the genus *Sarcoptes*, *Demodex*, and *Psoroptes* were the most prevalent mites infesting small ruminants in the country ([Asmare et al., 2016a](#)).

##### **Methods of controlling scabies in goats**

The treatment and prevention of *Sarcoptes scabiei* are carried out using various synthetic chemical products such as ivermectin, diazinon, phoxim, and coumaphos (Benti et al., 2020). Dipping with other insecticides can also be useful in combating parasites and preventing secondary bacterial complications (Benti et al., 2020). Maintaining hygiene in housing and avoiding overcrowding can minimize the accumulation and spread of mites (Benti et al., 2020). Currently, there is no commercial vaccine available to protect animals against scabies caused by mites (Benti et al., 2020). Farmers also use traditional methods, such as plants, for scabies control in Africa (Dassou et al., 2014; Yasmine et al., 2015). Prevention and control of scabies in small ruminants remain an important concern for animal and human health, as well as for the livelihoods of farmers (Salifou et al., 2013).

## Trypanosomiasis

### *Impact of trypanosomiasis on the production performance of goats*

In a study in Nigeria to investigate the susceptibility of Sahelian goats to experimental *Trypanosoma vivax* infection, Akinwale et al. (2006) observed a reduction in hematocrit levels two weeks following infestation. In Burkina Faso, it was provided evidence of the impact of the *Trypanosoma* spp. on hematocrit levels by comparing the mean hematocrit levels of infected and uninfected animals (Ye, 2012). Infected animals had a mean hematocrit level of 17.75%, while uninfected animals had a level of 26.03% in small ruminants, suggesting a significant decrease in hematocrit levels in infected animals, compared to uninfected ones. Similar observations were made by Ezebuoro et al. (2009) in Nigeria on the prevalence of trypanosomiasis in commercial livestock, where the hematocrit was 20.29% in trypanosomiasis-infected goats and 31.56% in uninfected goats. In a study by Faye et al. (2004), which examined the impact of *Trypanosoma congolense* infection on the reproductive performance of West African Dwarf (WAD) goats, results showed that the infestation resulted in elevated rectal temperature (38.8°C), abortions in 27.8% of infected goats, and a decrease in birth weight of offspring born to infected goats. In addition, a mortality rate of 61.5% was observed in kids born alive from infected goats during their first week of life. Furthermore, the concentrations of pregnancy-associated glycoprotein (PAG) and plasma progesterone were found to be lower in infected animals compared to the control group. Milk production and quality in dairy goats are also affected by trypanosomiasis. According to Lopes et al. (2016), goats infected with *Trypanosoma vivax* showed a rapid decline in milk production, a flattened lactation curve, reduced lactation persistence, as well as a significant decrease in milk fat and protein content. These results express that *Trypanosoma vivax* (*T. vivax*) infestation can have negative economic consequences on the milk production of goats.

### *Diagnosis methods of trypanosomiasis in goats*

Animal trypanosomiasis caused by *T. vivax*, *T. congolense*, and *T. brucei* is the most significant vector-borne disease in ruminants (Morrison et al., 2023). These diseases are characterized by fever, anemia, and weight loss (Tariq et al., 2022; Morrison et al., 2023). Several diagnostic methods are used to diagnose *Trypanosoma* infection in animals when there are no clinical signs. Classical approaches to identifying *Trypanosoma* spp. involve microscopic examination of fresh or stained blood smears (Tariq et al., 2022). *Trypanosomes* can be identified in blood samples using a light microscope at 40X magnification (Tariq et al., 2022). Another step is to examine blood samples stained with Giemsa stain (Mafie et al., 2018). The Primo Star iLED LED microscope from Carl Zeiss, and FIND is a breakthrough, enabling fluorescence and bright-field microscopy (Tariq et al., 2022). The use of acridine orange fluorescence and Giemsa staining improves sensitivity for detecting trypanosomes in blood (Tariq et al., 2022). The formalin gel test (FGT) and ELISA are used to detect antibodies (Tariq et al., 2022). The formalin gel test (FGT) involves mixing one milliliter of serum with a solution of concentrated formalin (Tariq et al., 2022). If the serum immediately coagulates and becomes opaque, the result is considered positive (Tariq et al., 2022). ELISA involves antigen preparation, washing, incubation, and detection (Tariq et al., 2022). Polymerase chain reaction (PCR) is a more sensitive molecular method, particularly real-time PCR, which can identify subspecies (Tariq et al., 2022).

### *Prevalence of trypanosomiasis in goats across some tropical African countries*

In southern Cameroon, Simo et al. (2005) observed a prevalence of 20% for *T. brucei*, 4.2% for *T. gambiense*, 15.2% for *T. vivax*, and 7.2% for *T. congolense* in goats. In a study conducted in a peri-urban area of Togo, Bastiaensen et al. (2003) observed an average prevalence of trypanosomiasis of 8.41% in goats. In the agropastoral zones of Sidéradougou, Samorogouan, and Barani in southern Burkina Faso, Sow et al. (2014) recorded a trypanosomiasis prevalence of 2.95%. The *trypanosomes* responsible for these infestations were primarily *T. vivax* or *T. congolense*. In Kenya, goats are most commonly carriers of *T. vivax* (O Ng'ayo et al., 2005). In the Mongo regions of southern Gabon, Maganga et al. (2020) recorded a prevalence of 7.8% of goats infected with *T. vivax*, *T. simiae*, *T. simiae Tsavo*, *T. congolense*, and *T. brucei*.

### *Methods of controlling trypanosomiasis in goats*

Chemotherapy represents the main means of controlling *trypanosome* infestation in goats (Jaiswal et al., 2015). There are several chemical compounds used to treat trypanosomiasis, and among these, diminazene aceturate is the most commonly used trypanocide (Tariq et al., 2022). In addition to diminazene aceturate, other compounds such as

isometamidium chloride, suramin, quinapyramine sulfate, and quinapyramine chloride are also available (Jaiswal et al., 2015, Tariq et al., 2022). Traditional medicine can also be considered an effective alternative to modern medicine in the fight against trypanosomiasis (Andre et al., 2017). According to Andre et al. (2017), aqueous extracts of *Guiera senegalensis* leaves showed trypanocidal activity against *T. brucei* in Burkina Faso.

Prevention and control of trypanosomiasis in goats require continuous monitoring of animal health, vector control measures, and appropriate treatments to minimize economic losses and impacts on food security. Therefore, clinical and experimental trials aimed at improving the prevention, diagnosis, and treatment of these diseases are necessary to help control and eradicate animal trypanosomiasis in Africa.

## VIRAL DISEASES

### Peste des petits ruminants

#### *Impact of peste des petits ruminants on the production performance of goats*

In Senegal, a study on the effect of peste des petits ruminants (PPR) on the productivity of goat herds showed that herds exposed to PPR experienced a threefold increase in natural mortality rates, four times more abortions in females, and a 30% decrease in birth rates and fertility (Grech-Angelini, 2012). In Nigeria, cases of PPR-related abortions were reported by Asuku et al. (2022). This study highlights the negative impact of the disease on sheep and goat production, emphasizing the risks of mortality and potential economic losses associated with abortions. Mortality rates of 53% have been recorded among goats in Nigeria (Ameh et al., 2000). In Africa, 25 countries have reported PPR epidemics in sheep and goats (African Union-Interafrican Bureau for Animal Resources, 2014). The significant outbreaks reported in Zambia, Tunisia, Uganda, Mongolia, Georgia, Liberia, Kenya, China, and Algeria, as well as on island environments in the Maldives and Comoros, resulted in a mortality of over 17,000 small ruminants (Jebara et al., 2012). A study conducted by Kindji (2006) in Northeast Benin on the socioeconomic impact of PPR estimated economic losses due to mortality at 8,224,07 USD, those related to morbidity at 647,30 USD, and 213,82 USD for the treatment of affected small ruminants. During the 2006-2008 PPR epidemic in Turkana, Kenya, more than a million animals perished and the total value, in terms of lost production, was estimated at USD 2.4 million (Njeumi et al., 2020). In the United Republic of Tanzania, during the same epidemic between 2006 and 2008, a total of 64,661 animals were slaughtered. Meanwhile, in Côte d'Ivoire, during the same period, affected animals were sold at half their usual market price (OIE, 2017).

#### *Clinical signs and methods of diagnosing peste des petits ruminants in goats*

Phylogeographic studies have traced the major axes of PPR dispersion and differentiation (Muniraju et al., 2014). According to Misinzo et al. (2015), *lineage I and II* viruses are predominantly present in West and Central Africa, while *lineage III* viruses are more common in East Africa and the southern Middle East. *Lineage IV* viruses are mainly localized in Asia and the Middle East, although some strains have been identified in North and East Africa (Misinzo et al., 2015). The clinical diagnosis of PPR is complex due to similarities with other diseases, such as *vesicular stomatitis* (SPPV) and *stomatitis virus* (GTPV) infections (Zewdie et al., 2021), rinderpest, pneumonic pasteurellosis and contagious caprine pleuropneumonia (Kinimi et al., 2020). The symptoms and lesions do not allow a clear distinction between these diseases. Skin lesions evolve from papules to nodules, vesicles, and pustules, then form scabs (Zewdie et al., 2021). Affected animals become weak, lose their appetite, run a high fever, and have breathing difficulties due to lesions in the respiratory tract and lungs. Lesions also appear in the mouth, nose, and eyelids, accompanied by excessive salivation. Mucous membranes become necrotic and ulcerated, leading to diarrhea in the case of nodules in the intestines (Zewdie et al., 2021). Histopathologically, skin changes (hyperkeratosis, acanthosis, hyperkeratinization, edema, degeneration of sebaceous glands and hair follicles) as well as lung lesions and proliferative alveolitis with occasional cytoplasmic inclusions in alveolar cells and macrophages are observed (Zewdie et al., 2021). In addition, inoculation experiments show consistent lesions and antigens in skin, lungs, and lymph nodes following inoculation of vesicular SPPV and GTPV in sheep and goats, with immunohistochemical detection of viral antigens (Zewdie et al., 2021). Serological tests include methods such as serum neutralization tests (VNT, SNT), indirect immunofluorescence assay (IFAT), and agar gel immunodiffusion assay (AGID, Kinimi et al., 2020; Zewdie et al., 2021). More sensitive and sophisticated tests, such as immunocapture ELISA and quantitative real-time PCR (qRT-PCR), are also available (Njeumi et al., 2020).

#### *Prevalence of peste des petits ruminants in goats across some tropical African countries*

Across the world, serological surveys have shown prevalence rates ranging from 30% to 45%, and even 55% in countries where PPR is enzootic (Diallo, 2006). Table 1 presents the prevalence rates of PPR in tropical Africa.

#### *Methods of controlling peste des petits ruminants in goats*

There is no specific treatment for this viral disease, but antibiotics can be administered to treat secondary bacterial infections, while careful nursing care is recommended to reduce morbidity and complications (Zewdie et al., 2021). Methods to combat PPR rely on sanitary prophylaxis measures. Vaccination against PPR is used as an effective method to control the spread of the disease and reduce mortality rates among animals in tropical Africa (Alemnew et al., 2022;

Loum Gazida et al., 2022). Capripox-inactivated vaccines have been shown to be safe and effective. However, they require two doses to confer prolonged immunity (Zewdie et al., 2021). It was reported that vaccination can protect PPR for up to 1 year after vaccination and possibly for the economic lifespan of vaccinated animals (Diallo, 2006). Njoya et al. (2005) reported a highly significant effect of PPR vaccination with Bovipestovax on reducing small ruminant mortality in a peasant farming environment in the Sudano-Sahelian zone of Northern Cameroon. In Ethiopia, Alemnew et al. (2022) indicated that vaccination led to a significant decrease in PPR prevalence and a reduction in mortality rates among animals. These findings confirm the report of Loum Gazida et al. (2022), which supports that increasing mass vaccination campaigns, maintaining the cold chain to preserve vaccines, and the serious commitment of vaccination agents are the only possibilities for completely eradicating PPR. It is therefore important to continue efforts to combat and prevent PPR, especially through herd vaccination, to reduce its impact on animal health, food security, and the livelihoods of local populations.

**Table 1.** Seroprevalence of Peste des Petits Ruminants in goats of Tropical Africa from 2006 to 2022

Countries	Regions	Sample (Number)	Prevalence (%)	Reference
Benin	Departments of Borgou and Alibori in Northeast Benin	330	24.08	(Kindji, 2006)
Burkina Faso	Soum, located in the north of the country	878	23.01	(Sow et al., 2008)
Chad	Southern region of the country	1,699	48.9	(Mahamat et al., 2018)
Djibouti	Ali Sabieh, Arta, Dikhil, Djibouti, Obock et Tadjourah	1,215	6.83	(Moumin et al., 2018)
Ethiopia	National survey	4,585	9.4	(Waret-Szkuta et al., 2008)
	Dugda and Adami Tullu districts in the Showa administrative zone and Dodota district in the Arsi zone of the Oromia	407	46.68	(Gari et al., 2017)
Ghana	Regional State 74 villages in the 31 districts of the 10 regions of the country	1,534	45.50	(Otsyina et al., 2013)
Kenya	Turkana County	538	40	(Kihu et al., 2015)
Mali	The entire territory except for the Kidal region in the far northeast of the country	1,784	42.4	(Kamissoko et al., 2013)
Niger	Arid zone (Niamey, Tillabéry, and Tahoua) of the country	266	47.9	(Farougou et al., 2013)
Nigeria	The semi-arid region in the northeast of the country	1,571	50.4	(El-Yuguda et al., 2013)
	South-East, South-South, Centre-North, North-West and South-West agro-ecological zones	3,489	22.93	(Woma et al., 2016)
Rwanda	Bugesera, Kirehe, and Nyagatare districts in the Eastern Province of Rwanda; Gicumbi and Musanze districts in the Northern	316	13.6	(Shyaka et al., 2021)
Sudan	Province Northern, Eastern, Central, and Western states	1,459	48.2	(Intisar et al., 2017)
	Arusha, Manyara, and Kilimanjaro regions in the north of the country	892	49.5	(Swai et al., 2009)
Tanzania	Ngorongoro in the northern region of Arusha, and Ulanga, Kilombero, and Mvomero in the southeastern region of Morogoro	238 and 323	48.3 and 10.8	(Torsson et al., 2017)
Uganda	Karenga district in the Karamoja region in the northeast of the country	569	43.8	(Akwongo et al., 2022)

## Rift valley fever

### *Impact of Rift Valley fever on the production performance of goats*

The Rift Valley fever virus (RVFV) is now believed to be endemic in certain countries in Africa with tropical or subtropical climates, such as Senegal, Gambia, Mauritania, Zimbabwe, Namibia, South Africa (Chevalier et al., 2009; Paweska et al., 2010). Several species of mosquitoes, including the Aedes and Culex genera, act as vectors for transmitting the virus to animals (Pepin et al., 2010). In goats and sheep, this disease is primarily characterized by high stillbirth rates (20-30% in adults) and numerous abortions two weeks after herd infection (Adam et al., 2010; Archer et al., 2011). During an epidemiological study conducted by Munyua et al. (2010) on Rift Valley Fever in livestock of Kenya, the predominant symptoms were death, hemorrhagic syndrome, and miscarriage. These manifestations were observed in all species, affecting up to 30% of infected herds. Other clinical markers included labored breathing, coughing, loss of appetite, debility, and extensive nasal and oral secretions. In terms of disease and mortality rates, goats showed the highest percentages (4.6% and 1.9%), followed by sheep (1.5% and 0.3%, respectively). The high mortality and abortion rates within infected herds result in significant economic losses (Adam et al., 2010; Archer et al., 2011).

### ***Impact of Rift Valley fever on human health***

In humans, transmission mainly occurs through direct contact with blood or other body fluids from infected animals (Arsevska et al., 2016). From November 30, 2006 to March 12, 2007, 684 cases were reported with a mortality rate of 23% in Kenya (WHO, 2007). During the first reported Rift Valley fever epidemic in Niger, 346 suspected human cases were reported, of which 38 were confirmed, resulting in 32 deaths (Doutchi et al., 2017). Patients had common risk factors such as mosquito bites, direct contact with dead or aborted animals, and regular consumption of milk from these animals (Doutchi et al., 2017). Ibrahim et al. (2021) observed a seroprevalence of 13.2% in humans during a study conducted on livestock and humans in the Somali region of Ethiopia. Due to the risks to human and animal health, as well as significant economic losses, it is crucial to maintain continuous epidemiological surveillance and implement prevention and control measures to prevent the spread of this disease.

### ***Methods of diagnosing Rift Valley fever in goats***

Real-time reverse transcription-polymerase chain reaction (qRT-PCR) is the current diagnostic method for Rift Valley Fever (RVF). It is a precise and sensitive molecular approach used to detect viral RNA (Sado et al., 2022). ELISA tests are also frequently used to detect antibodies produced in response to Rift Valley Fever virus infection (Ibrahim et al., 2021; Sado et al., 2022). The RVFV neutralization test is also used to assess the presence of RVFV-specific neutralizing antibodies in serum samples (Troupin et al., 2022).

### ***Prevalence of Rift Valley fever in goats across some tropical African countries***

The prevalence of RVF varies widely from region to region, ranging from 0.0% to 25.1% (Table 2). High-prevalence areas include the south-eastern shore of Lake Chad in Chad (18.8%), Zambezia province in Mozambique (25.1%), and Garissa County in Kenya (25.8%, Table 2). On the other hand, some regions report little or no prevalence (Table 2).

### ***Methods of controlling Rift Valley fever in goats***

Means of controlling RVFV include epidemiological surveillance to detect cases early, vector control through the destruction of mosquito habitats and the use of insecticides, and restricting the movement of infected animals (Fawzy and Helmy, 2019). Routine vaccination is considered the main means of controlling RVFV infections in animals in endemic countries to prevent human infections, socioeconomic losses, and epidemics (Fawzy and Helmy, 2019). Several vaccines are used, including the formalin-inactivated vaccine with alum adjuvant (Menya/sheep/258), the binary ethylenamine-inactivated vaccine with alum adjuvant (ZH501 RVF) from the Veterinary Serum and Vaccines Research Institute (VSVRI), as well as the live attenuated neurotropic Smithburn strain produced by VSVRI (Arsevska et al., 2016; Fawzy and Helmy, 2019).

**Table 2.** Seroprevalence of Rift Valley Fever in goats of Tropical Africa from 2014 to 2022

Countries	Regions	Sample (Number)	Prevalence (%)	Reference
Burkina Faso	Provinces of Yatenga, Seno, and Soum in the northern regions of the country	120	6.66	(Boussini et al., 2014)
	Tsinga livestock market in the Yaoundé 2 district	168	2.4	(Sado et al., 2022)
Cameroon	Lagdo, Pitoa, Boklé, Garoua, Kismatari, Poli, Touboro, and Dembo	355	2.3	(Poueme et al., 2019)
Chad	Southeastern shore of Lake Chad	144	18.8	(Abakar et al., 2014)
Congo	Mongala, Sud Ubangi, Nord Ubangi, Kwilu, Lomami, South Kivu, and Tanganyika	672	0.0 to 23.81	(Tshilenge et al., 2019)
Gabon	Mongo County in the southwest of the country	106	4.72	(Maganga et al., 2017)
Ethiopia	Adadle Woreda in the Somali region of the country	297	6.3	(Ibrahim et al., 2021)
Guinea	Prefectures of Beyla, Boffa, Boké, Coyah, Dabola, Dalaba, Faranah, Forécariah, Gaoual, Guéckédou, Kindia, Koundara, Kouroussa, Macenta, Mamou, Mandiana, and N'zérékoré	408	1.00	(Troupin et al., 2022)
Kenya	Garissa County	271	25.8	(Nanyingi et al., 2017)
	Nyandarua County	19	10.53	(Wanjama et al., 2022)
Mali	Mopti and Sikasso	290	3.1	(Dione et al., 2022)
Mauritania	Central and southern parts of the country	294	1.4	(Rissmann et al., 2017)
Mozambique	Zambezia Province	187	25.1	(Blomström et al., 2016)
Nigeria	Bodija Municipal Abattoir in Ibadan, southwest of the country	44	2.3	(Opayele et al., 2019)
Central African Republic	Ngawi market in the central commercial area of Bangui and the village of Ndangala located south of Bangui	219	5	(Nakouné et al., 2016)
Tanzania	East and West of the Rift Valley in the country	531	22	(Sindato et al., 2015)

## BACTERIAL DISEASES

### Respiratory mycoplasmosis (Contagious caprine pleuropneumonia)

#### *Impact of respiratory mycoplasmosis on the production performance of goats*

Respiratory mycoplasmosis is characterized by severe serofibrinous pleuropneumonia, high morbidity (approximately 100%), and mortality (80-100%, [Yatoo et al., 2019](#)). [Akwuobu et al. \(2016\)](#) conducted an experimental study to investigate the effects of *Mycoplasma ovipneumoniae* and *Mycoplasma arginini* infections on WAD goats under one year old. The results indicated that all infected goats exhibited respiratory signs, such as coughing and nasal discharge, along with observed lung lesions in each case. Mortalities were also recorded in the infected groups, suggesting that infection with these bacteria is dangerous in young goats.

#### *Methods of diagnosing respiratory mycoplasmosis in goats*

Methods for identifying contagious caprine pleuropneumonia (CCPP) encompass differences in approaches. Firstly, serological tests are employed based on rapid latex agglutination techniques using the CapriLAT kit (RAI6224, [Olorunshola et al., 2020](#)). These tests are specifically designed to detect the presence of antibodies to PPCC in biological samples, specifically blood serum ([Olorunshola et al., 2020](#)). However, despite their usefulness, it should be noted that bacterial culture is considered the most indisputable reference for the diagnosis of CCPP. This method provides direct confirmation of the presence of the bacteria. Nevertheless, it is important to note that this method is slow ([Abd-Elrahman et al., 2020](#)). Mycoplasma growth and isolation are performed using agar and broth culture media (M0535-250G, Sigma Aldrich, USA). For selective isolation of *Mycoplasma* species, a special approach is employed, namely incubation on a specific agar medium (CC1A, Mycoplasma Experience Ltd. Product) is carried out for 7 days ([Abd-Elrahman et al., 2020](#)). In addition to these methods, CCPP can be identified by PCR. This molecular biology technique amplifies the DNA of the bacteria present in the sample, enabling it to be detected with a high degree of accuracy ([Abd-Elrahman et al., 2020](#)).

#### *Prevalence of respiratory mycoplasmosis in goats across some tropical African countries*

A pooled estimate of CCPP prevalence of 23.21% in tropical Africa was reported by [Ahaduzzaman \(2021\)](#), including 21.87% in Ethiopia, 47.22% in Kenya, 8.06% in Mauritius, 3.70% in Nigeria, and 37.13% in Tanzania. In a study carried out in Nigeria concerning the separation of *Mycoplasma mycoides* subspecies *Mycoides* in small ruminants, [Egwu et al. \(2012\)](#) detected various strains of mycoplasma in the lungs of both sheep and goats, whether they were affected by the condition or not. The identified strains encompassed *Mycoplasma ovipneumoniae* (30%), *Mycoplasma mycoides* subspecies *capri* (29.5%), *Mycoplasma mycoides* subspecies *mycoides* (13.5%), *Mycoplasma capricolum* (11.5%), with the least frequently isolated being *Mycobacterium bovis* (1.5%). [Akwuobu et al. \(2014\)](#) reported a prevalence of 29.8% for CCPP in a separate study in north-central Nigeria, during which they identified two species of mycoplasma, namely *Mycoplasma ovipneumoniae* and *Mycoplasma arginini*. Contagious caprine pleuropneumonia is prevalent in the districts of Agago and Otuke in Uganda, with seroprevalence rates of 17.7% and 23.3% in unvaccinated goats, respectively ([Atim et al., 2016](#)).

#### *Methods of controlling respiratory mycoplasmosis in goats*

Contagious caprine pleuropneumonia is treated with several antibiotics, among which macrolides, particularly tylosin, are considered the most effective ([Yatoo et al., 2019](#)). The antibiotics marbofloxacin and spiramycin have also proved effective against CCPP ([Abd-Elrahman et al., 2020](#)). The use of oxytetracycline has also been effective in treating CCPP, but prolonged use can lead to undesirable side effects in young goats, such as congenital malformations and an increased risk of potential resistance ([Yatoo et al., 2019](#)). Additionally, prophylactic use of minimal antibiotic doses may enable mycoplasmas to tolerate these treatments better ([Yatoo et al., 2019](#)). Other antibiotics such as fluoroquinolones (enrofloxacin, danofloxacin), aminoglycosides (streptomycin), and pleuromutilin (tiamulin) are also used to treat CCPP ([Yatoo et al., 2019](#)). In addition to antibiotic treatments, there are preventive measures to combat CCPP in goats. Vaccination is considered an effective method for the prevention of disease. Several vaccines have been developed against CCPP, including a whole-cell inactivated vaccine, a live attenuated vaccine, and a recombinant subunit vaccine ([Yatoo et al., 2019](#)). Farmers also employ traditional methods for CCPP control. In Kenya, the majority of surveyed farmers use medicinal plants (*Solanum aculeastrum*, *Albizia coriaria*, *Ekebergia capensis*, *Piliostigma thonningii*, *Euclea divinorum*) to treat respiratory disease symptoms, such as deep dry cough, extended neck, fever, and weight loss ([Kama-Kama et al., 2016](#)).

## Brucellosis

#### *Impact of brucellosis on the production performance of goats*

Brucellosis can have detrimental consequences for agricultural economies, both in the short and long term, by causing abortions and infertility in animals, a decrease in milk production, as well as the birth of weakened offspring ([Dahmani et al., 2022](#)). A study conducted by [Mahboub et al. \(2013\)](#) into the impact of brucellosis infection on sheep and goats revealed that infection had no impact on the body weight of infected animals. Nevertheless, the number of offspring was reduced in goats infected with *Brucella melitensis* ([Mahboub et al., 2013](#)). In addition to its significant



effect on performance, brucellosis also has adverse health consequences, as it interferes with crucial biological processes and functions of the liver and kidneys, which is manifested by alterations in the concentration of biochemical parameters in the blood (Mahboub et al., 2013). In a survey on the socioeconomic impact of brucellosis on animal production and reproductive performance in Kenya, it was reported that brucellosis had a significant impact on different aspects such as milk production (54.1%), meat (54.4%), infertility (66.0%), selling cost (64.3%), and treatment cost (62.3%). In Ethiopia, Tadege et al. (2015) assessed the link between brucellosis prevalence and reproductive problems in goats and sheep. The results revealed a significant association between brucellosis seropositivity and the presence of reproductive abnormalities such as abortion, retained fetal membranes, and the birth of weak offspring. Furthermore, pregnant females were more vulnerable to brucellosis infection due to the growth stimulation of the bacteria in their reproductive tract. These findings are similar to those of the study by Tea et al. (2020) in the Dalaba prefecture of Guinea, where similar effects of brucellosis were observed in small ruminants.

#### ***Impact of brucellosis on human health***

Human infection can be transmitted by coming into direct contact with vaginal fluids, placental material, and fetuses that have been aborted by infected animals, or by consuming unprocessed milk or dairy products from these animals (Ducrotoy et al., 2014). Edao et al. (2020) found a prevalence of 2.6% of brucellosis among individuals associated with animal production systems in southern Ethiopia. In a study conducted in Chad on brucellosis in nomadic pastoralists and their livestock, a prevalence of 3.5% of this disease in humans was reported (Schelling et al., 2003). These studies also highlighted a significant association between human brucellosis and assistance during parturition and the presence of seropositive animals within a household. Similar observations were reported in Eritrea by Omer et al. (2002). The results of these studies emphasize the importance of implementing preventive and control measures for brucellosis in animals to reduce the transmission of this disease to humans and limit its socioeconomic impact.

#### ***Methods of diagnosing brucellosis in goats***

Diagnosis of brucellosis in small ruminants relies on a range of specific methods. Among these, the Rose Bengal Plate Test (RBPT) is the most commonly used serological tool in tropical Africa (Kamga et al., 2020; Tea et al., 2020; Lokamar et al., 2020). Rose Bengal Plate Test relies on visible agglutination when antibodies directed against *Brucella* antigens react, forming characteristic “dewy” lumps (Kamga et al., 2020; Tea et al., 2020; Lokamar et al., 2020). Enzyme-linked immunosorbent Assay is a serological method also frequently used to detect the presence of specific antibodies produced in response to *Brucella* infection. Enzyme-linked immunosorbent Assay offers advantages in terms of sensitivity and quantification of immune reactions and is used as a confirmatory test for RBPT-positive samples (Kamga et al., 2020; Legesse et al., 2023). The Serum Agglutination Test (SAT) is also a diagnostic method for brucellosis based on the agglutination reaction between *Brucella* antigens and specific antibodies present in serum (Bertu et al., 2010). Molecular PCR is also widely used for its sensitivity and specificity in detecting *Brucella* DNA (Legesse et al., 2023).

#### ***Prevalence of brucellosis in goats across some tropical African countries***

Studies on the prevalence of brucellosis in goats have been carried out in countries such as Burkina Faso, Cameroon, Ethiopia, Ghana, Guinea, Kenya, Nigeria, and Togo. Prevalence rates vary considerably from one region to another, ranging from 1.1% to 36.84% (Table 3).

#### ***Methods of controlling brucellosis in goats***

Different strategies are used to combat brucellosis. It is essential to remove and destroy parturition material, including aborted fetuses and placentas, by incineration to avoid exposure to other animals, such as dogs, farm cats, and children (Ganter, 2015). Daily cleaning of feces maintains hygiene, while regular disinfection, recommended at least three times a year, reduces the incidence of disease within the herd (Burns et al., 2023). The use of vaccines, such as *B. abortus* and *B. melitensis* in cattle, is proving useful in reducing disease prevalence, but their use is recommended as a last resort after other control methods have failed (Burns et al., 2023). The Rev. 1 *B. melitensis* vaccine controls brucellosis in small ruminants, mainly sexually immature females, offering long-lasting protection while potentially causing abortion in pregnant females and excretion of the organism in milk (Tizard, 2021). The Rev.1 vaccine was developed using streptomycin as a selective agent from a virulent field strain of *B. melitensis* (Tizard, 2021). It is administered by subcutaneous injection or deposition in the conjunctiva of lambs and kids aged three to five months. Conjunctival vaccination is generally considered safer than subcutaneous injection (Tizard, 2021). As a general rule, the entire flock should be vaccinated simultaneously at the end of the lambing or lactation season, especially when rearing under extensive conditions (Tizard, 2021).

The World Organization for Animal Health does not recommend antibiotic treatment of animals for brucellosis, underlining the importance of alternative preventive and control measures (Wareth et al., 2021; Burns et al., 2023).

**Table 3.** Seroprevalence of brucellosis in goats of tropical Africa from 2013 to 2022

Countries	Regions	Sample (Number)	Prevalence (%)			Reference
			Diagnostic test			
			SAT	RBPT	iELISA	
Burkina Faso	Bam Province	300			4.3	(Tialla, 2022)
Cameroon	Western, Central, Southern, and Southwestern regions of the southern part of the country	452		1.3	1.1	(Kamga et al., 2020)
Ethiopia	Tigray Regional State in the northern part of the country	495		5.5		(Teklue et al., 2013)
Ghana	Northern, Ashanti, and Greater Accra regions	286		10		(Jarikre et al., 2015)
Guinea	Dalaba Prefecture	90		10.3		(Tea et al., 2020)
Kenya	Baringo County	155		36.84		(Lokamar et al., 2020)
Nigeria	Plateau State	851	5.9	10.1		(Bertu et al., 2010)
Togo	Northern part of the country	221			8.8	(Dean et al., 2013)

RBPT: Rose Bengal Plate Test, SAT: Serum Agglutination Test, iELISA: Indirect enzyme-linked immunosorbent assay

## CONCLUSION

Gastrointestinal parasites are prevalent in goats, resulting in growth retardation and reduced slaughter weight. Sarcoptic mange mites have detrimental effects on goat production and reproduction, and severe infections can even lead to death. Trypanosomiasis causes decreased hematocrit levels, abortions, low birth weight, and high mortality rates in goats. Peste des petits ruminants and Rift Valley fever have high seroprevalence rates in certain endemic regions. Contagious Caprine Pleuropneumonia presents high morbidity and mortality rates, while brucellosis reduces litter size and disrupts vital organs in goats. The present study can be considered for an overall understanding of major goat diseases and their control in tropical Africa.

## DECLARATIONS

### Acknowledgments

The authors would like to express their gratitude to the University of Abomey-Calavi for providing financial support.

### Availability of data and materials

All data are presented in the published manuscript.

### Fundings

This study was funded by the University of Abomey-Calavi through the Competitive Research Fund Phase III program (PFCR3/CS/UAC).

### Authors' contributions

Kétomon Pierre Challaton drafted the first version of the manuscript. Aboudou Habirou Kifouly contributed to data collection. Kadoéito Cyrille Boko, Coovi Guénoilé Akouedegni, Goué Géorcelin Alowanou, Mawulé Sylvie Hounzangbé-Adoté contributed to the critical revision of the manuscript. All authors checked the last draft of the manuscript and confirmed it before submission to the journal.

### Conflict of interests

The authors declare that they have no conflict of interest.

### Ethical considerations

All ethical issues have been checked by the authors, including plagiarism, double submission, and data originality.

## REFERENCES

- Abakar MF, Naré NB, Schelling E, Hattendorf J, Alfaraoukh IO, and Zinsstag J (2014). Seroprevalence of Rift Valley fever, Q fever, and brucellosis in ruminants on the southeastern shore of Lake Chad. *Vector-Borne and Zoonotic Diseases*, 14(10): 757-762. DOI: <https://www.doi.org/10.1089/vbz.2014.1585>
- Abd-Elrahman AH, Khafaga AF, and Abas OM (2020). The first identification of contagious caprine pleuropneumonia (CCPP) in sheep and goats in Egypt: molecular and pathological characterization. *Tropical Animal Health and Production*, 52: 1179-1186. DOI: <https://www.doi.org/10.1007/s11250-019-02116-5>
- Adam AA, Karsany MS, and Adam I (2010). Manifestations of severe Rift Valley fever in Sudan. *International Journal of Infectious Diseases*, 14(2): E179-E180. DOI: <https://www.doi.org/10.1016/j.ijid.2009.03.029>

- Adeyemo P, Léger E, Hollenberg E, Diouf N, Sène M, Webster JP, and Häslér B (2022). Estimating the financial impact of livestock schistosomiasis on traditional subsistence and transhumance farmers keeping cattle, sheep and goats in northern Senegal. *Parasites & Vectors*, 15: 101. DOI: <https://www.doi.org/10.1186/s13071-021-05147-w>
- African Union-Interafrican Bureau for animal resources (2014). Panafrican animal resources yearbook, pp. 164. Available at: <http://repository.auiibar.org/handle/123456789/533>
- Ahaduzzaman MD (2021). Contagious caprine pleuropneumonia (CCPP): A systematic review and meta-analysis of the prevalence in sheep and goats. *Transboundary and Emerging Diseases*, 68(3): 1332-1344. DOI: <https://www.doi.org/10.1111/tbed.13794>
- Akinwale OP, Nock IH, Esievo KA, Edeghere HU, and Olukosi YA (2006). Study on the susceptibility of Sahel goats to experimental *Trypanosoma vivax* infection. *Veterinary Parasitology*, 137(3-4): 210-213. DOI: <https://www.doi.org/10.1016/j.vetpar.2006.01.032>
- Akwongo CJ, Quan M, and Byaruhanga C (2022). Prevalence, Risk Factors for Exposure, and Socioeconomic Impact of Peste Des Petits Ruminants in Karenga District, Karamoja Region, Uganda. *Pathogens*, 11(1): 54. DOI: <https://www.doi.org/10.3390/pathogens11010054>
- Akwuobu CA, Ayling RD, Chah KF, and Oboegbulem SI (2014). Studies into the prevalence of *Mycoplasma* species in small ruminants in Benue State, North-central Nigeria. *Tropical Animal Health and Production*, 46: 1087-1092. DOI: <https://www.doi.org/10.1007/s11250-014-0613-6>
- Akwuobu CA, Chah KF, Oboegbulem SI, and Rabo JS (2016). Pathogenicity of local isolates of *Mycoplasma ovipneumoniae* and *Mycoplasma arginini* strains in experimental West African Dwarf goats. *Journal of Advanced Veterinary and Animal Research*, 3(3): 242-251. DOI: <http://www.doi.org/10.5455/javar.2016.c161>
- Alemnew E, Asfaw T, Demis C, Alemu B, Yitagesu E, Wondifra Y, Ayele F, and Aydefruh D (2022). Evaluation of the immune response of small ruminant flocks to ovine Pasteurellosis and Peste Des Petits Ruminants vaccines in North Shewa, Ethiopia. *Journal of Science and Inclusive Development*, 4(1): 55-69. DOI: <https://www.doi.org/10.20372/jsid/2022-114>
- Alowanou GG, Adenilé AD, Akouédegni GC, Bossou AC, Zinsou FT, Akakpo GCA, Kifouly HA, Rinaldi L, Georg von Samson-Himmelstjerma G, Cringoli G et al. (2021). A comparison of Mini-FLOTAC and McMaster techniques in detecting gastrointestinal parasites in West Africa Dwarf sheep and goats and crossbreed rabbits. *Journal of Applied Animal Research*, 49(1): 30-38. DOI: <https://www.doi.org/10.1080/09712119.2021.1876703>
- Ameh JA, Egwu GO, and Tijjani AN (2000). Mortality in sahelian goats in Nigeria. *Preventive Veterinary Medicine*, 44(1-2): 107-111. DOI: [https://www.doi.org/10.1016/S0167-5877\(99\)00108-7](https://www.doi.org/10.1016/S0167-5877(99)00108-7)
- Andre Z, Adama K, Zakaria B, Herve VS, Aristide T, Hamadou TH, and Gaston BAM (2017). Prevalence of *trypanosoma brucei brucei* and potential *in vitro* trypanocidal activity of aqueous extracts of some medicinal plants in the pastoral area of gaongho in Burkina Faso. *Journal of Animal Health and Production*, 5(3): 107-114. DOI: <http://www.doi.org/10.17582/journal.jahp/2017/5.3.107.114>
- Archer BN, Weyer J, Paweska J, Nkosi D, Leman P, Tint KS, and Blumberg L (2011). Outbreak of Rift Valley fever affecting veterinarians and farmers in South Africa. *South African Medical Journal*, 101(4): 263-266. DOI: <https://hdl.handle.net/10520/EJC67552>
- Armson B, Ekiri AB, Alafiatayo R, and Cook AJ (2020). Small ruminant production in Tanzania, Uganda, and Ethiopia: a systematic review of constraints and potential solutions. *Veterinary Sciences*, 8(1): 5. DOI: <https://www.doi.org/10.3390/vetsci8010005>
- Arsenopoulos KV, Fthenakis GC, Katsarou EI, and Papadopoulos E (2021). Haemonchosis: A challenging parasitic infection of sheep and goats. *Animals*, 11(2): 363. DOI: <https://www.doi.org/10.3390/ani11020363>
- Arsevka E, Lancelot R, El Mamy B, and Cetre-Sossah C (2016). Situation épidémiologique de la fièvre de la Vallée du Rift en Afrique de l'Ouest et du Nord [Epidemiological situation of Rift Valley fever in West and North Africa]. *Bulletin Epidémiologique, Santé Animale et Alimentation*, 74: 25-29. Available at: <https://agritrop.cirad.fr/580830/1/BEP-mg-BE74-art6.pdf>
- Arul N, Arumugam AP, Padmanaban V, Narayanan D, and Karuppaiah R (2023). Successful Management of Concurrent Scabies and Dermatophytosis in a Chippiparai Pup. *Small Animal Advances*, 2(2): 29-32. DOI: <https://doi.org/10.58803/saa.v2i2.11>
- Asmare K, Abebe R, Sheferaw D, Krontveit RI, and Barbara W (2016a). Mange mite infestation in small ruminants in Ethiopia: Systematic review and meta-analysis. *Veterinary Parasitology*, 218: 73-81. DOI: <https://www.doi.org/10.1016/j.vetpar.2016.01.017>
- Asmare K, Abayneh T, Mekuria S, Ayelet G, Sibhat B, Skjerve E, Szonyi B, and Wieland B (2016b). A meta-analysis of contagious caprine pleuropneumonia (CCPP) in Ethiopia. *Acta Tropica*, 158: 231-239. DOI: <https://www.doi.org/10.1016/j.actatropica.2016.02.023>
- Asuku SO, Bamanga UM, Ali RI, Alkali IM, Ibrahim, A, Abba A, Mustapha AR, Bukar MM, and Waziri MA (2022). Abortions in peste des petits ruminants (PPR) infected flocks in Yola North of Adamawa and Askira Uba areas of Borno States, Nigeria. *Journal of Sustainable Veterinary & Allied Sciences*, 3(2): 114-118. DOI: <http://www.doi.org/10.54328/covm.josvas.2022.084>
- Atim SA, Ayebazibwe C, Mwiine FN, Erume J, and Tweyongyere R (2016). A survey for contagious caprine pleuropneumonia in Agago and Otuke districts in Northern Uganda. *Open Journal of Veterinary Medicine*, 6(1): 62773. DOI: <http://www.doi.org/10.4236/ojvm.2016.61002>
- Attindehou S, Salifou S, Biaou CF, Gbati OB, Adamou-Nrsquo M, and Pangui LJ (2012). Epidemiology of haemonchosis in sheep and goats in Benin. *African Journal of Parasitology Research*, 7(1): 1-5. Available at: <https://www.internationalscholarsjournals.com/articles/epidemiology-of-haemonchosis-in-sheep-and-goats-in-benin.pdf>
- Barda B, Zepherine H, Rinaldi L, Cringoli G, Burioni R, Clementi M, and Albonico M (2013). Mini-FLOTAC and Kato-Katz: helminth eggs watching on the shore of Lake Victoria. *Parasites & Vectors*, 6: 220. DOI: <https://www.doi.org/10.1186/1756-3305-6-220>
- Baron MD, Diallo A, Lancelot R, and Libeau G (2016). Peste des petits ruminants virus. *Advances in Virus Research*, 95: 1-42. DOI: <https://www.doi.org/10.1016/bs.aivir.2016.02.001>
- Barry AM, Pandey VS, Bah S, and Dorny P (2002). Etude épidémiologique des helminthes gastro-intestinaux des caprins en Moyenne Guinée [Epidemiological study of gastrointestinal helminths of goats in Middle Guinea]. *Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux*, 55(2): 99-104. DOI: <https://www.doi.org/10.19182/remvt.9839>
- Bastiaensen P, Dorny P, Batawui K, Boukaya A, Napala A, and Hendrickx G (2003). Parasitisme des petits ruminants dans la zone périurbaine de Sokodé, Togo. II. Caprins [Parasitism of small ruminants in the peri-urban area of Sokodé, Togo. II. Goats]. *Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux*, 56(1-2): 51-56. DOI: <https://www.doi.org/10.19182/remvt.9875>
- Benti E, Sori T, Degu T, and Fesseha H (2020). Manage mites' infestation in small ruminants in Ethiopia-review. *World Applied Sciences Journal*, 38(5): 395-403. DOI: <https://www.doi.org/10.5829/idosi.wasj.2020.395.403>
- Bertu WJ, Ajogi I, Bale JOO, Kwaga JKP, and Ocholi RA (2010). Sero-epidemiology of brucellosis in small ruminants in Plateau State, Nigeria. *African Journal of Microbiology Research*, 4(19): 1935-1938. Available at: <https://academicjournals.org/journal/AJMR/article-full-text-pdf/7ECC43914313.pdf>
- Blomström AL, Scharin I, Stenberg H, Figueiredo J, Nhambirre O, Abilio A, Berg M, and Fafetine J (2016). Seroprevalence of Rift Valley fever virus in sheep and goats in Zambézia. *Mozambique Infection Ecology & Epidemiology*, 6(1): 31344. DOI: <https://www.doi.org/10.3402/iee.v6.31343>
- Boussini H, Lamien CE, Nacoulma OG, Kaboré A, Poda G, and Viljoen G (2014). Prevalence of Rift Valley fever in domestic ruminants in the central and northern regions of Burkina Faso. *Scientific and technical review - International Office of Epizootics*, 33(3): 893-901. Available at: <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=e904cbee172f5f4b58fa21aaf444671ded10eae7>

- Burns RJL, Le KK, Siengsanun-Lamont J, and Blacksell S D (2023). A review of coxiellosis (Q fever) and brucellosis in goats and humans: Implications for disease control in smallholder farming systems in Southeast Asia. *One Health*, 16: 100568. DOI: <https://www.doi.org/10.1016/j.onehlt.2023.100568>
- Chalchisa N, Muktar Y, and Keffale M (2015). Prevalence of gastrointestinal strongyles of small ruminants in and around Haramaya, Ethiopia. *Middle-East Journal of Scientific Research*, 23: 2833-2840.
- Challaton KP, Akouedegni CG, Boko KC, Alowanou GG, Houndonougbo PV, Kifouly AH, and Hounzangbe-Adote MS (2023). Evaluation of the gastrointestinal parasite burden of goats in traditional breeding in Benin. *Journal of Animal Health and Production*, 11(2): 144-154. DOI: <http://www.doi.org/10.17582/journal.jahp/2023/11.2.144.154>
- Chevalier V, Thiongane Y, and Lancelot R (2009). Endemic transmission of Rift Valley fever in Senegal. *Transboundary and Emerging Diseases*, 56(9-10): 372-374. DOI: <https://www.doi.org/10.1111/j.1865-1682.2009.01083.x>
- Chuka A D, Precious OZ and Chinwe C (2020). Diagnosis of mange in West African Dwarf (WAD) and Red Sokoto (RS) goats. *Comparative Clinical Pathology*, 29(4): 799-805. DOI: <https://www.doi.org/10.1007/s00580-020-03116-w>
- Cringoli G, Maurelli MP, Levecke B, Bosco A, Vercruyssen J, Utzinger J, and Rinaldi L (2017). The Mini-FLOTAC technique for the diagnosis of helminth and protozoan infections in humans and animals. *Nature Protocols*, 12(9): 1723-1732. DOI: <https://www.doi.org/10.1038/nprot.2017.067>
- Cringoli G, Rinaldi L, Maurelli MP, and Utzinger J (2010). FLOTAC: New multivalent techniques for qualitative and quantitative copromicroscopic diagnosis of parasites in animals and humans. *Nature Protocols*, 5(3): 503-515. DOI: <https://www.doi.org/10.1038/nprot.2009.235>
- Dahmani A, Khelifi-Touhami NA, Khelifi-Touhami MKT, and Ouchene N (2022). Brucellosis: A retrospective sero-epidemiological study. *Journal of Veterinary Physiology and Pathology*, 1(2): 43-48. DOI: <https://www.doi.org/10.58803/jvpp.v1i2.1>
- Dassou H, Ogné C, Yédomonhan H, Adomou A, Tossou M, Doungnon J, and Akoègninou A (2014). Diversité, usages vétérinaires et vulnérabilité des plantes médicinales au Nord-Bénin [Diversity, veterinary uses and vulnerability of medicinal plants in North Benin]. *International Journal of Biological and Chemical Sciences*, 8(1): 189-210. DOI: <http://www.doi.org/10.4314/ijbcs.v8i1.18>
- Davou KP, Dogo GA, Tanko J, Bialla M, and Kogi CA (2017). Epidemiology of ectoparasites infestation in Jos North, Plateau State, Nigeria. *Saudi Journal of Medical and Pharmaceutical Sciences*, 3: 206-210. DOI: <http://www.doi.org/10.21276/sjmps.2017.3.3.15>
- Dean AS, Bonfoh B, Kulo AE, Boukaya GA, Amidou M, Hattendorf J, Pilo P, and Schelling E (2013). Epidemiology of brucellosis and Q fever in linked human and animal populations in northern Togo. *PLoS ONE*, 8(8): e71501. DOI: <https://www.doi.org/10.1371/journal.pone.0071501>
- Dedieu B, Aubin J, Duteurtre G, Alexandre G, Vayssières J, Bommel P, Faye B, Mahieu M, Fanchone A, and Ickowicz A (2011). Conception et évaluation de systèmes d'élevage durables en régions chaudes, à l'échelle de l'exploitation [Design and evaluation of sustainable breeding systems in hot regions, at farm level]. *INRA Productions Animales*, 24: 113-128. Available at: <https://hal.science/hal-01189570/document>
- Devadharshini J, Mathivathani C, and Sankar Das S (2022). Gastrointestinal parasites of non-descript goats (*Capra hircus*) in Semi-Arid Zone of Southern India. *Small Animal Advances*, 1(1): 22-26. DOI: <https://www.doi.org/10.58803/saa.v1i1.5>
- Diallo A (2006). Control of peste des petits ruminants and poverty alleviation?. *Journal of Veterinary Medicine*, 53(s1): 11-13. DOI: <https://www.doi.org/10.1111/j.1439-0450.2006.01012.x>
- Dias de Castro LL, Abrahão CL, Buzatti A, Molento MB, Bastianetto E, Rodrigues DS, Lopes LB, Xavier Silva M, de Freitas MG, Conde MH et al. (2017). Comparison of McMaster and Mini-FLOTAC fecal egg counting techniques in cattle and horses. *Veterinary Parasitology: Regional Studies and Reports*, 10: 132-135. DOI: <https://www.doi.org/10.1016/j.vprsr.2017.10.003>
- Dione MM, Séry A, Sidibé CAK, Wieland B, and Fall A (2022). Exposure to multiple pathogens-serological evidence for Rift Valley fever virus, *Coxiella burnetii*, Bluetongue virus, and *Brucella* spp. in cattle, sheep, and goat in Mali. *PLoS Neglected Tropical Diseases*. 16(4): e0010342. DOI: <https://www.doi.org/10.1371/journal.pntd.0010342>
- Doutchi M, Ali AA, O Maidagi IA, Mohamed AO, Sibongwere D, Danbouzoua N, Kinda M, Kojan R, Shepherd S, and Ouattara A (2017). Aspects Épidémiologiques, Cliniques et Évolutifs des Cas Compliqués de Fièvre de la Vallée du Rift au District Sanitaire de Tchintabaraden (Niger) [Epidemiological, clinical and evolutionary aspects of complicated cases of Rift Valley Fever in the Tchintabaraden Health District (Niger)]. *Health Sciences and Disease*, 18(2): 3-7. Available at: [http://hsd-fmsb.org/index.php/hsd/article/view/818/pdf\\_424](http://hsd-fmsb.org/index.php/hsd/article/view/818/pdf_424)
- Ducrottoy MJ, Bertu WJ, Ocholi RA, Gusi AM, Bryssinckx W, Welburn S, and Moriyon I (2014). Brucellosis as an emerging threat in developing economies: lessons from Nigeria. *PLOS Neglected Tropical Diseases*, 8(7): e3008. DOI: <https://www.doi.org/10.1371/journal.pntd.0003008>
- Edao BM, Ameni G, Assefa Z, Berg S, Whatmore AM, and Wood JL (2020). Brucellosis in ruminants and pastoralists in Borena, Southern Ethiopia. *PLoS Neglected Tropical Diseases*, 14(7): e0008461. DOI: <https://www.doi.org/10.1371/journal.pntd.0008461>
- Egwu G, Adamu M, Mshelia G, and Bukar-Kolo Y (2012). Isolates of *Mycoplasma mycoides* subspecies *mycoides* (SC) in small ruminants in Sahel zone of Nigeria and its implications on disease control. *African Journal of Biotechnology*, 11(23): 6396-6401. DOI: <https://www.doi.org/10.5897/AJB12.001>
- El-Yuguda AD, Saka SB, Ambali AG, and Egwu GO (2013). Seroprevalence of peste des petits ruminants among domestic small and large ruminants in the semi-arid region of North-eastern Nigeria. *Veterinary World*, 6(10): 807-8011. DOI: <https://www.doi.org/10.14202/vetworld.2013.807-811>
- Ezebuio O, Abenga J, and Ekejindu G (2009). The prevalence of trypanosome infection in trade cattle, goats and sheep slaughtered at the Kaduna abattoir. *African Journal of Clinical and Experimental Microbiology*, 10(1): 15-25. DOI: <https://www.doi.org/10.4314/ajcem.v10i1.7503>
- Faihun AML, Azando EVB, Attakpa EY, Akouèdègnè CG, and Hounzangbe-Adoté MS (2017). Etude comparative de la charge parasitaire des petits ruminants et du guib harnaché dans quatre campements riverains à la forêt classée de Wari-Marou au Nord-Est du Bénin [Comparative study of the parasitic load of small ruminants and harnessed guib in four camps bordering the Wari-Marou classified forest in northeastern Benin]. *Tropicacultura*, 35(1): 51-60. Available at: <https://popups.uliege.be/2295-8010/index.php?id=1154&file=1>
- Fajimi AK and Taiwo AA (2005). Herbal remedies in animal parasitic diseases in Nigeria: A review. *African Journal of Biotechnology*, 4(4): 303-307. Available at: <https://www.ajol.info/index.php/ajb/article/view/15099>
- Farougou S, Gagara M, and Mensah GA (2013). Prevalence of peste des petits ruminants in the arid zone in the Republic of Niger. *Onderstepoort Journal of Veterinary Research*, 80(1): 544. DOI: <https://www.doi.org/10.4102/ojvr.v80i1.544>
- Fawzy M and Helmy YA (2019). The one health approach is necessary for the control of Rift Valley fever infections in Egypt: A comprehensive review. *Viruses*, 11(2): 139. DOI: <https://www.doi.org/10.3390/v11020139>
- Faye D, Sulon J, Kane Y, Beckers JF, Leak S, Kaboret Y, and Geerts S (2004). Effects of an experimental Trypanosoma congolense infection on the reproductive performance of West African Dwarf goats. *Theriogenology*, 62(8): 1438-1451. DOI: <https://www.doi.org/10.1016/j.theriogenology.2004.02.007>
- Ganter M (2015). Zoonotic risks from small ruminants. *Veterinary Microbiology*, 181(1-2): 53-65. DOI: <https://www.doi.org/10.1016/j.vetmic.2015.07.015>
- Gari G, Serda B, Negesa, D, Lemma F, and Asgedom H (2017). Serological investigation of Peste Des Petits Ruminants in East Shewa and Arsi Zones, Oromia Region, Ethiopia. *Veterinary medicine international*, 2017: 976907. DOI: <https://www.doi.org/10.1155/2017/9769071>
- Githiori JB, Höglund J, Waller PJ, and Baker RL (2004). Evaluation of anthelmintic properties of some plants used as livestock dewormers against *Haemonchus contortus* infections in sheep. *Parasitology*, 129(2): 245-253. <https://www.doi.org/10.1017/S003182004005566>

- Grech-Angelini DS (2012). Etude de l'effet de la peste des petits ruminants sur la productivité des troupeaux caprins au Sénégal [Study of the effect of Peste des Petits Ruminants on the productivity of goat herds in Senegal]. Master II, Institut Pasteur, Ecole des Hautes Etudes de Santé Publique, Centre National des Arts et Métiers. Pp. 59. Available at: <http://www.secheresse.info/spip.php?article28935>
- Hounzangbe-Adote MS (2001). L'élevage face à la pharmacopée en médecine vétérinaire au sud du Bénin [Livestock farming and veterinary pharmacopoeia in southern Benin]. Bulletin de la Recherche Agronomique du Bénin, 33: 1-9. Available at: [http://www.slire.net/download/1204/hounzangbe-adote\\_bra\\_033\\_2001.pdf](http://www.slire.net/download/1204/hounzangbe-adote_bra_033_2001.pdf)
- Ibrahim M, Schelling E, Zinsstag J, Hattendorf J, Andargie E, and Tschopp R (2021). Seroprevalence of brucellosis, Q-fever and Rift Valley fever in humans and livestock in Somali Region, Ethiopia. PLoS Neglected Tropical Diseases, 15(1): e0008100. DOI: <https://www.doi.org/10.1371/journal.pntd.0008100>
- Intisar KS, Ali YH, Haj M, Sahar MAT, Shaza MM, Baraa AM, Ishag OM, Nouri M, Taha KM, Nada EM et al. (2017). Peste des petits ruminant's infection in domestic ruminants in Sudan. Tropical Animal Health and Production, 49: 747-754. DOI: <https://www.doi.org/10.1007/s11250-017-1254-3>
- Ipungu L, Kayuma M, Ngoy K, Ivudi K, Ilunga N, and Kindiao B (2017). Fréquence des nématodes gastro-intestinaux chez la chèvre élevée dans la commune annexe de la ville de Lubumbashi [Frequency of gastrointestinal nematodes in goats reared in the outlying district of Lubumbashi]. Journal of Animal & Plant Sciences, 33(1): 5243-5248. Available at: <https://m.elewa.org/Journals/wp-content/uploads/2017/06/1.Ipungu.pdf>
- Jaiswal A, Vikrant S, Neha, and Verma A (2015). Insight into trypanosomiasis in animals: various approaches for its diagnosis, treatment and control: A review Asian Journal of Animal Sciences, 9(5): 172-186. DOI: <https://www.doi.org/10.3923/ajas.2015.172.186>
- Jarikre T, Emikpe B, Folitse R, Odom T, Fuseini A, and Shaibu E (2015). Prevalence of brucellosis in small ruminants in three regions of Ghana. Bulgarian Bulgarian Journal of Veterinary Medicine, 18(1): 49-55. DOI: <https://www.doi.org/10.15547/bjvm.828>
- Jebara KB, Cáceres P, Berlingieri F, and Weber-Vintzel L (2012). Ten years' work on the world organisation for animal health (oie) worldwide animal disease notification system. Preventive veterinary medicine. 107(3-4): 149-159. DOI: <https://www.doi.org/10.1016/j.prevetmed.2012.08.008>
- Kama-Kama F, Midiwo J, Nganga J, Maina N, Schiek E, Omosa LK, and Naessens J (2016). Selected ethno-medicinal plants from Kenya with in vitro activity against major African livestock pathogens belonging to the Mycoplasma mycoides cluster. Journal of Ethnopharmacology, 192: 524-534. DOI: <https://www.doi.org/10.1016/j.jep.2016.09.034>
- Kamba RM, Silatsa BA, Farikou O, Kuate JR, and Simo G (2020). Detection of Brucella antibodies in domestic animals of southern Cameroon: Implications for the control of brucellosis. Veterinary Medicine and Science, 6(3): 410-420. DOI: <https://www.doi.org/10.1002/vms3.264>
- Kamissoko B, Sidibé CAK, Niang M, Samaké K, Traoré A, Diakité A, Sangaré O, Diallo A, and Libeau G (2013). Séroprévalence de la peste des petits ruminants au Mali [Seroprevalence of peste des petits ruminants in Mali]. Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux, 66(1): 5-10. Available at: [https://agritrop.cirad.fr/570672/1/document\\_570672.pdf](https://agritrop.cirad.fr/570672/1/document_570672.pdf)
- Kebede MC (2013). Effect of small ruminant ectoparasites in the tanning industry in Ethiopia: A review. Journal of Animal Science Advances, 3: 424-430. Available at: <https://core.ac.uk/download/pdf/199937304.pdf>
- Kihu SM, Gachohi JM, Ndungu EK, Gitao GC, Bebora LC, John NM, Wairire GG, Maingi N, Wahome RG, and Ireri R (2015). Sero-epidemiology of Peste des petits ruminants virus infection in Turkana County, Kenya. BMC Veterinary Research, 11: 87. DOI: <https://www.doi.org/10.1186/s12917-015-0401-1>
- Kindji SL (2006). La Peste des petits ruminants (PPR) et son incidence socio-économique au Nord-Est du Bénin (Départements du Borgou et de l'Alibori) [Peste des Petits Ruminants (PPR) and its socio-economic impact in northeastern Benin (Borgou and Alibori departments)]. Doctoral thesis, Cheikh Anta Diop university of Dakar, 14: 1-144. Available at: <https://beep.ird.fr/greenstone/collect/eismv/index/assoc/TD06-14.dir/TD06-14.pdf>
- Kinimi E, Odongo S, Muyltermans S, Kock R, and Misinzo G (2020). Paradigm shift in the diagnosis of peste des petits ruminants: scoping review. Acta Veterinaria Scandinavica, 62: 17. DOI: <https://www.doi.org/10.1186/s13028-020-0505-x>
- Kipronoh AK, Ombui JN, Kiara HK, Binopal YS, Gitonga E, and Wesonga HO (2016). Prevalence of contagious caprine pleuro-pneumonia in pastoral flocks of goats in the Rift Valley region of Kenya. Tropical Animal Health and Production, 48(1): 151-155. DOI: <https://www.doi.org/10.1007/s11250-015-0934-0>
- Koudandé OD (2006). Lutte contre la gale des chèvres en milieu villageois au Sud-Bénin [Mange treatment of goats in villages environment in Bénin]. Bulletin de la Recherche Agronomique du Bénin, 51: 12-19. Available at: [http://www.slire.net/download/830/articles\\_51\\_n\\_2\\_lutte\\_contre\\_la\\_gale.pdf](http://www.slire.net/download/830/articles_51_n_2_lutte_contre_la_gale.pdf)
- Lancelot R, Zundel E, and Ducrot C (2011). Spécificités de la santé animale en régions chaudes: Le cas des maladies infectieuses majeures en Afrique [Specificities of animal health in hot regions: the case of major infectious diseases in Africa]. INRA Productions Animales, 24(1): 65-76. Available at: [https://eurl-ppr.cirad.fr/FichiersComplementaires/PPR-ENG/res/lancelot\\_2011.pdf](https://eurl-ppr.cirad.fr/FichiersComplementaires/PPR-ENG/res/lancelot_2011.pdf)
- Legesse A, Mekuriaw A, Gelaye E, Abayneh T, Getachew B, Weldemedhin W, Tesgera T, Deresse G, and Birhanu K (2023). Comparative evaluation of RBPT, I-ELISA, and CFT for the diagnosis of brucellosis and PCR detection of Brucella species from Ethiopian sheep, goats, and cattle sera. BMC microbiology, 23: 216. DOI: <https://www.doi.org/10.1186/s12866-023-02962-2>
- Lokamar PN, Kutwah MA, Atieli H, Gumo S, and Ouma C (2020). Socioeconomic impacts of brucellosis on livestock production and reproduction performance in Koibatek and Marigat regions, Baringo County, Kenya. BMC Veterinary Research, 16: 61. Available at: <https://bmcvetres.biomedcentral.com/articles/10.1186/s12917-020-02283-w>
- Lopes FC, de Paiva KAR, Coelho WAC, Nunes FVA, da Silva JB, da Escóssia CdGM., de Macêdo Praça L, Silva JBA, Freitas CIA, and Batista JS (2016). Lactation curve and milk quality of goats experimentally infected with Trypanosoma vivax. Experimental Parasitology, 167: 17-24. DOI: <https://www.doi.org/10.1016/j.exppara.2016.04.006>
- Lotfalizadeh N, Sadr S, Moghaddam S, Saberi Najjar M, Khakshoor A, and Simab PA (2022). The innate immunity defense against gastrointestinal nematodes: Vaccine development. Farm Animal Health and Nutrition, 1(2): 31-38. DOI: <https://www.doi.org/10.58803/fahn.v1i2.10>
- Loum Gazida R, Oyetola WD, Sall B, Gaye AM, Fall M, Moustapha L, and Alamedjji RB (2022). Évaluation du statut immunitaire post vaccinal des petits ruminants dans le cadre de la lutte contre la Peste des Petits Ruminants au Sénégal [Evaluation of the post-vaccination immune status of small ruminants as part of the fight against Peste des Petits Ruminants in Senegal]. Journal of Applied Biosciences, 179: 18762-18772. DOI: <https://www.doi.org/10.35759/JABs.179.8>
- Mafie E, Rupa FH, Takano A, Suzuki K, Maeda K, and Sato H (2018). First record of Trypanosoma dionisii of the T. cruzi clade from the Eastern bent-winged bat (Miniopterus fuliginosus) in the Far East. Parasitology Research, 117(3): 673-680. DOI: <https://www.doi.org/10.1007/s00436-017-5717-2>
- Maganga GD, Abessolo Ndong A., Mikala Okouyi CS, Makiala Mandanda S, N'Dilimabaka N, Pinto A, Agossou E, Cossic B, Akue J-P, and Leroy EM (2017). Serological evidence for the circulation of Rift Valley fever virus in domestic small ruminants in southern Gabon. Vector-Borne and Zoonotic Diseases, 17(6): 443-446. DOI: <https://www.doi.org/10.1089/vbz.2016.2065>
- Maganga GD, Boundenga L, Ologui-Minkue-Edzo EJ, Kombila LB, Mebaley TGN, Kumulungui B, and Mavoungou JF (2020). Frequency and diversity of trypanosomes in sheep and goats from Mongo County in South Gabon, Central Africa. Veterinary World, 13(11): 2502-2507. DOI: <https://www.doi.org/10.14202/vetworld.2020.2502-2507>

- Mahamat O, Doungous T, Kebkiba B, Oumar HA, Oussiguéré A, and Yacoub AH (2018). Seroprevalence, geographical distribution, and risk factors of Peste des petits ruminants in the Republic of Chad. *Journal of Advanced Veterinary and Animal Research*, 5(4): 420–425. DOI: <https://www.doi.org/10.5455/javar.2018.e293>
- Mahboub H, Helal M, Abd Eldaim M, Abd El-Razek M, and Elsfify A (2013). Seroprevalence of abortion causing agents in Egyptian sheep and goat breeds and their effects on the animal's performance. *Journal of Agricultural Science*, 5(9): 92-101. DOI: <http://www.doi.org/10.5539/jas.v5n9p92>
- Malla ME, Payne VK, Cedric Y, Christelle Nadia NA, Leonelle M, and Rosine M (2021). Prevalence of gastrointestinal parasites in sheep and goats of Bui and Donga-Mantung divisions of the North West region of Cameroon. *Asian Journal of Research in Animal and Veterinary Sciences*, 7(4): 1-15. Available at: <http://stmtdigitallib.com/id/eprint/327/1/149-Article%20Text-271-2-10-20220921.pdf>
- Misinzio G, Kgotlele T, Muse EA, Van Doorselaere J, Berg M, and Munir M (2015). Peste des petits ruminants virus lineage II and IV from goats in southern Tanzania during an outbreak in 2011. *British Journal of Virology*, 2(1):1-4. Available at: <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=32a56d2dcbfd9e306f01708393376615c7e4ada>
- Missohou A, Nahimana G, Ayssiwede SB, and Sembene M (2016). Elevage caprin en Afrique de l'Ouest: Une synthèse [Goat breeding in West Africa: A summary]. *Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux*, 69(1): 3-18. DOI: <https://www.doi.org/10.19182/remvt.31167>
- Monau P, Raphaka K, Zvinorova-Chimboza P, and Gondwe, T (2020). Sustainable utilization of indigenous goats in Southern Africa. *Diversity*, 12(1): 20. DOI: <https://www.doi.org/10.3390/d12010020>
- Morrison LJ, Steketeke PC, Tetey MD, and Matthews KR (2023). Pathogenicity and virulence of African trypanosomes: From laboratory models to clinically relevant hosts. *Virulence*, 14(1): 2150445. DOI: <https://www.doi.org/10.1080/21505594.2022.2150445>
- Moumin G, Moussa C, Teshale S, and Gezahegne M (2018). Seroprevalence and risk factors for peste des petits ruminants in sheep and goats in Djibouti. *Revue Scientifique et Technique*, 37(3): 961-969. DOI: <https://www.doi.org/10.20506/37.3.2899>
- Mpofu TJ, Nephawe KA, and Mtileni B (2022). Prevalence and resistance to gastrointestinal parasites in goats: A review, *Veterinary World*, 15(10): 2442-2452. DOI: <https://www.doi.org/10.14202/vetworld.2022.2442-2452>
- O Ng'ayo M, Njiru ZK, Kenya EU, Muluvi GM, Osir EO, and Masiga DK (2005). Detection of trypanosomes in small ruminants and pigs in western Kenya: important reservoirs in the epidemiology of sleeping sickness?. *Kinetoplastid Biology and Disease*, 4: 5. DOI: <https://www.doi.org/10.1186/1475-9292-4-5>
- Muniraju M, Munir M, Parthiban AR, Banyard A, Bao J, Wang Z, Ayeabazibwe C, Ayelet G, El Harrak M, Mahapatra M et al. (2014). Molecular evolution of peste des petits ruminants virus. *Emerging Infectious Diseases*. DOI: <http://www.doi.org/10.3201/eid2012.140684>
- Munyua P, Murithi RM, Wainwright S, Githinji J, Hightower A, Mutonga D, Macharia J, Ithondeka PM, Musaa J, Breiman RF et al. (2011). Rift Valley fever outbreak in livestock in Kenya, 2006–2007. *The American Journal of Tropical Medicine and Hygiene*, 83(2 Suppl): 58-64. DOI: <http://www.doi.org/10.4269/ajtmh.2010.09-0292>
- Nair MRR, Sejian V, Silpa MV, Fonsêca VFC, de Melo Costa CC, Devaraj C, Krishnan G, Bagath M, Nameer PO, and Bhatta R (2021). Goat as the ideal climate-resilient animal model in tropical environment: revisiting advantages over other livestock species. *International Journal of Biometeorology*, 65: 2229-2240. DOI: <https://www.doi.org/10.1007/s00484-021-02179-w>
- Nakouné E, Kamgang B, Berthet N, Manirakiza A, and Kazanji M (2016). Rift Valley fever virus circulating among ruminants, mosquitoes and humans in the Central African Republic. *PLoS Neglected Tropical Diseases*, 10(10): e0005082. DOI: <https://www.doi.org/10.1371/journal.pntd.0005082>
- Nanyingi MO, Muchemi GM, Thumbi SM, Ade F, Onyango CO, Kiama SG, and Bett B (2017). Seroepidemiological survey of Rift Valley fever virus in ruminants in Garissa, Kenya. *Vector-borne and zoonotic diseases*, 17(2): 141-146. DOI: <https://www.doi.org/10.1089/vbz.2016.1988>
- Njeumi F, Bailey D, Soula JJ, Diop B, and Tekola BG (2020). Eradicating the scourge of peste des petits ruminants from the world. *Viruses*, 12(3): 313. DOI: <https://www.doi.org/10.3390/v12030313>
- Njoya A, Awa DN, Tama ACN, Cardinale E, and Mamoudou A (2005). Evaluation d'une stratégie de réduction de la mortalité des petits ruminants en zone soudano-sahélienne du Nord-Cameroun [Evaluation of a strategy to reduce small ruminant mortality in the Sudano-Saharan zone of North Cameroon]. *Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux*, 58(1-2): 89-94. DOI: <https://www.doi.org/10.19182/remvt.9946>
- Noel ML, Scare JA, Bellow JL, and Nielsen MK (2017). Accuracy and precision of mini-FLOTAC and McMaster techniques for determining equine strongyle egg counts. *Journal of Equine Veterinary Science*, 48: 182-187. DOI: <https://www.doi.org/10.1016/j.jevs.2016.09.006>
- Ogundiya AI, Bemji MN, Adebambo OA, Dipeolu MA, Onagbesan OM, James IJ, and Osinowo OA (2012). Prevalence of mange among West African Dwarf sheep and goats and associated haematological and biochemical parameters. *Tropical Animal Health and Production*, 44: 1263-1269. DOI: <https://www.doi.org/10.1007/s11250-011-0067-z>
- Olorunshola ID, Daodu BO, Ajiboye B, Folaranmi EB, Nicholas RA, Adegboye DS, and Peters AR (2020). Seroprevalence of contagious bovine pleuropneumonia and contagious caprine pleuropneumonia in the Middle-Belt of Nigeria. *African Journal of Microbiology Research*, 14(1): 25-31 DOI: <https://www.doi.org/10.5897/AJMR2019.9262>
- Omer M, Assefaw T, Skjerve E, Teklegiorgis T, and Woldehiwet Z (2002). Prevalence of antibodies to *Brucella spp.* and risk factors related to high-risk occupational groups in Eritrea. *Epidemiology & Infection*, 129(1): 85-91. DOI: <https://www.doi.org/10.1017/S0950268802007215>
- Opayele AV, Ndiana LA, Odaibo GN, and Olaleye DO (2019). Serological evidence of Rift Valley fever virus infection in slaughtered ruminants in Nigeria. *Journal of Immunoassay and Immunochemistry*, 40(4): 367-377. DOI: <https://www.doi.org/10.1080/15321819.2019.1609498>
- Organisation Mondiale de la Santé Animale (OIE) (2017). Programme mondial d'éradication de la peste des petits ruminants, Phase quinquennale (2017-2021), pp. 11-88. Available at: <https://www.woah.org/app/uploads/2021/03/fr-gep-ppr-finalweb.pdf>
- Otsyina H, Arthur C, Ayim-Akunnor M, and Obese F (2013). Sero-prevalence of pestes des petits ruminant (PPR) in sheep, goats and cattle in Ghana. *Bulletin of Animal Health and Production in Africa*, 61(3): 473-479. Available at: <https://www.ajol.info/index.php/bahpa/article/view/105283>
- Pepin M, Bouloy M, Bird BH, Kemp A, and Paweska J (2010). Rift Valley fever virus (Bunyaviridae: Phlebovirus): An update on pathogenesis, molecular epidemiology, vectors, diagnostics and prevention. *Veterinary Research*, 41(6): 61. DOI: <https://www.doi.org/10.1051/vetres/2010033>
- Poueme R, Stoek F, Nloga N, Awah-Ndukum J, Rissmann M, Schulz A, Wade A, Kouamo J, Moctar M, Eisenbarth A et al. (2019). Seroprevalence and associated risk factors of Rift Valley Fever in domestic small ruminants in the North Region of Cameroon. *Veterinary Medicine International*, 2019: 8149897. DOI: <https://www.doi.org/10.1155/2019/8149897>
- Rissmann M, Eiden M, El Mamy B, Isselmou K, Doumbia B, Ziegler U, Homeier-Bachmann T, Yahya B, and Groschup M (2017). Serological and genomic evidence of Rift Valley fever virus during inter-epidemic periods in Mauritania. *Epidemiology & Infection*, 145(5): 1058-1068. DOI: <https://www.doi.org/10.1017/S0950268816003022>
- Sado FY, Tchegnana HS, Kamgang B, Djonabaye D, Nakouné E, McCall PJ, Ndip RN, and Wondji CS (2022). Seroprevalence of Rift Valley fever virus in domestic ruminants of various origins in two markets of Yaoundé, Cameroon. *PLoS neglected tropical diseases*. e0010683. DOI: <https://www.doi.org/10.1371/journal.pntd.0010683>
- Sadr S, Simab PA, Kasaei M, Landi MG, Borji H, and Adhami G (2022). Potential of anthelmintic herbal drugs against gastrointestinal nematodes in farm animals: A review. *Farm Animal Health and Nutrition*, 1(1): 26-30. DOI: <https://www.doi.org/10.58803/fahn.v1i1.9>

- Salifou S, Attindéhou S, Salifou CFA, and Pangui LJ (2013). Prevalence and zoonotic aspects of small ruminant mange in the lateritic and waterlogged zones, southern Benin. *Revista Brasileira de Parasitologia Veterinária*, 22(2): 243-247. DOI: <https://www.doi.org/10.1590/S1984-29612013000200043>
- Sargison N and Scott P (2003). Survey of sheep nematode parasite control methods in south-east Scotland. *Veterinary Records*, 152: 51-52. DOI: <https://www.doi.org/10.1136/vr.152.2.51>
- Schelling E, Diguimbaye C, Daoud S, Nicolet J, Boerlin P, Tanner M, and Zinsstag J (2003). Brucellosis and Q-fever seroprevalences of nomadic pastoralists and their livestock in Chad. *Preventive Veterinary Medicine*, 61(4): 279-293. DOI: <https://www.doi.org/10.1016/j.prevetmed.2003.08.004>
- Shyaka A, Ugirabe MA, and Wensman JJ (2021). Serological evidence of exposure to peste des petits ruminants in small ruminants in Rwanda. *Frontiers in Veterinary Science*, 8: 651978. DOI: <https://www.doi.org/10.3389/fvets.2021.651978>
- Simo G, Herder S, Njikou F, Asonganyi T, Tilley A, and Cuny G (2005). *Trypanosoma brucei* s.l: characterisation of stocks from Central Africa by PCR analysis of mobile genetic elements. *Experimental Parasitology*, 110(4): 353-362. DOI: <https://www.doi.org/10.1016/j.exppara.2005.04.001>
- Sindato C, Pfeiffer DU, Karimuribo ED, Mboera LE, Rweyemamu MM, and Paweska JT (2015). A spatial analysis of Rift Valley fever virus seropositivity in domestic ruminants in Tanzania. *PloS ONE*, 10(7): e0131873. DOI: <https://www.doi.org/10.1371/journal.pone.0131873>
- Sow A, Ouattara L, Compaoré Z, Doukom BR, Paré M, Poda G, and Nyambré J (2008). Prévalence sérologique de la peste des petits ruminants dans la province du Soum au nord du Burkina Faso [Serological prevalence of peste des petits ruminants in Soum province, northern Burkina Faso]. *Revue D'élevage et de Médecine Vétérinaire des Pays Tropicaux*, 61(1): 5-9. DOI: <https://www.doi.org/10.19182/remvt.10012>
- Sow A, Ouédraogo S, Sidibé I, Kalandi M, Zabré ZM, and Sawadogo GJ (2014). Enquête de base parasitologique de la trypanosomose animale dans trois zones agropastorales du Burkina Faso [Parasitological baseline survey of animal trypanosomosis in three agropastoral zones of Burkina Faso]. *Bulletin Animal Health and Production*, 62: 241-250. Available at: <http://edocs.maseno.ac.ke/bitstream/handle/123456789/2829/2.pdf?sequence=1&isAllowed=y#page=44>
- Swai ES, Kapaga A, Kivaria F, Tinuga D, Joshua G, and Sanka P (2009). Prevalence and distribution of peste des petits ruminants virus antibodies in various districts of Tanzania. *Veterinary Research Communications*, 33(8): 927-936. DOI: <https://www.doi.org/10.1007/s11259-009-9311-7>
- Tadeg WM, Gudeta FR, Mekonen TY, Asfaw YT, Birru AL, and Reda AA (2015). Seroprevalence of small ruminant brucellosis and its effect on reproduction at Tellalak District of Afar region, Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 7(4): 111-116. DOI: <https://www.doi.org/10.5897/JVMAH2014.0287>
- Tariq M, Khan MS, Mubashir M, Safdar M, Özaslın M, Farooq Z, Rizwan MA, Kaleem M, Ullah Q, Siddique F et al. (2022). Review on Trypanosomiasis and their prevalence in ruminants. *Zeugma Biological Science*, 3(2): 12-31. Available at: <https://dergipark.org.tr/en/download/article-file/2320070>
- Tassou AW (2009). Infestation des ruminants domestiques par les acariens et insectes dans le Nord-Bénin : impact et connaissance paysanne de lutte [Infestation of domestic ruminants by mites and insects in northern Benin: impact and farmers' knowledge of control methods]. Doctoral thesis, Cheikh Anta Diop university of Dakar, pp. 117. Available at: <https://beep.ird.fr/collect/eismv/index/assoc/TD09-12.dir/TD09-12.pdf>
- Tchetan E, Olounlade AP, Houehanou TD, Azando EVB, Kaneho JA, Houinato MRB, and Gbaguidi FA (2021). Ethnoveterinary knowledge of sheep and goat farmers in Benin (West Africa): effect of socioeconomic and environmental factors. *Heliyon*, 7(7): e07656. DOI: <https://www.doi.org/10.1016/j.heliyon.2021.e07656>
- Tea MA, Soromou LW, Beavogui S, and Keyra M (2020). Epidemiological survey AND effects of Brucellosis in small ruminants in selected pastoral lands of dalaba, Guinea. *Journal of Global Biosciences*, 9(6): 7560-7572. Available at: <https://www.mutagens.co.in/jgb/vol.09/06/090610.pdf>
- Tekle T, Tolosa T, Tuli G, Beyene B, and Hailu B (2013). Sero-prevalence and risk factors study of brucellosis in small ruminants in Southern Zone of Tigray Region, Northern Ethiopia. *Tropical Animal Health and Production*, 45(8): 1809-1815. DOI: <https://www.doi.org/10.1007/s11250-013-0439-7>
- Terefe D, Demissie D, Beyene D, and Haile S (2012). A prevalence study of internal parasites infecting Boer goats at Adami Tulu agricultural research center, Ethiopia. *Journal of Veterinary Medicine and Animal Health*, 4(2): 12-16. Available at: [https://academicjournals.org/article/article1379668204\\_Terefe%20et%20al%20\(1\).pdf](https://academicjournals.org/article/article1379668204_Terefe%20et%20al%20(1).pdf)
- Tialla D (2022). The first study on seroprevalence and risk factors for zoonotic transmission of ovine and caprine brucellosis in the Province of Bam, Burkina Faso. *Veterinary World*, 15(2): 262-267. DOI: <https://www.doi.org/10.14202/vetworld.2022.262-267>
- Tizard IR (2021). Sheep and goat vaccines. *Vaccines for Veterinarians*, 2021: 215-224e1. DOI: <https://www.doi.org/10.1016/B978-0-323-68299-2.00026-5>
- Torsson E, Berg M, Misinzo G, Herbe I, Kgotlele T, Päärni M, Roos N, Blomström AL, Ståhl K, and Johansson Wensman J (2017). Seroprevalence and risk factors for peste des petits ruminants and selected differential diagnosis in sheep and goats in Tanzania. *Infection Ecology & Epidemiology*, 7(1): 1368336. DOI: <https://www.doi.org/10.1080/20008686.2017.1368336>
- Troupin C, Ellis I, Doukouré B, Camara A, Keita M, Kagbadouno M, Bart JM, Diallo M, Lacôte S, Marianneau P et al. (2022). Seroprevalence of brucellosis, Q fever and Rift Valley fever in domestic ruminants in Guinea in 2017–2019. *BMC Veterinary Research*, 18(1): 64. DOI: <https://www.doi.org/10.1186/s12917-022-03159-x>
- Tshilenge GM, Mulumba MLK, Misinzo G, Noad R, and Dundon WG (2019). Rift Valley fever virus in small ruminants in the Democratic Republic of the Congo. *Onderstepoort Journal of Veterinary Research*, 86(1): a1737. DOI: <https://www.doi.org/10.4102/ojvr.v86i1.1737>
- Van Wyk JA and Mayhew E (2013). Morphological identification of parasitic nematode infective larvae of small ruminants and cattle: A practical lab guide. *Onderstepoort Journal of Veterinary Research*, 80(1): 539. DOI: <https://www.doi.org/10.4102/ojvr.v80i1.539>
- Wanjama J, Aboge G, Muneri C, and Nanyingi M (2022). Sero-Epidemiological Survey of Rift Valley Fever Virus in Ruminants in Nyandarua County, Kenya. *East African Agricultural and Forestry Journal*, 86(1-2): 11-11. Available at: [file:///C:/Users/PC/Downloads/555-Article%20Text-943-1-10-20220826%20\(2\).pdf](file:///C:/Users/PC/Downloads/555-Article%20Text-943-1-10-20220826%20(2).pdf)
- Wareth G, El-Diasty M, Abdel-Hamid NH, Holzer K, Hamdy MER, Moustafa S, Shahein MA, Melzer F, Beyer W, Pletzw MW et al. (2021). Molecular characterization and antimicrobial susceptibility testing of clinical and non-clinical *Brucella melitensis* and *Brucella abortus* isolates from Egypt. *One Health*, 13: 100255. DOI: <https://www.doi.org/10.1016/j.onehlt.2021.100255>
- Waret-Szkuta A, Roger F, Chavernac D, Yigezu L, Libeau G, Pfeiffer DU, and Guitián J (2008). Peste des Petits Ruminants (PPR) in Ethiopia: Analysis of a national serological survey. *BMC Veterinary Research*, 4: 34. DOI: <https://www.doi.org/10.1186/1746-6148-4-34>
- Woma TY, Quan M, Ekong PS, Bwala DG, Ibu JO, Ta'ama L, Dyek DY, Saleh L, Shamaki D, Kalla D et al. (2016). Serosurvey of peste des petits ruminants virus in small ruminants from different agro-ecological zones of Nigeria: Original research. *Onderstepoort Journal of Veterinary Research*, 83(1): 1-9. Available at: <https://journals.co.za/doi/epdf/10.4102/ojvr.v83i1.1035>
- World Health Organization (2007). Outbreaks of Rift Valley fever in Kenya, Somalia and United Republic of Tanzania, December 2006–April 2007. *Weekly Epidemiological Record= Relevé épidémiologique hebdomadaire*, 82(20): 169-178. Available at: [https://apps.who.int/iris/bitstream/handle/10665/240937/WER8220\\_169-178.PDF](https://apps.who.int/iris/bitstream/handle/10665/240937/WER8220_169-178.PDF)
- Yasine A, Kumsa B, Hailu Y, and Ayana D (2015). Mites of sheep and goats in Oromia Zone of Amhara Region, North Eastern Ethiopia: species, prevalence and farmers awareness. *BMC Veterinary Research*, 11: 122. DOI: <https://www.doi.org/10.1186/s12917-015-0433-6>

- Yatoo MI, Parray OR, Bashir ST, Muheet, Bhat R, Gopalakrishnan A, Karthik K, Dhama K, and Singh SV (2019). Contagious caprine pleuropneumonia—a comprehensive review. *The Veterinary Quarterly*, 39(1): 1-25. DOI: <https://www.doi.org/10.1080/01652176.2019.1580826>
- Ye A (2012). Contribution à la connaissance des pathologies des petits ruminants dans trois communes du Houet (Dandé, Padema et Satiri) [Contribution to knowledge of small ruminant pathologies in three Houet communes (Dandé, Padema and Satiri)]. Mémoire de fin d'études. IDR/UPB, Option Elevage., Burkina Faso, pp. 63.
- Zajac AM and Garza J (2020). Biology, epidemiology, and control of gastrointestinal nematodes of small ruminants. *Veterinary Clinics of North America: Food Animal Practice*, 36(1): 73-87. DOI: <https://www.doi.org/10.1016/j.cvfa.2019.12.005>
- Zewdie G, Derese G, Getachew B, Belay H, and Akalu M (2021). Review of sheep and goat pox disease: Current updates on epidemiology, diagnosis, prevention and control measures in Ethiopia. *Animal Diseases*, 1: 28. DOI: <https://www.doi.org/10.1186/s44149-021-00028-2>

**Publisher's note:** [Scienceline Publication](#) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access:** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023